

- Feynman, R. P., and Hibbs, A. R. (1965). *Quantum Mechanics and Path Integrals*, McGraw-Hill, New York, pp. 96ff.
- Fisher, R. S. (1973). *Statistical Methods and Scientific Inference*, 3rd ed., Hafner Press, New York, pp. 153ff.
- Gibbs, J. W. (1902). *Elementary Principles in Statistical Mechanics*, Yale University Press, New Haven.
- Khinchin, A. I. (1949). *Mathematical Foundations of Statistical Mechanics*, Dover, New York.
- Klein, M. J. (1985). *Paul Ehrenfest: The Making of a Theoretical Physicist*, 3rd ed., North-Holland, Amsterdam.
- Kolmogorov, A. N. (1956). *Foundations of the Theory of Probability*, 2nd English ed., Chelsea, New York, p. 42.
- Koopman, D. O. (1936). *Transactions of the American Mathematical Society*, **39**, 399.
- Kullback, S. (1959). *Information Theory and Statistics*, Dover, New York.
- Kullback, S., and Khairat, M. A. (1966). *Annals of Mathematical Statistics*, **37**, 279.
- Kullback, S., and Leibler, R. A. (1951). *Annals of Mathematical Statistics*, **22**, 79.
- Landau, L. D., and Lifshitz, E. M. (1969). *Statistical Physics*, 2nd ed., Pergamon Press, Oxford.
- Lavenda, B. H. (1987). The implication of Bayes' theorem in statistical mechanics, submitted for publication.
- Lavenda, B. H., and Scherer, C. (1987a). *Rivista del Nuovo Cimento*, in press.
- Lavenda, B. H., and Scherer, C. (1987b). The statistical inference approach to generalized thermodynamics, submitted for publication.
- Le Cam, L. (1953). *University of California Publications in Statistics*, Vol. 1, p. 277.
- Lorentz, H. A. (1916). *Les Théories Statistique en Thermodynamique*, Teubner, Leipzig, pp. 10 and 16.
- Mandelbrot, B. (1956). *IRE Transactions on Information Theory*, **IT-2**, 190.
- Mandelbrot, B. (1962). *Annals of Mathematical Statistics*, **33**, 1021.
- Mandelbrot, B. (1964). *Journal of Mathematical Physics* **5**, 164.
- Neyman, J., and Pearson, E. S. (1933). *Proceedings of the Cambridge Philosophical Society*, **29**, 492.
- Perrin, F. (1939). *Mécanique Statistique Quantique*, Gauthier-Villars, Paris.
- Pitman, E. J. G. (1936). *Proceedings of the Cambridge Philosophical Society*, **32**, 567.
- Rao, C. R. (1945). *Bulletin of the Calcutta Mathematical Society*, **37**, 81.
- Savage, L. (1962). In *The Foundations of Statistical Inference*, Methuen, London.
- Szilard, L. (1925). *Zeitschrift für Physik*, **32**, 753.
- Wald, A. (1947). *Sequential Analysis*, Wiley, New York, Chapter 3.

Prolongation Structure and Painlevé Property of the Gürses–Nutku Equations

Ayşe Kalkanlı¹

Received November 27, 1986

It is shown that the Gürses–Nutku equations have a finite prolongation algebra for any value of the parameter K . The Painlevé property of these equations is also examined.

A generalization of the KdV equation found by Gürses and Nutku (1981) is given by

$$u_t + 6uu_x + u_{xxx} - \lambda_x u_{xx} = 0 \quad (1a)$$

$$\lambda_t + 2u\lambda_x + 2Ku_x = 0 \quad (1b)$$

where K is an arbitrary constant. These equations arise as the embedding equations of a two-dimensional surface into a flat, three-dimensional space. Gürses and Nutku discussed the equivalence between the two-dimensional integrable systems and surface theory at the metric level. In a recent work, Chowdhury and Paul (1985) studied the prolongation structure of equations (1) without being aware of Gürses and Nutku (1981). They showed that the prolongation algebra closes for $K = -1, 1, 2$. In this work we restudy the prolongation algebra of Gürses–Nutku equations and show that it closes for any value of K . We also show that a nontrivial Bäcklund transformation cannot be found by prolongation techniques of Wahlquist and Estabrook (1975). We reduced the partial differential equations (1) to ordinary differential equations by transformation of variables and applied the Painlevé test of Ablowitz *et al.* (1980). We also applied the Painlevé test for partial differential equations introduced by Weiss *et al.* (1983) to the Gürses–Nutku equations. In both cases these equations fail when $K \neq 0$. Therefore they are not of P -type in their present form.

¹Department of Physics, Middle East Technical University, 06531, Ankara, Turkey.

Introducing the notation

$$u_x = p, \quad p_x = q \quad (2)$$

we can write equations (1) as

$$u_t + 6up + q_x - \lambda_x q = 0 \quad (3a)$$

$$\lambda_t + 2u\lambda_x + 2Kp = 0 \quad (3b)$$

These first-order partial differential equations can be associated with the set of four 2-forms,

$$\begin{aligned} \alpha_1 &= du \wedge dt - p dx \wedge dt \\ \alpha_2 &= dp \wedge dt - q dx \wedge dt \\ \alpha_3 &= dq \wedge dt + 6up dx \wedge dt - q d\lambda \wedge dt - du \wedge dx \\ \alpha_4 &= -d\lambda \wedge dx + 2u d\lambda \wedge dt + 2Kp dx \wedge dt \end{aligned} \quad (4)$$

The set $\{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$ constitutes a closed ideal. That is,

$$d\alpha_a = \sum_{b=1}^4 f_{ab} \alpha_b, \quad a, b = 1, \dots, 4 \quad (5)$$

where f_{ab} are some 1-forms. We now seek a set of 1-forms

$$W_k = dy_k + F_k dx + G_k dt, \quad k = 1, \dots, n \quad (6)$$

with $F_k(u, p, q, \lambda, y_k)$ and $G_k(u, p, q, \lambda, y_k)$ having the property that the prolonged ideal $\{\alpha_1, \alpha_2, \alpha_3, \alpha_4, w_1, \dots, w_n\}$ is closed, that is,

$$dw_k = \sum_{a=1}^4 g_{ka} \alpha_a + \sum_{i=1}^n n_k^i \wedge w^i \quad (7)$$

where n is the number of prolongation variables, and g_{ka} and n_k^i are some sets of 0-forms and 1-forms, respectively. This requirement gives the set of partial differential equations for F_k and G_k :

$$\begin{aligned} F_p &= F_q = 0, \quad F_u + G_q = 0 \\ G_\lambda + 2uF_\lambda - qF_u &= 0 \end{aligned} \quad (8)$$

$$pG_u + qG_p + 6upF_u + 2KpF_\lambda + [F, G] = 0$$

where

$$[F, G] = \frac{\partial F}{\partial y^i} G^i - \frac{\partial G}{\partial y^i} F^i$$

The solutions of equations (8) are

$$\begin{aligned} F &= u\beta_0 e^{-\lambda} + \gamma_0 e^\lambda \\ G &= -q\beta_0 e^{-\lambda} + p\delta_0 - 2u(u\beta_0 e^{-\lambda} + \gamma_0 e^\lambda) + \xi_0 \end{aligned} \quad (9)$$

where $\beta_0, \gamma_0, \delta_0$, and ξ_0 are independent of u, p, q , and λ and satisfy the commutator relations

$$\begin{aligned} [\gamma_0, \beta_0] &= \delta_0 \\ [\delta_0, \gamma_0] &= 2\gamma_0(K-1) \\ [\delta_0, \beta_0] &= 2\beta_0(1-K) \\ [\xi_0, \gamma_0] &= [\xi_0, \beta_0] = [\xi_0, \delta_0] = 0 \end{aligned} \quad (10)$$

The above algebra is finite-dimensional and no “closing off” is needed. With the choice $\xi_0 = 0$, which does not alter our conclusion, equations (9) and (10) reduce to the results given by Chowdhury and Paul (1985) for the special values $K = -1, 1, 2$, provided F and G in (9) are expressed in terms of

$$p = Ke^{\lambda/K} \quad (11)$$

Choosing $\gamma_0 = (1-K)X_1$, $\beta_0 = -X_2$, and $\delta_0 = (K-1)X_0$, one can write F and G in (9) in terms of the generators of $SL(2, R)$ algebra, satisfying the commutation relations

$$[X_0, X_1] = 2X_1, \quad [X_0, X_2] = -2X_2, \quad [X_1, X_2] = X_0 \quad (12)$$

Using the 2×2 matrix representation of these generators, we can construct an $SL(2, R)$ -valued connection 1-form

$$\Gamma = \begin{pmatrix} \Theta_0 & \Theta_1 \\ \Theta_2 & -\Theta_0 \end{pmatrix} \quad (13)$$

where

$$\begin{aligned} \Theta_0 &= (K-1)u_x dt \\ \Theta_1 &= (1-K)e^\lambda(dx - 2u dt) \\ \Theta_2 &= -ue^{-\lambda} dx + e^{-\lambda}(2u^2 + u_{xx}) dt \end{aligned} \quad (14)$$

This connection defines a linear equation $d\Psi = -\Gamma\Psi$, where Ψ is a column vector with components Ψ_1 and Ψ_2 . This associated linear equation does not contain any spectral parameter. To introduce such a parameter, we can perform an $SL(2, R)$ gauge transformation

$$\Gamma' = \Sigma \Gamma \Sigma^{-1} + \Sigma d\Sigma^{-1} \quad (15)$$

where $\det \Sigma = 1$. As a simple example, we choose

$$\Sigma = \begin{pmatrix} e^{i\varphi x} & 0 \\ 0 & e^{-i\varphi x} \end{pmatrix} \quad (16)$$

In the case the linear eigenvalue equations and the associated time evolution equations can be reduced to the following scattering problem:

$$\begin{aligned}\Psi_{xx} &= (2i\xi + \lambda_x)\Psi_x + [\xi^2 - i\xi\lambda_x + (K-1)u]\Psi \\ \Psi_t &= [2ui\xi + (1-K)u_x]\Psi - 2u\Psi_x\end{aligned}\quad (17)$$

Assuming that one particular solution of the prolonged ideal $\{\alpha_i, w_k\}$ is known, another solution of equations (1) can be written as

$$\begin{aligned}\tilde{u} &= \tilde{u}(u, p, q, y^i) \\ \tilde{\lambda} &= \tilde{\lambda}(\lambda, y^i) \\ \tilde{p} &= \tilde{p}(u, p, q, y^i) \\ \tilde{q} &= \tilde{q}(u, p, q, y^i)\end{aligned}\quad (18)$$

Substituting these into the set of forms

$$\begin{aligned}\tilde{\alpha}_1 &= d\tilde{u} \wedge dt - \tilde{p} dx \wedge dt \\ \tilde{\alpha}_2 &= d\tilde{p} \wedge dt - \tilde{q} dx \wedge dt \\ \tilde{\alpha}_3 &= d\tilde{p} \wedge dt + 6\tilde{u}\tilde{p} dx \wedge dt - \tilde{q} d\tilde{\lambda} \wedge dt - d\tilde{u} \wedge dx \\ \tilde{\alpha}_4 &= -d\tilde{\lambda} \wedge dx + 2\tilde{u} d\tilde{\lambda} \wedge dt + 2K\tilde{p} dx \wedge dt\end{aligned}\quad (19)$$

and requiring these be in the ring of the prolonged ideal, we have the following set of differential equations:

$$\begin{aligned}\tilde{u}_q &= \tilde{p}_q = \tilde{u}_p = 0 \\ p\tilde{u}_u &- \tilde{u}_{y^i} F^i - \tilde{p} = 0 \\ p\tilde{p}_u + q\tilde{p}_p - \tilde{p}_{y^i} F^i - \tilde{q} &= 0 \\ p\tilde{q}_u + q\tilde{q}_p - 6up\tilde{q}_q + 6\tilde{u}\tilde{p} - \tilde{q}_{y^i} F^i + \tilde{q}\tilde{\lambda}_{y^i} F^i - \tilde{u}_{y^i} G^i &= 0 \\ q\tilde{q}_q - \tilde{q}\tilde{\lambda}_\lambda &= 0 \\ \tilde{q}_q - \tilde{u}_u &= 0 \\ \tilde{\lambda}_\lambda(\tilde{u} - u) &= 0 \\ 2K(\tilde{p} - p\tilde{\lambda}_\lambda) - \tilde{\lambda}_{y^i} G^i - 2\tilde{u}\tilde{\lambda}_{y^i} F^i &= 0\end{aligned}\quad (20)$$

The solutions of these equations are trivial, i.e.,

$$\tilde{u} = u, \quad \tilde{\lambda} = \lambda, \quad \tilde{q} = q, \quad \tilde{p} = p \quad (21)$$

which means that there exists no Bäcklund transformation other than the identity map. This supports the claim that for nonlinear partial differential equations admitting only finite-dimensional prolongation algebra Bäcklund transformations other than the identity mapping seem to be absent (Leo *et al.*, 1983).

There is a close connection between solvable nonlinear evolution equations and nonlinear ordinary differential equations of Painlevé type. An ODE is said to possess the Painlevé property when all movable singularities are simple poles. Ablowitz *et al.* (1980) conjecture that a nonlinear ordinary differential equation obtained by an exact reduction of a nonlinear partial differential equation of inverse scattering transform class is either *P*-type or it must be related by a simple transformation to an ordinary differential equation that is *P*-type.

Introducing new variables

$$v = x^2 u, \quad z = x^3/t \quad (22)$$

we can write equations (1) in the form of a single nonlinear ordinary differential equation,

$$\begin{aligned}[27z^3 v''' + zv'(18v + 24 - z) - 12v(v+2)](6v - z) \\ - 6K(2v - 3v'z)(6v - 6v'z + 9z^2 v'') = 0\end{aligned}\quad (23)$$

where prime denotes derivative with respect to z . Using the algorithm for nonlinear ordinary differential equations introduced by Ablowitz *et al.* (1980), we obtain the following results:

Substituting

$$v \sim (z - z_0)^{-\alpha} v_0, \quad z_0 \text{ is arbitrary} \quad (24)$$

into equation (23), we find

$$\alpha = -2, \quad v_0 = -9z_0^2(K+2) \quad (25)$$

The resonances occur at $r_1 = -1$, $r_2 = 6$, $r_3 = 2(K+2)$.

It is obvious that for different values of K we have different resonance values. For $K \neq 0$, substituting

$$v = (z - z_0)^{-2} \sum_{j=0}^6 v_j (z - z_0)^j \quad (26)$$

into the full equation (23) and requiring that the coefficients of $(z - z_0)^j$ must vanish identically, we find that v_j corresponding to the resonance values are not arbitrary. This means that at the resonances there must be logarithmic branch points. Therefore, equation (23) is not of *P*-type for $K \neq 0$.

A different exact reduction of equations (1) to an ODE may be obtained by looking for a self-similar solution

$$u = \frac{v(z)}{(3t)^{2/3}}, \quad z = \frac{x}{(3t)^{1/3}} \quad (27)$$

where $v(z)$ satisfies

$$v'''(2v - z) + 2Kv'v'' + v'(12v^2 - 8vz + z^2) + 2vz - 4v^2 = 0 \quad (28)$$

This can be integrated once:

$$v''(2v - z) + (K - 1)(v')^2 + v' + 4v^3 - 4v^2z + z^2v + C = 0 \quad (29)$$

where C is an arbitrary integration constant.

This ODE is not of P -type, because the Painlevé expansion does not contain the correct number of arbitrary constants.

Another exact reduction of equations (1) to a different ODE may be obtained by looking for a traveling wave solution

$$u = v(z), \quad z = x - ct \quad (30)$$

where $v(z)$ satisfies

$$v''' + 6vv' - cv' + \frac{2Kv'}{2v - c} v'' = 0 \quad (31)$$

which can also be integrated once,

$$v''(2v - c) + (K - 1)(v')^2 + 4v^3 - 4cv^2 + c^2v - C' = 0 \quad (32)$$

where C' is an arbitrary integration constant. Again this ODE does not have the correct number of arbitrary constant; hence, it is not of P -type for $K \neq 0$. This ODE admits solutions of the form of the elliptic functions, but the total number of arbitrary constants is reduced to two, instead of the three needed. The ODEs (23), (28), and (31) obtained by exact reductions of Gürses–Nutku equations, which are IST class, are not of P -type in their present form.

A different and more recent Painlevé test due to Weiss *et al.* (1983) is given as follows: A partial differential equation has the Painlevé property when its solutions are “single-valued” about the movable singularity manifold. If the singularity manifold is determined by

$$\Phi(x^0, x^1, \dots, x^n) = 0 \quad (33)$$

and $u^a (a = 1, \dots, N)$ satisfy a system of partial differential equations (N -equations), then the Painlevé expansion is given by

$$u^a = \Phi^{\alpha_a} \sum_{k=0}^{\infty} u_{(k)}^a(x^0, x^1, \dots, x^n) \Phi^k \quad (34)$$

where $u_{(k)}^a$ are analytic functions of (x^0, x^1, \dots, x^n) in a neighborhood of the manifold (33). The substitution of (34) into the partial differential equations under consideration determines the possible values of α_a and gives the recursion relations for $u_{(k)}^a$.

A set of partial differential equations is said to have the Painlevé property in the sense of Weiss *et al.* provided the α_a are integers, the recursion relations are consistent, and the series expansion (34) contains the correct number of arbitrary functions. So there are basically three steps to the algorithm. Applying these steps to equations (1), we obtain the following:

1. Leading order analysis:

$$\begin{aligned} u &\sim \Phi^{\alpha_1} u_0, & \lambda &\sim \Phi^{\alpha_2} \lambda_0, \\ \alpha_1 &= -2, & \alpha_2 &= 2K \\ u_0 &= -(2+K)\Phi_x^2, & \lambda_0 &= \Phi^{-2K} \end{aligned} \quad (35)$$

Here K must take negative values in order for λ_0 to be analytic on $\Phi = 0$.

2. Resonances: Substituting

$$u = \Phi^{-2} u_0 + \beta_1 \Phi^{r-2}, \quad \lambda = \Phi^{2K} \lambda_0 + \beta_2 \Phi^{r+2K} \quad (36)$$

into the leading terms of the original equation and requiring that β_1 and β_2 remain arbitrary, we have

$$(2+K)(r+2K)[(r-2)(r-3)(r-4) - 12(r-2) + 12(2+K)] = 0 \quad (37)$$

The roots of this equation determine the resonances. We must always have the root $r_1 = -1$, since it represents the arbitrariness of the singular manifold $\Phi = 0$. The other roots should be integers. $r_1 = -1$ is a root of the equation (37) when $K = 0$. For this case the other roots are found to be $r_2 = 0$, $r_3 = 4$, $r_4 = 6$. We know that the case $K = 0$ is related to the KdV equation, which is in fact P -type. When $K = 1/2$, $r_1 = -1$ is again a root of equation (37), but this time λ_0 is not analytic on $\Phi = 0$. Therefore equations (1) are not of P -type when $K \neq 0$.

In conclusion, we have showed that the coupled differential equations (1) have no nontrivial Bäcklund transformation. Even though the coupled partial differential equations (1) have a nontrivial prolongation structure and have a Lax pair, we have showed that they do not pass the Painlevé tests of Ablowitz *et al.* and Weiss *et al.* In that respect we should reexamine the Painlevé tests for coupled partial differential equations. Presumably we must expand each field, u and λ for our case, about distinct singular manifolds.

The research reported in this paper is supported in part by the Scientific and Technical Research Council of Turkey.

REFERENCES

- Ablowitz, M. J., Ramani, A., and Segur, H. (1980). *Journal of Mathematical Physics*, **21**, 715.
 Chowdhury, A. Roy, and Paul, S. (1985). *International Journal of Theoretical Physics*, **24**, 633.
 Gürses, M., and Nukku, Y. (1981). *Journal of Mathematical Physics*, **22**, 1393.
 Leo, M., Leo, R. A., Soliani, G., Solombrino, L., and Martina, L. (1983). *Journal of Mathematical Physics*, **24**, 1720.
 Wahlquist, H. D., and Estabrook, F. B. (1975). *Journal of Mathematical Physics*, **16**, 1.
 Weiss, J., Tabor, M., and Carnevale, G. (1983). *Journal of Mathematical Physics*, **24**, 522.

Stochastic Theory of Ignition Processes

Ariel Fernández¹ and Herschel Rabitz¹

Received May 21, 1987

We consider a closed gaseous system immersed in a heat bath undergoing a thermal explosion. The effects of instantaneous fluctuations in the temperature on the heat removal mechanism and on the reaction rate are considered. The intensity of the fluctuations in situations far from equilibrium is determined by calculating the temperature self-correlation. This quantity scales with the inverse of an effective volume Ω obtained from generalized fluctuation-dissipation theory. This determines a virtual system corresponding to the localized ignition process, possibly leading to a global runaway. The induction period is identified with the Kramers mean passage time for diffusion across a kinetic barrier. The induction period is thus shown to be dependent on the fluctuation volume Ω . The diffusion process is hastened by the critical fluctuations. The explosive decomposition of ethyl azide was selected to test the theory and the results exhibit very good agreement with experimental data. Our treatment resolves the previous discrepancy between the predictions rooted in the classical Frank-Kamenetsky treatment and the premature ignition observed experimentally.

1. INTRODUCTION

The aim of this work is to provide a rigorous framework for demonstrating the stochastic contribution to ignition processes in gaseous closed systems. That is, we shall describe the phenomenon of ignition in a statistical sense. In an ensemble of realizations of the combustion experiment, some replicas of the system will present a delayed explosion and others an accelerated ignition. In order to appreciate the results presented in this work, we emphasize the physical difference between critical nonequilibrium fluctuations responsible for ignition and fluctuations near thermodynamic equilibrium. Critical fluctuations are assumed to be *localized in space* and the physical consequences of such an assumption will be examined. This issue is discussed in a general context in Graham (1975), Fernández (1985a),

¹Department of Chemistry, Princeton University, Princeton, New Jersey 08544.