

REPRODUCING KERNEL HILBERT SPACES

A THESIS

SUBMITTED TO THE DEPARTMENT OF MATHEMATICS
AND THE INSTITUTE OF ENGINEERING AND SCIENCE
OF BILKENT UNIVERSITY

IN PARTIAL FULFILLMENT OF THE REQUIREMENTS
FOR THE DEGREE OF
MASTER OF SCIENCE

By

Baver Okutmuştur

August, 2005

I certify that I have read this thesis and that in my opinion it is fully adequate, in scope and in quality, as a thesis for the degree of Master of Science.

Assist. Prof. Dr. Aurelian Gheondea (Supervisor)

I certify that I have read this thesis and that in my opinion it is fully adequate, in scope and in quality, as a thesis for the degree of Master of Science.

Prof. Dr. Mefharet Kocatepe

I certify that I have read this thesis and that in my opinion it is fully adequate, in scope and in quality, as a thesis for the degree of Master of Science.

Assoc. Prof. Dr. H. Turgay Kaptanođlu

Approved for the Institute of Engineering and Science:

Prof. Dr. Mehmet B. Baray
Director of the Institute Engineering and Science

ABSTRACT

REPRODUCING KERNEL HILBERT SPACES

Baver Okutmuştur
M.S. in Mathematics
Supervisor: Assist. Prof. Dr. Aurelian Gheondea
August, 2005

In this thesis we make a survey of the theory of reproducing kernel Hilbert spaces associated with positive definite kernels and we illustrate their applications for interpolation problems of Nevanlinna-Pick type. Firstly we focus on the properties of reproducing kernel Hilbert spaces, generation of new spaces and relationships between their kernels and some theorems on extensions of functions and kernels. One of the most useful reproducing kernel Hilbert spaces, the Bergman space, is studied in details in chapter 3. After giving a brief definition of Hardy spaces, we dedicate the last part for applications of interpolation problems of Nevanlinna-Pick type with three main theorems: interpolation with a finite number of points, interpolation with an infinite number of points and interpolation with points on the boundary. Finally we include an Appendix that contains a brief recall of the main results from functional analysis and operator theory.

Keywords: Reproducing kernel, Reproducing kernel Hilbert spaces, Bergman spaces, Hardy spaces, Interpolation, Riesz theorem.

ÖZET

DOĞURAN ÇEKİRDEKLİ HİLBERT UZAYLARI

Baver Okutmuştur

Matematik, Yüksek Lisans

Tez Yöneticisi: Yrd. Doç. Dr. Aurelian Gheondea

Ağustos, 2005

Bu tezde, doğuran çekirdekli Hilbert uzayları teorisini pozitif tanımlı çekirdekler ile beraber inceledik ve bunun uygulamalarını Nevallina-Pick interpolasyon problemleri üzerinde örnekledik. Öncelikle, doğuran çekirdekli Hilbert uzaylarının özelliklerini, üretilen yeni uzaylar ve onların çekirdekleri arasındaki ilişkileri ve genişletilen çeşitli fonksiyon ve çekirdeklerle ilgili bazı teoremleri inceledik. Sıkça kullanılan doğuran çekirdekli Hilbert uzaylarından biri olan Bergman uzayı 3. kısımda detaylarıyla işlendi. Hardy uzayının kısa bir tanımıyla başladığımız son kısım, Nevallina-Pick interpolasyon problemlerinin uygulamalarını içeren üç ana teorem ile son buldu. Bunlar: sınırlı sayıda nokta ile interpolasyon, sınırsız sayıda nokta ile interpolasyon ve sınır noktalarında interpolasyon. Son olarak Appendix kısmı bu tezde sıkça kullandığımız fonksiyonel analiz ve operator teori ile ilgili temel esasların kısa bir özetine ayrıldı.

Anahtar sözcükler: Doğuran çekirdekler, Doğuran çekirdekli Hilbert uzayları, Bergman uzayları, Hardy uzayları, İnterpolasyon, Riesz teoremi.

Contents

1	Introduction	1
2	Reproducing Kernel Hilbert Spaces	4
2.1	Definition, Uniqueness and Existence	4
2.2	Operations with Reproducing Kernel Hilbert Spaces	15
2.3	Extension of Functions and Kernels	25
3	Spaces of Analytic Functions	33
3.1	Sesqui-analytic kernels	33
3.2	Bergman Spaces	42
3.3	Szegö Kernel	48
4	Interpolation Theorems	60
4.1	General definition of Hardy spaces	60
4.2	Interpolation Inside Unit Disc	66
4.3	Interpolation on the Boundary	73

A Hilbert Spaces	76
A.1 Definitions	76
A.2 Projection	83
A.3 Weak Topology	88
A.4 Self-adjoint Operators	91

Chapter 1

Introduction

The reproducing kernel was used for the first time at the beginning of the 20th century by S. Zaremba in his work on boundary value problems for harmonic and biharmonic functions. In 1907, he was the first who introduced, in a particular case, the kernel corresponding to a class of functions, and stated its reproducing property. But he did not develop any theory and did not give any particular name to the kernels he introduced.

In 1909, J. Mercer examined the functions which satisfy reproducing property in the theory of integral equations developed by Hilbert and he called this functions as 'positive definite kernels'. He showed that this positive definite kernels have nice properties among all continuous kernels of integral equations.

However, for a long time these results were not investigated. Then the idea of reproducing kernels appeared in the dissertations of three Berlin mathematicians G. Szegö (1921), S. Bergman (1922) and S. Bochner (1922). In particular, S. Bergman introduced reproducing kernels in one and several variables for the class of harmonic and analytic functions and he called them 'kernel functions'.

In 1935, E.H. Moore examined the positive definite kernels in his general analysis under the name of positive Hermitian matrix.

Later, the theory of reproducing kernels was systematized by N.Aronszajn

around 1948.

The original idea of Zaremba to apply the kernels to the solution of boundary value problems was developed by S. Bergman and M. Schiffer. In these investigations, the kernels were proved to be powerful tool for solving boundary value problems of partial differential equations of elliptic type. Moreover, by application of kernels to conformal mapping of multiply-connected domains, very beautiful results were obtained by S. Bergman and M. Schiffer.

Several important results were achieved by the use of these kernels in the theory of one and several complex variables, in conformal mapping of simply- and multiply-connected domains, in pseudo-conformal mappings, in the study of invariant Riemannian metrics and in other subjects.

Meanwhile, in probability theory, the theory of positive definite kernels was used by A.N. Kolmogorov, E. Parzen and others.

There are also several papers and lecture notes on this subject; B. Burbea (1987), E. Hille (1972), S. Saitoh (1988), H. Dym (1989) and T. Ando (1987). Most part of this thesis owes to T. Ando's lecture notes [1] in its diversity of tools and results. We also used H. Dym, S. Saitoh and N. Aronszajn's works especially for the second chapter. Moreover, we used partially the books of P.L. Duren [4], P. Koosis [7], P.L. Duren and A. Schuster's [5] for complementing with result on Bergman and Hardy spaces.

The thesis is organized as follows:

In Chapter 2, after giving definitions and properties of reproducing kernel Hilbert spaces with some theorems, we focus on generation of new spaces and relationship between their kernels. Also, some extension theorems of functions and kernels are proven.

In Chapter 3, we present some of the most useful reproducing kernel Hilbert spaces consisting of analytic functions. A special role is played by the Bergman spaces and Bergman kernels that we present in detail.

Chapter 4 is dedicated to applications to interpolation problems of Nevanlinna-Pick type. We start with a brief definition of Hardy spaces. Then we prove three main theorems: interpolation with a finite number of points, interpolation with an infinite number of points, and interpolation with points on the boundary.

The Appendix part contains some elementary facts from functional analysis and operator theory in Hilbert spaces which can be found in textbooks, e.g. in J. Conway [3] and J. Weidman [9].

Chapter 2

Reproducing Kernel Hilbert Spaces

2.1 Definition, Uniqueness and Existence

Definition 2.1.1. Let \mathcal{H} be a Hilbert space of functions on a set X . Denote by $\langle f, g \rangle$ the inner product and let $\|f\| = \langle f, f \rangle^{1/2}$ be the norm in \mathcal{H} , for f and $g \in \mathcal{H}$. The complex valued function $K(y, x)$ of y and x in X is called a *reproducing kernel of \mathcal{H}* if the followings are satisfied:

- (i) For every x , $K_x(y) = K(y, x)$ as a function of y belongs to \mathcal{H} .
- (ii) *The reproducing property:* for every $x \in X$ and every $f \in \mathcal{H}$,

$$f(x) = \langle f, K_x \rangle. \tag{2.1}$$

So applying (2.1) to the function K_x at y , we get

$$K_x(y) = \langle K_x, K_y \rangle, \text{ for } x, y \in X,$$

and by (i),

$$K(y, x) = \langle K_x, K_y \rangle, \text{ for } x, y \in X.$$

By the above relations, for $x \in X$ we obtain $\|K_x\| = \langle K_x, K_x \rangle^{1/2} = K(x, x)^{1/2}$.

Definition 2.1.2. A Hilbert space \mathcal{H} of functions on a set X is called a *reproducing kernel Hilbert space* (sometimes abbreviated by RKHS) if there exists a reproducing kernel K of \mathcal{H} , cf. Definition 2.1.1.

The Hilbert space with reproducing kernel K is denoted by $\mathcal{H}_K(X)$. Correspondingly norm will be denoted by $\|\cdot\|_K$ (or sometimes by $\|\cdot\|_{\mathcal{H}_K}$) and inner product will be denoted by $\langle \cdot, \cdot \rangle_K$ (or sometimes by $\langle \cdot, \cdot \rangle_{\mathcal{H}_K}$), if there is a need of distinction.

Theorem 2.1.3. *If a Hilbert space \mathcal{H} of functions on a set X admits a reproducing kernel, then the reproducing kernel $K(y, x)$ is uniquely determined by the Hilbert space \mathcal{H} .*

Proof. Let $K(y, x)$ be a reproducing kernel of \mathcal{H} . Suppose that there exists another kernel $K'(y, x)$ of \mathcal{H} . Then, for all $x \in X$, applying (ii) for K and K' we get

$$\begin{aligned} \|K_x - K'_x\|^2 &= \langle K_x - K'_x, K_x - K'_x \rangle \\ &= \langle K_x - K'_x, K_x \rangle - \langle K_x - K'_x, K'_x \rangle \\ &= (K_x - K'_x)(x) - (K_x - K'_x)(x) \\ &= 0 \end{aligned}$$

Hence $K_x = K'_x$, that is, $K_x(y) = K'_x(y)$ for all $y \in X$. This means that $K(x, y) = K'(x, y)$ for all $x, y \in X$. \square

Theorem 2.1.4. *For a Hilbert space \mathcal{H} of functions on X , there exists a reproducing kernel K for \mathcal{H} if and only if for every x of X , the evaluation linear functional $\mathcal{H} \ni f \mapsto f(x)$ is a bounded linear functional on \mathcal{H} .*

Proof. Suppose that K is the reproducing kernel for \mathcal{H} . By reproducing property and Schwarz inequality of the scalar product, for all $x \in X$,

$$|f(x)| = |\langle f, K_x \rangle| \leq \|f\| \|K_x\| = \|f\| \langle K_x, K_x \rangle^{1/2} = \|f\| K(x, x)^{1/2}$$

that is, the evaluation at x is a bounded linear functional on \mathcal{H} .

Conversely, if for all $x \in X$ the evaluation $\mathcal{H} \ni f \mapsto f(x)$ is a bounded linear functional on \mathcal{H} , then by the Riesz Representation Theorem, for all $x \in X$, there exists a function g_x belonging to \mathcal{H} such that

$$f(x) = \langle f, g_x \rangle.$$

If we put K_x instead of g_x , then for all $y \in X$, we get $K_x(y) = g_x(y)$. Hence K is a reproducing kernel for \mathcal{H} . \square

Definition 2.1.5. Let X be an arbitrary set and K be a kernel on X , that is, $K: X \times X \rightarrow \mathbb{C}$. The kernel K is called *Hermitian* if for any finite set of points $\{y_1, \dots, y_n\} \subseteq X$ and any complex numbers $\epsilon_1, \dots, \epsilon_n$ we have

$$\sum_{i,j=1}^n \bar{\epsilon}_j \epsilon_i K(y_j, y_i) \in \mathbb{R}$$

and K is called *positive definite* if

$$\sum_{i,j=1}^n \bar{\epsilon}_j \epsilon_i K(y_j, y_i) \geq 0.$$

Equivalently, the last inequality means that for any finitely supported family of complex numbers $\{\epsilon_x\}_{x \in X}$ we have

$$\sum_{x,y \in X} \bar{\epsilon}_y \epsilon_x K(y, x) \geq 0. \quad (2.2)$$

In brief, sometimes we will denote this by $[K(y, x)] \geq 0$ on X , or equivalently, we will say that K is a *positive definite matrix in the sense of E. H. Moore*.

Theorem 2.1.6. *The reproducing kernel $K(y, x)$ of a reproducing kernel Hilbert space \mathcal{H} is a positive matrix in the sense of E. H. Moore.*

Proof. We have

$$\begin{aligned} 0 &\leq \left\| \sum_{i=1}^n \epsilon_i K_{y_i} \right\|^2 = \left\langle \sum_{i=1}^n \epsilon_i K_{y_i}, \sum_{j=1}^n \epsilon_j K_{y_j} \right\rangle \\ &= \sum_{i=1}^n \sum_{j=1}^n \epsilon_i \bar{\epsilon}_j \langle K_{y_i}, K_{y_j} \rangle = \sum_{i=1}^n \sum_{j=1}^n \epsilon_i \bar{\epsilon}_j K(y_j, y_i). \end{aligned}$$

Hence,

$$\sum_{i,j=1}^n K(y_j, y_i) \bar{\epsilon}_j \epsilon_i \geq 0. \quad \square$$

Remark 2.1.7. Given a reproducing kernel Hilbert space \mathcal{H} and its kernel $K(y, x)$ on X , then for all $x, y \in X$ we have the followings:

- (i) $K(y, y) \geq 0$.
- (ii) $K(y, x) = \overline{K(x, y)}$.
- (iii) $|K(y, x)|^2 \leq K(y, y)K(x, x)$, (Schwarz Inequality).
- (iv) Let $x_0 \in X$. Then the followings are equivalent:
 - (a) $K(x_0, x_0) = 0$.
 - (b) $K(y, x_0) = 0$ for all $y \in X$.
 - (c) $f(x_0) = 0$ for all $f \in \mathcal{H}$.

Indeed, (i) and (ii) can be easily seen. For (iii), we use the Schwarz Inequality in \mathcal{H} and get

$$\begin{aligned} |K(y, x)|^2 &= |\langle K_x, K_y \rangle|^2 \leq \|K_x\| \|K_y\| \|K_x\| \|K_y\| = \|K_x\|^2 \|K_y\|^2 \\ &= \langle K_x, K_x \rangle \langle K_y, K_y \rangle = K(x, x)K(y, y) \end{aligned}$$

which is the desired result.

As for (iv), it follows by (iii) that $K(x_0, x_0) = 0$ is equivalent with $K(y, x_0) = 0$ for all $y \in X$. Further, by the reproducing property, $K(y, x_0) = 0$ for all $y \in X$ if and only if $f(x_0) = 0$, for all f .

The following theorem can be regarded as a converse of Theorem 2.1.3.

Theorem 2.1.8. *For any positive definite kernel $K(y, x)$ on X , there exists a uniquely determined Hilbert space \mathcal{H}_K of functions on X , admitting the reproducing kernel $K(y, x)$.*

Proof. We denote by \mathcal{H}_0 the space of all functions f on X such that there exists a finite set of points x_1, x_2, \dots, x_n of X and complex numbers $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n$,

$$f(y) = \sum_{i=1}^n \varepsilon_i K(y, x_i),$$

for all $y \in X$. Let $g(\cdot) = \sum_{i=1}^m \mu_j K(\cdot, y_j)$ be in \mathcal{H}_0 . Define the inner product of the functions f and g from \mathcal{H}_0 by

$$\langle f, g \rangle_{\mathcal{H}_0} = \sum_{i=1}^n \sum_{j=1}^m \varepsilon_i \bar{\mu}_j \langle K(\cdot, x_i), K(\cdot, y_j) \rangle_{\mathcal{H}_0} = \sum_{i=1}^n \sum_{j=1}^m \varepsilon_i \bar{\mu}_j K(y_j, x_i). \quad (2.3)$$

Then,

$$\langle f, K(\cdot, x) \rangle_{\mathcal{H}_0} = \sum_{i=1}^n \varepsilon_i \langle K(\cdot, x_i), K(\cdot, x) \rangle = \sum_{i=1}^n \varepsilon_i K(x, x_i) = f(x) \quad (2.4)$$

for all $x \in X$, that is, \mathcal{H}_0 has the reproducing property. This implies that the definition of the inner product in (2.3) does not depend on the representations of the functions f and g in \mathcal{H}_0 . Moreover, it is easy to see that $\langle \cdot, \cdot \rangle_{\mathcal{H}_0}$ is linear in the first variable and Hermitian. Since K is positive definite it follows that $\langle f, f \rangle_{\mathcal{H}_0} \geq 0$ for all $f \in \mathcal{H}_0$, hence we have the Schwarz Inequality for $\langle \cdot, \cdot \rangle_{\mathcal{H}_0}$. In addition, if $\langle f, f \rangle_{\mathcal{H}_0} = 0$, $\|f\| = 0$ and then by (2.4) for all $x \in X$,

$$|f(x)| \leq \|f\| \|K(\cdot, x)\| = 0,$$

which implies that $f \equiv 0$. Thus, $(\mathcal{H}_0, \langle \cdot, \cdot \rangle_{\mathcal{H}_0})$ is a pre-Hilbert space.

Now denote by \mathcal{H} abstract the completion of \mathcal{H}_0 to a Hilbert space. We will show that \mathcal{H} has a unique representation as a Hilbert space with reproducing kernel $K(y, x)$. Consider first any Cauchy sequence $(f_n)_{n \geq 1}$ in \mathcal{H}_0 . Then for any $x \in X$ we have

$$\begin{aligned} |f_m(x) - f_n(x)| &= |\langle f_m, K_x \rangle_{\mathcal{H}_0} - \langle f_n, K_x \rangle_{\mathcal{H}_0}| \\ &= |\langle f_m - f_n, K_x \rangle_{\mathcal{H}_0}| \\ &\leq \|f_m - f_n\|_{\mathcal{H}_0} K(x, x)^{1/2}. \end{aligned}$$

So, there exists the function $f: X \rightarrow \mathbb{C}$ such that for all $x \in X$,

$$\lim_{n \rightarrow \infty} f_n(x) = f(x). \quad (2.5)$$

Moreover, we have

$$\|f\|_{\mathcal{H}} = \lim_{n \rightarrow \infty} \|f_n\|_{\mathcal{H}_0}$$

and for any two Cauchy sequences (f_n) and (g_n) in \mathcal{H}_0 , denoting by f and, respectively g , the corresponding pointwise limit of (f_n) and (g_n) , we have

$$\langle f, g \rangle_{\mathcal{H}} = \lim_{n, m \rightarrow \infty} \langle f_n, g_m \rangle_{\mathcal{H}_0}.$$

We can easily see that, for any two Cauchy sequences (f_n) and (g_n) , these limits exist and are independent of the approximating sequences (f_n) and (g_n) of the limits f and g , respectively.

Let us note that (2.5) yields a concrete representation of \mathcal{H} as a space of functions on X . In addition, K has the reproducing property with respect to \mathcal{H} . To see this, let $f \in \mathcal{H}$ and $(f_n) \subset \mathcal{H}$ such that $f_n \rightarrow f$ as $n \rightarrow \infty$ strongly. Then for all $x \in X$,

$$\begin{aligned} f(x) &= \lim_{n \rightarrow \infty} f_n(x) = \lim_{n \rightarrow \infty} \langle f_n, K_x \rangle_{\mathcal{H}_0} \\ &= \langle \lim_{n \rightarrow \infty} f_n, K_x \rangle_{\mathcal{H}_0} = \langle f, K_x \rangle_{\mathcal{H}} \end{aligned}$$

It remains to show the uniqueness of the Hilbert space \mathcal{H} admitting the reproducing kernel K . Suppose \mathcal{H}_1 is another Hilbert space with the same reproducing kernel K . By definition, for any $x \in X$, $K_x \in \mathcal{H}_1$ and then we have $\mathcal{H}_0 \subseteq \mathcal{H}_1$. Also, for any $f, g \in \mathcal{H}_0$, because of the reproducing property we have

$$\langle f, g \rangle_{\mathcal{H}_0} = \langle f, g \rangle_{\mathcal{H}_1}. \quad (2.6)$$

If $f \in \mathcal{H}_1$ such that $0 = \langle f, K_x \rangle_{\mathcal{H}_1} = f(x)$ for all $x \in X$, then $f \equiv 0$. Thus, the family $\{K_x : x \in X\}$ is total in \mathcal{H}_1 . So for any $f \in \mathcal{H}_1$, we can take a Cauchy sequence $(f_n)_{n \geq 1}$ in \mathcal{H}_0 such that $\lim_{n \rightarrow \infty} f_n = f$. Hence, (2.6) is valid in \mathcal{H}_0 .

Now since we have $\mathcal{H}_0 \subseteq \mathcal{H}_1$ and (2.6), we obtain $\mathcal{H} \subseteq \mathcal{H}_1$. Also from the construction of \mathcal{H} , we get $\mathcal{H}_1 \subseteq \mathcal{H}$. Thus, we have $\mathcal{H}_1 = \mathcal{H}$.

Finally, we have to show that the inner products and the norms are equal in \mathcal{H} and \mathcal{H}_1 . Consider any $f, g \in \mathcal{H}_1$ and any Cauchy sequences $(f_n)_{n \geq 1}$ and $(g_n)_{n \geq 1}$ in \mathcal{H}_0 which converge to f and g respectively. We have

$$\langle f, g \rangle_{\mathcal{H}_1} = \lim_{n \rightarrow \infty} \langle f_n, g_n \rangle_{\mathcal{H}_1} = \lim_{n \rightarrow \infty} \langle f_n, g_n \rangle_{\mathcal{H}_0} = \langle f, g \rangle_{\mathcal{H}}$$

and hence the norms in \mathcal{H} and \mathcal{H}_1 are equal. \square

Theorem 2.1.9. *Every sequence of functions $(f_n)_{n \geq 1}$ which converges strongly to a function f in $\mathcal{H}_K(X)$, converges also in the pointwise sense, that is, $\lim_{n \rightarrow \infty} f_n(x) = f(x)$, for any point $x \in X$. Further, this convergence is uniform on every subset of X on which $x \mapsto K(x, x)$ is bounded.*

Proof. For $x \in X$, using the reproducing property and the Schwartz Inequality,

$$\begin{aligned} |f(x) - f_n(x)| &= |\langle f, K_x \rangle - \langle f_n, K_x \rangle| \\ &= |\langle f - f_n, K_x \rangle| \\ &\leq \|f - f_n\| \cdot \|K_x\| \\ &= \|f - f_n\| \cdot K(x, x)^{1/2}. \end{aligned}$$

Therefore, $\lim_{n \rightarrow \infty} f_n(x) = f(x)$, for any point $x \in X$.

Moreover, it is clear from the above inequality that this convergence is uniform on every subset of X on which $x \mapsto K(x, x)$ is bounded. \square

In the following we will use the following notation: given X an abstract nonempty set and H and K two Hermitian kernels on X , we denote

$$[H(y, x)] \leq [K(y, x)] \text{ on } X, \quad (2.7)$$

whenever for any natural number n , any finite set $\{x_1, \dots, x_n\} \subseteq X$ and any complex numbers $\epsilon_1, \dots, \epsilon_n$ we have

$$\sum_{i,j=1}^n \bar{\epsilon}_j \epsilon_i H(x_j, x_i) \leq \sum_{i,j=1}^n \bar{\epsilon}_j \epsilon_i K(x_j, x_i). \quad (2.8)$$

Theorem 2.1.10. *A complex valued function g on X belongs to the reproducing kernel Hilbert space $\mathcal{H}_K(X)$ if and only if there exists $0 \leq \gamma < \infty$ such that,*

$$[g(y)\overline{g(x)}] \leq \gamma^2 [K(y, x)] \text{ on } X. \quad (2.9)$$

The minimum of all such γ coincides with $\|g\|$.

Proof. By the reproducing property, $g \in \mathcal{H}_K$ and $\|g\| \leq \gamma$ is equivalent with the existence of $f \in \mathcal{H}_K(X)$ such that $\|f\| \leq \gamma$ and $g(x) = \langle f, K_x \rangle$ for $x \in X$. By

applying the Abstract Interpolation Theorem (see Theorem A.2.6) we obtain the inequality (2.9). The converse implication is also a consequence of the Abstract Interpolation Theorem. \square

Theorem 2.1.11. *Let $K^{(1)}(y, x)$ and $K^{(2)}(y, x)$ be two positive definite kernels on X . Then the following assertions are mutually equivalent:*

- (i) $\mathcal{H}_{K^{(1)}}(X) \subseteq \mathcal{H}_{K^{(2)}}(X)$, (set inclusion).
- (ii) There exists $0 \leq \gamma < \infty$ such that

$$[K^{(1)}(y, x)] \leq \gamma^2 [K^{(2)}(y, x)].$$

If this is the case, the inclusion map J in (i) is continuous, and its norm is given by the minimum of γ in (ii).

Proof. Denote the norm and the inner product in $\mathcal{H}_{K^{(i)}}(X)$ by $\|\cdot\|_i$ and $\langle \cdot, \cdot \rangle_i$, respectively.

Let (i) be satisfied. Set $J : \mathcal{H}_{K^{(1)}}(X) \longrightarrow \mathcal{H}_{K^{(2)}}(X)$, the inclusion map.

Claim: J is a closed and continuous operator.

Suppose that $f_n \rightarrow g$ in $\mathcal{H}_{K^{(1)}}(X)$ and $f_n \rightarrow h$ in $\mathcal{H}_{K^{(2)}}(X)$. As point evaluations are continuous in $\mathcal{H}_{K^{(i)}}(X)$, ($i = 1, 2$), we get

$$f_n(x) \rightarrow g(x) \text{ and } f_n(x) \rightarrow h(x)$$

which implies that $g(x) = h(x)$ for all x , since the limit is unique. So J is closed. Since J is closed, we know that by the Closed Graph Theorem any closed linear operator between Hilbert spaces is continuous. Hence J is continuous, as claimed.

Now, for all $f \in \mathcal{H}_{K^{(1)}}(X)$ and for all $x \in X$, by reproducing property we have $f(x) = \langle f, K_x^{(1)} \rangle_1$ and $(Jf)(x) = \langle Jf, K_x^{(2)} \rangle_2$. Then by using this and the inclusion property of J , for all $x \in X$, we have

$$\langle f, J^* K_x^{(2)} \rangle_1 = \langle Jf, K_x^{(2)} \rangle_2 = (Jf)(x) = \langle f, K_x^{(1)} \rangle_1$$

and hence we obtain $J^*K_x^{(2)} = K_x^{(1)}$ for all $x \in X$.

Finally, for any $\gamma \geq \|J\|$ and any finitely supported family of complex numbers $\{\epsilon_x\}_{x \in X}$, we have

$$\begin{aligned} \sum_{x,y} \epsilon_x \bar{\epsilon}_y K^{(1)}(y,x) &= \left\langle \sum_x \epsilon_x K_x^{(1)}, \sum_y \epsilon_y K_y^{(1)} \right\rangle = \left\| \sum_x \epsilon_x K_x^{(1)} \right\|_1^2 \\ &= \left\| J^* \left(\sum_x \epsilon_x K_x^{(2)} \right) \right\|_1^2 \leq \gamma^2 \left\| \sum_x \epsilon_x K_x^{(2)} \right\|^2 \\ &= \gamma^2 \left\langle \sum_x \epsilon_x K_x^{(2)}, \sum_y \epsilon_y K_y^{(2)} \right\rangle \\ &= \gamma^2 \sum_{x,y} \epsilon_x \bar{\epsilon}_y K^{(2)}(y,x) \end{aligned}$$

Hence,

$$[K^{(1)}(y,x)] \leq \gamma^2 [K^{(2)}(y,x)].$$

Conversely, suppose that (ii) is satisfied for some $0 \leq \gamma < \infty$. This means that for any finitely supported family of complex numbers $\{\epsilon_x\}_{x \in X}$, that is denoted by $[\epsilon_x]$,

$$\sum_{x,y} \epsilon_x \bar{\epsilon}_y K^{(1)}(y,x) \leq \gamma^2 \sum_{x,y} \epsilon_x \bar{\epsilon}_y K^{(2)}(y,x).$$

Taking the minimum of γ in Theorem 2.1.10, we have the norm of any function f on X given by

$$\|f\|_i^2 = \sup_{[\epsilon_x]} \frac{|\sum_x \epsilon_x f(x)|^2}{\sum_{x,y} \epsilon_x \bar{\epsilon}_y K^{(i)}(y,x)}, \quad (i = 1, 2),$$

with $\|f\|_i = \infty$ if f is not in $\mathcal{H}_{K^{(i)}}(X)$. Now since $\{K_x^{(i)} : x \in X\}$ is total in $\mathcal{H}_{K^{(i)}}(X)$, ($i = 1, 2$) and using the Schwarz Inequality for the norms $\|f\|_1$ and $\|f\|_2$, we get

$$\|f\|_2 \leq \gamma \|f\|_1 \text{ for } f \in \mathcal{H}_{K^{(1)}}(X).$$

Hence, $\mathcal{H}_{K^{(1)}}(X) \subseteq \mathcal{H}_{K^{(2)}}(X)$ with $\|J\| \leq \gamma$. □

Suppose that there is a map ϕ from a set X to a Hilbert space \mathcal{H} such that $x \mapsto \phi_x$. Then ϕ can be used to define a positive definite kernel

$$K(y,x) = \langle \phi_x, \phi_y \rangle \text{ for } x, y \in X. \quad (2.10)$$

Theorem 2.1.12. *Let $\phi : X \mapsto \mathcal{H}$ and K be defined as in (2.10). Let T be the linear operator from \mathcal{H} to the space of functions on X , defined by*

$$(Tf)(x) = \langle f, \phi_x \rangle \text{ for } x \in X, f \in \mathcal{H}.$$

Then $\text{Ran}(T)$ coincides with $\mathcal{H}_K(X)$ and

$$\|Tf\|_K = \|P_{\mathcal{M}}f\| \text{ for } f \in \mathcal{H},$$

where \mathcal{M} is the orthogonal complement of $\ker(T)$, $P_{\mathcal{M}}$ is the orthogonal projection onto \mathcal{M} and $\|\cdot\|_K$ denotes the norm in $\mathcal{H}_K(X)$.

Proof. To see the positive definiteness of $K(y, x)$, let $X \ni x \mapsto \varepsilon_x$ be a complex valued function with finite support. Then,

$$\begin{aligned} \sum_{x,y} \overline{\varepsilon_y} \varepsilon_x K(y, x) &= \sum_{x,y} \overline{\varepsilon_y} \varepsilon_x \langle \phi_x, \phi_y \rangle = \sum_{x,y} \langle \varepsilon_x \phi_x, \varepsilon_y \phi_y \rangle \\ &= \langle \sum_x \varepsilon_x \phi_x, \sum_y \varepsilon_y \phi_y \rangle = \left\| \sum_x \varepsilon_x \phi_x \right\|^2 \geq 0 \text{ for } x, y \in X. \end{aligned}$$

Hence $K(y, x)$ is positive definite.

Let $x \in X$ and $K_x : X \rightarrow \mathbb{C}$. For all $y \in X$, $K_x(y) = \langle \phi_x, \phi_y \rangle = (T\phi_x)(y)$. So, $\text{Ran}(T)$ contains all the functions K_x , $x \in X$, where $K_x(y) = K(y, x) = \langle \phi_x, \phi_y \rangle$, $y \in X$. Since $\text{Ran}(T)$ is a linear space, then linear span of $\{K_x : x \in X\}$, that is, $\text{lin}\{K_x : x \in X\} = \mathcal{H}_0$, will be in $\text{Ran}(T)$, i.e. $\mathcal{H}_0 \subseteq \text{Ran}(T)$.

Claim: $T : \text{lin}\{\phi_x : x \in X\} \rightarrow \mathcal{H}_0$ is isometric.

Since $T\phi_x = K_x$, for all $x \in X$, then $T(\sum_x \varepsilon_x \phi_x) = \sum_x \varepsilon_x K_x$. Hence,

$$\begin{aligned} \langle T(\sum_x \varepsilon_x \phi_x), T(\sum_y \eta_y \phi_y) \rangle_K &= \langle \sum_x \varepsilon_x K_x, \sum_y \eta_y K_y \rangle_K = \sum_{x,y} \overline{\eta_y} \varepsilon_x K(y, x) \\ &= \sum_{x,y} \overline{\eta_y} \varepsilon_x \langle \phi_x, \phi_y \rangle_{\mathcal{H}} = \langle \sum_x \varepsilon_x \phi_x, \sum_y \eta_y \phi_y \rangle_{\mathcal{H}}. \end{aligned}$$

That is, $T \text{lin}\{\phi_x : x \in X\} \rightarrow \text{lin}\{K_x : x \in X\} = \mathcal{H}_0$ is isometric. Clearly, $T(\text{lin}\{\phi_x : x \in X\}) = \mathcal{H}_0$.

Now take f in $\ker(T)$. So $Tf = 0$, i.e. $(Tf)(x) = 0$ for all $x \in X$. But $(Tf)(x) = \langle f, \phi_x \rangle = 0$ for all $x \in X$ and T is linear which implies

$$f \perp \text{lin}\{\phi_x : x \in X\}.$$

If $f \in \text{lin}\{\phi_x : x \in X\}^\perp = \{\phi_x : x \in X\}^\perp$, then for all $x \in X$,

$$0 = \langle f, \phi_x \rangle = (Tf)(x).$$

That is, $Tf = 0$ and $\ker(T) = \text{lin}\{\phi_x : x \in X\}^\perp$. By this, we reach

$$\ker(T)^\perp = \text{lin}\{\phi_x : x \in X\}^{\perp\perp} = \overline{\text{lin}\{\phi_x : x \in X\}} =: \mathcal{M}.$$

As \mathcal{M} is a closed subspace, then \mathcal{H} can be written as $\mathcal{H} = \mathcal{M} \oplus \mathcal{M}^\perp$. Since

$$T : \text{lin}\{\phi_x : x \in X\} \longrightarrow \mathcal{H}_0 \subseteq \mathcal{H}_K(X)$$

is isometric and surjective and since \mathcal{H}_0 is dense in $\mathcal{H}_K(X)$, it follows that $T(\overline{\text{lin}\{\phi_x : x \in X\}}) \longrightarrow \overline{\mathcal{H}_0} = \mathcal{H}_K(X)$. Hence, $TM = \mathcal{H}_K(X) = T(\mathcal{M} \oplus \mathcal{M}^\perp) = T\mathcal{H} = \text{Ran}(T)$.

Finally, to see the equality of norms, take $f \in \mathcal{H} = \mathcal{M} \oplus \mathcal{M}^\perp$. It can be written as $f = P_{\mathcal{M}}f + (I - P_{\mathcal{M}})f$, where $I - P_{\mathcal{M}} = P_{\ker T}$. Then, since T is isometric on \mathcal{M} ,

$$\|Tf\|_K = \|T(P_{\mathcal{M}}f + P_{\ker T}f)\|_K = \|TP_{\mathcal{M}}f\|_K = \|P_{\mathcal{M}}f\|_K. \quad \square$$

The next result that concludes this section shows that the assumptions in (2.10) is by no means restrictive, if we consider positive definite kernels.

Theorem 2.1.13. (Kolmogorov Decomposition) *Let $K(y, x)$ be a positive definite kernel on an abstract set X . Then there exists a Hilbert space \mathcal{H} and a function $\phi: X \rightarrow \mathcal{H}$ such that*

$$K(y, x) = \langle \phi_x, \phi_y \rangle \text{ for } x, y \in X.$$

In addition, the Hilbert space \mathcal{H} can be chosen in such a way that the set $\{\phi_x\}_{x \in X}$ is total in \mathcal{H} and in this case the pair (ϕ, \mathcal{H}) is unique in the following sense: for any other pair (ψ, \mathcal{K}) , where $\psi: X \rightarrow \mathcal{K}$ and \mathcal{K} is a Hilbert space such that $\{\psi_x\}_{x \in X}$ is total in \mathcal{K} and $K(y, x) = \langle \psi_x, \psi_y \rangle_{\mathcal{K}}$ for all $x, y \in X$, there exists a unitary operator $U \in \mathcal{L}(\mathcal{H}, \mathcal{K})$ such that $U\phi_x = \psi_x$ for all $x \in X$.

Proof. Since K is positive definite, by Theorem 2.1.8 there exists the reproducing kernel space \mathcal{H}_K with reproducing kernel K . Let $\phi_x = K_x \in \mathcal{H}_K$ for all $x \in X$. By the reproducing property, for all $x, y \in X$ we have

$$K(y, x) = \langle K_x, K_y \rangle_{\mathcal{H}_K},$$

and $\{K_x\}_{x \in X}$ is a total subset of \mathcal{H}_K .

To prove uniqueness, let (ψ, \mathcal{K}) be a pair as in the statement and define $U\phi_x = \psi_x$ for all $x \in X$. Clearly U extends by linearity as a linear mapping $U: \text{lin}\{\phi_x: x \in X\} \rightarrow \text{lin}\{\psi_x: x \in X\}$. In addition, for any finitely supported families of complex numbers $\{\epsilon_x\}_{x \in X}$ and $\{\eta_y\}_{y \in X}$ we have

$$\begin{aligned} \langle U\left(\sum_{x \in X} \epsilon_x \phi_x\right), U\left(\sum_{y \in X} \eta_y \phi_y\right) \rangle_{\mathcal{K}} &= \langle \left(\sum_{x \in X} \epsilon_x \psi_x\right), \left(\sum_{y \in X} \eta_y \psi_y\right) \rangle_{\mathcal{K}} \\ &= \sum_{x, y \in X} \bar{\epsilon}_y \epsilon_x \langle \psi_x, \psi_y \rangle_{\mathcal{K}} \\ &= \sum_{x, y \in X} \bar{\epsilon}_y \epsilon_x K(y, x) = \sum_{x, y \in X} \bar{\epsilon}_y \epsilon_x \langle \phi_x, \phi_y \rangle_{\mathcal{H}_K} \\ &= \langle \left(\sum_{x \in X} \epsilon_x \phi_x\right), \left(\sum_{y \in X} \eta_y \phi_y\right) \rangle_{\mathcal{H}_K} \end{aligned}$$

which shows that U is isometric. Due to the fact that both families $\{\phi_x\}_{x \in X}$ and $\{\psi_y\}_{y \in X}$ are total in \mathcal{H}_K and, respectively, \mathcal{K} , it follows that U can be uniquely extended to a unitary operator $U: \mathcal{H}_K \rightarrow \mathcal{K}$. By definition, U satisfies the condition $U\phi_x = \psi_x$ for all $x \in X$. \square

2.2 Operations with Reproducing Kernel Hilbert Spaces

Let $K(y, x)$ be a positive definite kernel on X and $\mathcal{H} = \mathcal{H}_K(X)$ be the RKHS. Let \mathcal{M} be a closed subspace of $\mathcal{H}_K(X)$. We know \mathcal{M} is a Hilbert space since it is closed. As every point evaluation functional is continuous in $\mathcal{H}_K(X)$ and \mathcal{M} is a closed subspace, then every point evaluation functional is continuous also in \mathcal{M} . Thus, \mathcal{M} is a RKHS.

Denote by $P_{\mathcal{M}}$ the orthogonal projection onto \mathcal{M} . This means that, for $h \in \mathcal{H}_K(X)$, $P_{\mathcal{M}}(h) = h_{\mathcal{M}} \in \mathcal{M}$ where $h = h_{\mathcal{M}} + h_{\mathcal{M}^\perp}$, with $h_{\mathcal{M}} \in \mathcal{M}$, $h_{\mathcal{M}^\perp} \in \mathcal{M}^\perp$. If $P_{\mathcal{M}}K_x \in \mathcal{M}$, we have $f(x) = \langle f, P_{\mathcal{M}}K_x \rangle$ for all $f \in \mathcal{M}$. Then we have the reproducing kernel $K^{\mathcal{M}}(y, x)$ for \mathcal{M} as

$$K^{\mathcal{M}}(y, x) = \langle P_{\mathcal{M}}K_x, P_{\mathcal{M}}K_y \rangle = \langle P_{\mathcal{M}}P_{\mathcal{M}}^*K_x, K_y \rangle = \langle P_{\mathcal{M}}K_x, K_y \rangle.$$

Let $K^{(0)}(y, x)$ be the restriction of $K(y, x)$ to a subset X_0 of X , i.e. $K^{(0)}(y, x) = K(y, x)|_{X_0 \times X_0}$. Since $K(y, x)$ is positive definite, then so is $K^{(0)}(y, x)$. Then by Theorem 2.1.8 there exists a unique RKHS admitting $K^{(0)}(y, x)$ as its reproducing kernel. Denote this RKHS by $\mathcal{H}_{K^{(0)}}(X)$. The following theorem gives the relation between $\mathcal{H}_K(X)$ and $\mathcal{H}_{K^{(0)}}(X)$.

Theorem 2.2.1. *Let $K^{(0)}(y, x)$ be the restriction of $K(y, x)$ to a subset X_0 of X , $\mathcal{H}_{K^{(0)}}(X)$ be the RKHS admitting $K^{(0)}(y, x)$ as its reproducing kernel and $\mathcal{H}_K(X)$ be the RKHS with its reproducing kernel $K(y, x)$. Then*

$$\mathcal{H}_{K^{(0)}}(X_0) = \{f|_{X_0} : f \in \mathcal{H}_K(X)\} \quad (2.11)$$

and

$$\|h\|_{K^{(0)}} = \min\{\|f\|_K : f|_{X_0} = h\} \quad \text{for } h \in \mathcal{H}_{K^{(0)}}(X_0). \quad (2.12)$$

Proof. For $x, y \in X_0$, we have

$$K^{(0)}(y, x) = K(y, x) \text{ and so } \langle K_x^{(0)}, K_y^{(0)} \rangle_{K^{(0)}} = \langle K_x, K_y \rangle_K.$$

Define a map S such that $SK_x^{(0)} = K_x$ for all $x \in X_0$, which is uniquely extended to an isometry from the closed linear span of $\{K_x^{(0)} : x \in X_0\}$ that coincides with $\mathcal{H}_{K^{(0)}}(X_0)$ onto the closed linear span \mathcal{M} of $\{K_x : x \in X_0\}$.

Let $T = SP_{\mathcal{M}}$ where $P_{\mathcal{M}}$ is the orthogonal projection to \mathcal{M} , i.e. $P_{\mathcal{M}} : \mathcal{H}_K(X) \rightarrow \mathcal{M}$. We have

$$T : \mathcal{H}_K(X) = \mathcal{M} \oplus \mathcal{M}^\perp \rightarrow \mathcal{H}_{K^{(0)}}(X_0).$$

Since

$$(TK_x)(y) = K_x^{(0)}(y) = K_x(y) \text{ for } x, y \in X_0,$$

then

$$(Tf)(y) = f(y) \text{ for } f \in \mathcal{M} \text{ and } y \in X_0$$

and when $(Tf)(y) = 0$, $(Tf)(y) = f(y) = \langle f, K_y \rangle = 0$ for $f \in \mathcal{M}^\perp$ and $y \in X_0$. So T is the restriction map to X_0 and

$$(Tf)(x) = \langle Tf, K_x^{(0)} \rangle_{K^{(0)}} = \langle f, T^* K_x^{(0)} \rangle_K, \text{ for } x \in X_0.$$

Hence we have $(Tf)(x) = \langle f, \phi_x \rangle = \langle f, T^* K_x^{(0)} \rangle_K$ which gives us $\phi_x = T^* K_x^{(0)}$. By this with Theorem 2.1.12 and taking into account that T^* is isometric,

$$\langle \phi_x, \phi_y \rangle_K = \langle T^* K_x^{(0)}, T^* K_y^{(0)} \rangle_K = \langle K_x^{(0)}, K_y^{(0)} \rangle_{K^{(0)}} = K^{(0)}(y, x),$$

for all $x, y \in X_0$. □

Let $K^{(1)}(y, x)$ and $K^{(2)}(y, x)$ be two positive definite kernels. Then

$$K(y, x) = K^{(1)}(y, x) + K^{(2)}(y, x)$$

is also a positive definite kernel.

Let $\mathcal{H}_{K^{(1)}}$, $\mathcal{H}_{K^{(2)}}$ and \mathcal{H}_K be RKHSs with reproducing kernels $K^{(1)}(y, x)$, $K^{(2)}(y, x)$ and $K(y, x)$, respectively, with $K = K^{(1)} + K^{(2)}$.

Theorem 2.2.2. *Let $K^{(1)}(y, x)$ and $K^{(2)}(y, x)$ be two positive definite kernels and $K = K^{(1)} + K^{(2)}$. Then*

$$\mathcal{H}_K(X) = \mathcal{H}_{K^{(1)}}(X) + \mathcal{H}_{K^{(2)}}(X), \quad (\text{algebraic sum})$$

and for $f \in \mathcal{H}_{K^{(1)}}(X)$ and $g \in \mathcal{H}_{K^{(2)}}(X)$,

$$\|f + g\|_K^2 = \min\{\|f + h\|_{K^{(1)}}^2 + \|g - h\|_{K^{(2)}}^2 : h \in \mathcal{H}_{K^{(1)}}(X) \cap \mathcal{H}_{K^{(2)}}(X)\}. \quad (2.13)$$

Proof. We have

$$K(y, x) = K^{(1)}(y, x) + K^{(2)}(y, x) = \langle K_x^{(1)}, K_y^{(1)} \rangle_{K^{(1)}} + \langle K_x^{(2)}, K_y^{(2)} \rangle_{K^{(2)}}.$$

Consider the direct sum Hilbert space $\mathcal{H} = \mathcal{H}_{K^{(1)}} \oplus \mathcal{H}_{K^{(2)}}$. Since both $\mathcal{H}_{K^{(1)}}$ and $\mathcal{H}_{K^{(2)}}$ are Hilbert spaces, so is \mathcal{H} . Then, by the definition of inner product for direct sum, we have

$$\langle K_x, K_y \rangle_K = K(y, x) = K^{(1)}(y, x) + K^{(2)}(y, x) = \langle K_x^{(1)} \oplus K_x^{(2)}, K_y^{(1)} \oplus K_y^{(2)} \rangle_K.$$

Consider the map ϕ such that $\phi(x) = K_x^{(1)} \oplus K_x^{(2)}$. Then we have

$$K(y, x) = \langle K_x^{(1)} \oplus K_x^{(2)}, K_y^{(1)} \oplus K_y^{(2)} \rangle_K = \langle \phi_x, \phi_y \rangle.$$

Using the operator T in Theorem 2.1.12, where $(Tf)(x) = \langle f, \phi_x \rangle$, we get

$$\begin{aligned} (T(f \oplus g))(x) &= \langle f \oplus g, \phi_x \rangle = \langle f \oplus g, K_x^{(1)} \oplus K_x^{(2)} \rangle_K \\ &= \langle f, K_x^{(1)} \rangle_{K^{(1)}} + \langle g, K_x^{(2)} \rangle_{K^{(2)}} \\ &= f(x) + g(x). \end{aligned}$$

So, $T(f \oplus g) = f + g$. This shows by Theorem 2.1.12 that $\mathcal{H}_K(X) = \text{Ran}(T) = \mathcal{H}_{K^{(1)}}(X) + \mathcal{H}_{K^{(2)}}(X)$. Again by the same theorem,

$$\|f + g\|^2 = \|P_{\mathcal{M}}(f \oplus g)\|^2, \text{ where } \mathcal{M} = (\ker(T))^\perp.$$

Next we show that

$$\ker(T) = \{h \oplus (-h) : h \in \mathcal{H}_{K^{(1)}}(X) \cap \mathcal{H}_{K^{(2)}}(X)\}.$$

If $h \in \mathcal{H}_{K^{(1)}}(X) \cap \mathcal{H}_{K^{(2)}}(X)$, then $T(h \oplus (-h)) = h - h = 0$. Conversely, if $h_1 \oplus h_2 \in \ker(T)$, then $0 = T(h_1 \oplus h_2) = h_1 + h_2$ implies that $h_2 = -h_1$. Thus $h \in \mathcal{H}_{K^{(1)}}(X) \cap \mathcal{H}_{K^{(2)}}(X)$.

Then by Theorem 2.1.12 we have $\mathcal{M} = (\ker(T))^\perp$ which implies $\mathcal{M}^\perp = \ker(T)$. So, $h \oplus (-h) \in \mathcal{M}^\perp$. Consider the quotient

$$\mathcal{H}/\mathcal{M}^\perp := \{\tilde{h} + \mathcal{M}^\perp : \tilde{h} \in \mathcal{H}\}.$$

Let $f \in \mathcal{H}_{K^{(1)}}(X)$, $g \in \mathcal{H}_{K^{(2)}}(X)$ and $f \oplus g \in \mathcal{H} = \mathcal{H}_{K^{(1)}} \oplus \mathcal{H}_{K^{(2)}}$. Then, for $\hat{k} \in \mathcal{H}/\mathcal{M}^\perp$,

$$\hat{k} = \{\tilde{k} + \tilde{h} : \tilde{h} \in \mathcal{M}^\perp\},$$

where $\tilde{k} = f \oplus g \in \mathcal{H}$, $\tilde{h} = h \oplus (-h) \in \mathcal{M}^\perp$. Then,

$$\hat{h} = \{f \oplus g + h \oplus (-h) : f \oplus g \in \mathcal{H}, h \oplus (-h) \in \mathcal{M}^\perp\}.$$

Taking the norm of both sides, it follows that

$$\begin{aligned} \|\hat{h}\| &= \inf\{\|f \oplus g + h \oplus (-h)\| : f \oplus g \in \mathcal{H}, h \oplus (-h) \in \mathcal{M}^\perp\} = \|P_{\mathcal{M}}\tilde{k}\| \\ &= \|P_{\mathcal{M}}(f \oplus g)\| = \|f + g\|. \end{aligned}$$

Taking the square, we get

$$\|f + g\|^2 = \inf\{\|f \oplus g + h \oplus (-h)\|^2 : f \oplus g \in \mathcal{H}, \quad h \oplus (-h) \in \mathcal{M}^\perp\}.$$

Then,

$$\begin{aligned} \|f \oplus g + h \oplus (-h)\|^2 &= \langle f \oplus g + h \oplus (-h), f \oplus g + h \oplus (-h) \rangle \\ &= \langle f \oplus g, f \oplus g \rangle + \langle f \oplus g, h \oplus (-h) \rangle + \langle h \oplus (-h), f \oplus g \rangle + \langle h \oplus (-h), h \oplus (-h) \rangle \\ &= \langle f, f \rangle + \langle g, g \rangle + \langle f, h \rangle + \langle g, -h \rangle + \langle h, f \rangle + \langle -h, g \rangle + \langle h, h \rangle + \langle -h, -h \rangle \\ &= \|f + h\|^2 + \|g - h\|^2. \end{aligned}$$

Hence, we get

$$\|f + g\|^2 = \min\{\|f + h\|^2 + \|g - h\|^2 : f \in \mathcal{H}_{K^{(1)}}(X), g \in \mathcal{H}_{K^{(2)}}(X), \\ h \in \mathcal{H}_{K^{(1)}}(X) \cap \mathcal{H}_{K^{(2)}}(X)\}. \quad \square$$

Given Hilbert spaces $\mathcal{H}_{K^{(1)}}(X)$ and $\mathcal{H}_{K^{(2)}}(X)$, the intersection $\mathcal{H}_{K^{(1)}}(X) \cap \mathcal{H}_{K^{(2)}}(X)$ will be again a Hilbert space of functions on X with respect to the norm

$$\|f\|^2 := \|f\|_{K^{(1)}}^2 + \|f\|_{K^{(2)}}^2.$$

Since every point evaluation functional is continuous in both $\mathcal{H}_{K^{(1)}}(X)$ and $\mathcal{H}_{K^{(2)}}(X)$, letting $f \in \mathcal{H}_{K^{(1)}}(X) \cap \mathcal{H}_{K^{(2)}}(X)$, it follows that every point evaluation functional will be continuous in $\mathcal{H}_{K^{(1)}}(X) \cap \mathcal{H}_{K^{(2)}}(X)$. Therefore the intersection Hilbert space is a RKHS.

Theorem 2.2.3. *The reproducing kernel of the space*

$$\mathcal{H}_K(X) = \mathcal{H}_{K^{(1)}}(X) \cap \mathcal{H}_{K^{(2)}}(X)$$

is determined, as a quadratic form, by

$$\sum_{x,y} \overline{\varepsilon_y} \varepsilon_x K(y, x) = \inf \left\{ \sum_{x,y} \overline{\eta_y} \eta_x K^{(1)}(y, x) + \sum_{x,y} \overline{\zeta_y} \zeta_x K^{(2)}(y, x) : [\varepsilon_x] = [\eta_x] + [\zeta_x] \right\},$$

where $[\varepsilon_x]$ is an arbitrary complex valued function on X with finite support, and the same are true for $[\eta_x]$ and $[\zeta_x]$.

Proof. Consider the isometric map S , such that it embeds $\mathcal{H}_K(X)$ into the direct sum Hilbert space $\mathcal{H}_{K^{(1)}}(X) \oplus \mathcal{H}_{K^{(2)}}(X)$, $S : \mathcal{H}_K(X) \longrightarrow \mathcal{H}_{K^{(1)}}(X) \oplus \mathcal{H}_{K^{(2)}}(X)$ that is

$$Sf = f \oplus f \text{ for } f \in \mathcal{H}_K(X).$$

Let $P_{\mathcal{M}}$ be the orthogonal projection onto $\text{Ran}(S) := \mathcal{M}$. Then $P_{\mathcal{M}} = SS^*$, and by using the reproducing and algebraic direct sum properties, it follows

$$\begin{aligned} (Sf)(x) &= \langle Sf, K_x^{(1)} \oplus K_x^{(2)} \rangle = \langle f \oplus f, K_x^{(1)} \oplus K_x^{(2)} \rangle \\ &= \langle f, K_x^{(1)} \rangle_{K^{(1)}} + \langle f, K_x^{(2)} \rangle_{K^{(2)}} \\ &= f(x) + f(x) = 2f(x) \text{ where } f \in \mathcal{H}_K(X). \end{aligned}$$

So, $\langle Sf, K_x^{(1)} \oplus K_x^{(2)} \rangle = 2f(x)$, i.e. $\frac{1}{2} \langle Sf, K_x^{(1)} \oplus K_x^{(2)} \rangle = f(x)$. This implies

$$\frac{1}{2} \langle f, S^*(K_x^{(1)} \oplus K_x^{(2)}) \rangle = f(x)$$

or equivalently

$$\langle f, \frac{1}{2} S^*(K_x^{(1)} \oplus K_x^{(2)}) \rangle = f(x).$$

In other words, $K_x = \frac{1}{2} S^*(K_x^{(1)} \oplus K_x^{(2)})$ for $x \in X$. Then using this and the isometricity of S ,

$$\begin{aligned} \sum_{x,y} \overline{\varepsilon_y} \varepsilon_x K(y,x) &= \left\| \sum_x \varepsilon_x K_x \right\|_K^2 = \left\| \sum_x \varepsilon_x \frac{1}{2} S^*(K_x^{(1)} \oplus K_x^{(2)}) \right\|^2 \\ &= \left\| \frac{1}{2} SS^* \sum_x \varepsilon_x (K_x^{(1)} \oplus K_x^{(2)}) \right\|^2 \\ &= \left\| P_{\mathcal{M}} \left(\sum_x \varepsilon_x (K_x^{(1)} \oplus K_x^{(2)}) \right) \right\|^2. \end{aligned}$$

Now, since $\mathcal{M} = \text{Ran}(S) = (\ker(S))^\perp$, then $\mathcal{M} = (\{K_x^{(1)} \oplus (-K_x^{(2)}) : x \in X\})^\perp$ which implies

$$\begin{aligned} \mathcal{M}^\perp &= [\{K_x^{(1)} \oplus (-K_x^{(2)}) : x \in X\}]^{\perp\perp} \\ &= \overline{\text{lin}\{K_x^{(1)} \oplus (-K_x^{(2)}) : x \in X\}} \\ &= \{K_x^{(1)} \oplus (-K_x^{(2)}) : x \in X\}. \end{aligned}$$

So, the elements of the form $\frac{1}{2} \sum_x \lambda_x (K_x^{(1)} \oplus (-K_x^{(2)}))$ are dense in \mathcal{M}^\perp . Then, by using Theorem 2.1.12 and the property of orthogonal projection, we get

$$\begin{aligned}
& \|P_{\mathcal{M}}(\frac{1}{2} \sum_x \varepsilon_x (K_x^{(1)} \oplus K_x^{(2)}))\|^2 \\
&= \|\frac{1}{2} \sum_x \varepsilon_x (K_x^{(1)} \oplus K_x^{(2)}) \oplus \frac{1}{2} \sum_x \varepsilon_x (K_x^{(1)} \oplus K_x^{(2)})\|^2 \\
&= \langle \frac{1}{2} \sum_x \varepsilon_x (K_x^{(1)} \oplus K_x^{(2)}) \oplus \frac{1}{2} \sum_x \varepsilon_x (K_x^{(1)} \oplus K_x^{(2)}), \\
&\quad \frac{1}{2} \sum_x \varepsilon_x (K_x^{(1)} \oplus K_x^{(2)}) \oplus \frac{1}{2} \sum_x \varepsilon_x (K_x^{(1)} \oplus K_x^{(2)}) \rangle \\
&= \langle \frac{1}{2} \sum_x \varepsilon_x (K_x^{(1)} \oplus K_x^{(2)}), \frac{1}{2} \sum_x \varepsilon_x (K_x^{(1)} \oplus K_x^{(2)}) \rangle \\
&\quad + \langle \frac{1}{2} \sum_x \varepsilon_x (K_x^{(1)} \oplus K_x^{(2)}), \frac{1}{2} \sum_x \varepsilon_x (K_x^{(1)} \oplus K_x^{(2)}) \rangle \\
&= \|\frac{1}{2} \sum_x \varepsilon_x (K_x^{(1)} \oplus K_x^{(2)})\|^2 + \|\frac{1}{2} \sum_x \varepsilon_x (K_x^{(1)} \oplus K_x^{(2)})\|^2 \\
&= \frac{1}{2} \|\sum_x \varepsilon_x (K_x^{(1)} \oplus K_x^{(2)})\|^2.
\end{aligned}$$

Let $\frac{1}{2}(\varepsilon_x + \lambda_x) = \eta_x$, $\frac{1}{2}(\varepsilon_x - \lambda_x) = \delta_x$ and $\frac{1}{2}(\varepsilon_x + \lambda_x) + \frac{1}{2}(\varepsilon_x - \lambda_x) = \varepsilon_x$, that is,

$$[\varepsilon_x] = [\eta_x] + [\delta_x].$$

Finally, applying Theorem 2.2.2 to the function $\frac{1}{2} \sum_x \varepsilon_x (K_x^{(1)} \oplus K_x^{(2)})$, we get

$$\frac{1}{2} \|\sum_x \varepsilon_x (K_x^{(1)} \oplus K_x^{(2)})\|^2 = \inf_{[\lambda_x]} \|\left(\sum_x \frac{1}{2}(\varepsilon_x + \lambda_x) K_x\right) \oplus \left(\sum_x \frac{1}{2}(\varepsilon_x - \lambda_x) K_x\right)\|^2$$

where

$$\begin{aligned}
& \|\left(\sum_x \frac{1}{2}(\varepsilon_x + \lambda_x) K_x\right) \oplus \left(\sum_x \frac{1}{2}(\varepsilon_x - \lambda_x) K_x\right)\|^2 \\
&= \|\left(\sum_x \eta_x K_x^{(1)}\right) \oplus \left(\sum_x \delta_x K_x^{(2)}\right)\|^2 \\
&= \langle \sum_x \eta_x K_x^{(1)} \oplus \sum_x \delta_x K_x^{(2)}, \sum_x \eta_x K_x^{(1)} \oplus \sum_x \delta_x K_x^{(2)} \rangle \\
&= \|\sum_x \eta_x K_x^{(1)}\|_{K^{(1)}}^2 + \|\sum_x \delta_x K_x^{(2)}\|_{K^{(2)}}^2 \\
&= \sum_{x,y} \bar{\eta}_y \eta_x K^{(1)}(y, x) + \sum_{x,y} \bar{\delta}_y \delta_x K^{(2)}(y, x)
\end{aligned}$$

which completes the proof. \square

Remark 2.2.4. Consider the tensor product Hilbert space $\mathcal{H}_{K^{(1)}}(X) \otimes \mathcal{H}_{K^{(2)}}(X)$. Take $g \in \mathcal{H}_{K^{(1)}}(X)$, $h \in \mathcal{H}_{K^{(2)}}(X)$ and $x, x' \in X$. It follows

$$(g \otimes h)(x, x') = g(x)h(x') = \langle g, K_x^{(1)} \rangle \langle h, K_{x'}^{(2)} \rangle = \langle g \otimes h, K_x^{(1)} \otimes K_{x'}^{(2)} \rangle$$

which shows that the tensor product Hilbert space $\mathcal{H}_{K^{(1)}}(X) \otimes \mathcal{H}_{K^{(2)}}(X)$ is a RKHS on $X \times X$.

Consider the map $\phi : X \longrightarrow \mathcal{H}_{K^{(1)}}(X) \otimes \mathcal{H}_{K^{(2)}}(X)$ defined by $x \mapsto K_x^{(1)} \otimes K_x^{(2)}$. Then

$$\begin{aligned} K(y, x) &= \langle \phi_x, \phi_y \rangle = \langle K_x^{(1)} \otimes K_x^{(2)}, K_y^{(1)} \otimes K_y^{(2)} \rangle \\ &= \langle K_x^{(1)}, K_y^{(1)} \rangle \cdot \langle K_x^{(2)}, K_y^{(2)} \rangle \\ &= K^{(1)}(y, x) \cdot K^{(2)}(y, x) \quad \text{for } x, y \in X. \end{aligned}$$

Hence the pointwise product of two positive definite kernels is again a positive definite kernel.

Theorem 2.2.5. For the product kernel $K(y, x) = K^{(1)}(y, x) \cdot K^{(2)}(y, x)$, the RKHS $\mathcal{H}_K(X)$ consists of all functions f on X for which there are sequences $(g_n)_{n \geq 0}$ of functions in $\mathcal{H}_{K^{(1)}}(X)$ and $(h_n)_{n \geq 0}$ of functions in $\mathcal{H}_{K^{(2)}}(X)$ such that

$$\sum_1^\infty \|g_n\|_{K^{(1)}}^2 \|h_n\|_{K^{(2)}}^2 < \infty \quad \text{and} \quad \sum_1^\infty g_n(x)h_n(x) = f(x) \quad \text{for all } x \in X, \quad (2.14)$$

and the norm is given by

$$\|f\|_K^2 = \min \left\{ \sum_1^\infty \|g_n\|_{K^{(1)}}^2 \|h_n\|_{K^{(2)}}^2 \right\},$$

where the minimum is taken over the set of all sequences $(g_n)_{n \geq 0}$ and $(h_n)_{n \geq 0}$ satisfying (2.14).

Proof. Let T be an operator from $\mathcal{H}_{K^{(1)}} \otimes \mathcal{H}_{K^{(2)}}$ to the space of functions on X , associated with ϕ_x , as in Theorem 2.1.12 more precisely,

$$T : \mathcal{H}_{K^{(1)}} \otimes \mathcal{H}_{K^{(2)}} \longrightarrow \mathcal{F}(X) := \{f : X \longrightarrow \mathbb{C} : f \text{ complex function on } X\}.$$

Let $F \in \mathcal{H}_{K^{(1)}} \otimes \mathcal{H}_{K^{(2)}}$. Then F will be of the form

$$F = \sum_1^{\infty} g_n \otimes h_n \text{ with } g_n \in \mathcal{H}_{K^{(1)}} \text{ and } h_n \in \mathcal{H}_{K^{(2)}}.$$

Let $x \in X$ and $\phi_x := K_x^{(1)} \otimes K_x^{(2)}$, $\phi_x : X \longrightarrow \mathcal{H}_{K^{(1)}} \otimes \mathcal{H}_{K^{(2)}}$ and $K(y, x) = \langle \phi_x, \phi_y \rangle$.

It follows by Theorem 2.1.12

$$\begin{aligned} (TF)(x) &= \langle F, \phi_x \rangle = \langle F, K_x^{(1)} \otimes K_x^{(2)} \rangle = \left\langle \sum_{n=1}^{\infty} g_n \otimes h_n, K_x^{(1)} \otimes K_x^{(2)} \right\rangle \\ &= \sum_{n=1}^{\infty} \langle g_n, K_x^{(1)} \rangle \langle h_n, K_x^{(2)} \rangle \\ &= \sum_{n=1}^{\infty} g_n(x) h_n(x) = f(x). \end{aligned}$$

Finally,

$$\begin{aligned} \|F\|^2 &= \langle F, F \rangle = \left\langle \sum_{n=1}^{\infty} g_n \otimes h_n, \sum_{n=1}^{\infty} g_n \otimes h_n \right\rangle \\ &= \sum_{n=1}^{\infty} \langle g_n, g_n \rangle_{K^{(1)}} \langle h_n, h_n \rangle_{K^{(2)}} \\ &= \sum_{n=1}^{\infty} \|g_n\|_{K^{(1)}}^2 \|h_n\|_{K^{(2)}}^2. \end{aligned}$$

Taking the norm of $(TF)(x) = f(x)$, again by Theorem 2.1.12 we get

$$\|f(x)\|_K = \|TF\|_K = \|P_{\mathcal{M}}F\|_K = \min \left\{ \sum_1^{\infty} \|g_n\|_{K^{(1)}}^2 \|h_n\|_{K^{(2)}}^2 \right\}. \quad \square$$

Remark 2.2.6. If X consists of a finite number of points, say n , then the space of all functions on X , that is \mathbb{C}^n , has the canonical RKHS structure $(l_n^2, \langle \cdot, \cdot \rangle)$, where the point evaluation at i is induced by the inner product with e_i , ($i = 1, 2, \dots, n$). Moreover, for a positive definite kernel $K(j, i)$ on X , we have

$$K(j, i) = \langle L e_i, e_j \rangle, \quad (i, j = 1, 2, \dots, n) \quad (2.15)$$

where L is a uniquely determined linear operator on l_n^2 and is positive definite.

Theorem 2.2.7. *If $K(j, i)$ is a strictly positive definite kernel on $X = \{1, 2, \dots, n\}$ and the operator L on l_n^2 is defined as in (2.15), then L is a strictly positive definite operator and*

$$\langle f, g \rangle_K = \langle L^{-1}f, g \rangle \text{ for } f, g \in \mathbb{C}^n. \quad (2.16)$$

Proof. Given $K(j, i)$ a positive definite kernel on $X = \{1, 2, \dots, n\}$, consider the inclusion map $J : \mathcal{H}_K(X) \longrightarrow \mathbb{C}^n = l_n^2$. As a result of Theorem 2.1.11, J is continuous.

Let J^* be the adjoint of J . We have $J^* : l_n^2 \longrightarrow \mathcal{H}_K(X)$. Let $J^*(e_i) = K_i$, ($i = 1, 2, \dots, n$). By (2.15),

$$\langle Le_i, e_j \rangle = K(j, i) = \langle K_i, K_j \rangle_K = \langle J^*e_i, J^*e_j \rangle_K = \langle (JJ^*e_i), e_j \rangle$$

which gives

$$L = JJ^*.$$

Since K is a strictly positive definite kernel, it follows that $\dim(\mathcal{H}_K(X)) = n$ and J is a bijection. Hence,

$$\begin{aligned} \langle f, g \rangle_K &= \langle J^{-1}f, J^{-1}g \rangle_K = \langle (J^{-1})^*J^{-1}f, g \rangle \\ &= \langle (JJ^*)^{-1}f, g \rangle = \langle L^{-1}f, g \rangle \end{aligned}$$

and consequently we have

$$\langle f, g \rangle_K = \langle L^{-1}f, g \rangle. \quad \square$$

Each positive definite operator L on l_n^2 produces a positive definite kernel $K(j, i)$ on X by (2.15).

Theorem 2.2.8. *If L_i , ($i = 1, 2$) are two strictly positive definite operators on l_n^2 , then*

$$\langle (L_1^{-1} + L_2^{-1})f, f \rangle = \min \{ \langle L_1g, g \rangle + \langle L_2h, h \rangle : g + h = f \} \text{ for } f \in \mathbb{C}^n. \quad (2.17)$$

Proof. Let $K^{(1)}$ and $K^{(2)}$ be the kernels associated to L_1 and L_2 , respectively. By using (2.15) and the result of previous theorem, we have

$$\langle f, g \rangle_{K^{(i)}} = \langle L_i^{-1} f, g \rangle \text{ for } f, g \in \mathbb{C}^n.$$

Now consider the inner product $\langle f, f \rangle_{K^{(1)}+K^{(2)}}$ in Theorem 2.2.2,

$$\begin{aligned} \langle f, f \rangle_{K^{(1)}+K^{(2)}} &= \|f\|_{K^{(1)}+K^{(2)}}^2 = \min \{ \|g\|_{K^{(1)}}^2 + \|h\|_{K^{(2)}}^2 : g + h = f \} \\ &= \min \{ \langle g, g \rangle_{K^{(1)}} + \langle h, h \rangle_{K^{(2)}} : g + h = f \} \\ &= \min \{ \langle L_1 g, g \rangle + \langle L_2 h, h \rangle : g + h = f \} \text{ for } f, g \text{ and } h \in \mathbb{C}^n, \end{aligned}$$

and since by (2.16) in Theorem 2.2.7, we have $\langle f, f \rangle = \langle (L_1^{-1} + L_2^{-1})^{-1} f, f \rangle$, combining this with the above equations, we obtain

$$\begin{aligned} \langle f, f \rangle &= \langle (L_1^{-1} + L_2^{-1})^{-1} f, f \rangle \\ &= \min \{ \langle L_1 g, g \rangle + \langle L_2 h, h \rangle : g + h = f \text{ for } f \in \mathbb{C}^n \}. \quad \square \end{aligned}$$

2.3 Extension of Functions and Kernels

The following four theorems refer to extensions of a function (respectively a kernel), defined on a subset, to a function (respectively a kernel) on the whole set which obeys suitable restrictions.

Theorem 2.3.1. *Let $K(y, x)$ be a positive definite kernel on X and h a function on X_0 , where X_0 is a subset of X . If*

$$[h(y)\overline{h(x)}] \leq [K(y, x)] \text{ on } X_0, \quad (2.18)$$

then there is a function $\tilde{h} \in \mathcal{H}_K(X)$ such that

$$\|\tilde{h}\|_K \leq 1 \text{ and } \tilde{h}(x) = h(x) \text{ for } x \in X_0. \quad (2.19)$$

Proof. Let $K^{(0)}(y, x)$ be the restriction of $K(y, x)$ to X_0 . We know that $K^{(0)}(y, x)$ is positive definite because $K(y, x)$ is positive definite. By assumption, for h a function on X_0 , since the equation (2.18) is satisfied, then applying Theorem

2.1.10 with $\gamma = 1$, we have $h \in \mathcal{H}_{K^{(0)}}(X_0)$ and by the proof of Theorem 2.1.10, $\|h\|_{K^{(0)}} \leq \gamma = 1$. It follows by Theorem 2.2.1 that, for $\mathcal{H}_{K^{(0)}}(X_0)$ and $\mathcal{H}_K(X)$ are reproducing kernel Hilbert spaces we have

$$\mathcal{H}_{K^{(0)}}(X_0) = \{\tilde{h} \mid_{X_0} : \tilde{h} \in \mathcal{H}_K(X)\}$$

and

$$\|h\|_{K^{(0)}} = \min \{\|\tilde{h}\|_K : \tilde{h} \mid_{X_0} = h\} \text{ for } h \in \mathcal{H}_{K^{(0)}}(X_0),$$

equivalently,

$$\|h\|_{K^{(0)}} = \min \{\|\tilde{h}\|_K : \tilde{h}(x) = h(x), x \in X_0\} \text{ for } h \in \mathcal{H}_{K^{(0)}}(X_0) \text{ with } \|h\|_{K^{(0)}} \leq 1.$$

Hence,

$$\|\tilde{h}\|_K \leq 1. \quad \square$$

Theorem 2.3.2. *Let $K^{(i)}(y, x)$, ($i = 1, 2$) be positive definite kernels on X , h a function on $X_0 \subseteq X$.*

(i) *If*

$$[h(y)\overline{h(x)}] \leq [K^{(1)}(y, x) + K^{(2)}(y, x)] \text{ on } X_0,$$

then there are functions $\tilde{f} \in \mathcal{H}_{K^{(1)}}(X)$ and $\tilde{g} \in \mathcal{H}_{K^{(2)}}(X)$ such that

$$\|\tilde{f}\|_{K^{(1)}}^2 + \|\tilde{g}\|_{K^{(2)}}^2 \leq 1 \text{ and } h(x) = \tilde{f}(x) + \tilde{g}(x) \text{ for } x \in X_0. \quad (2.20)$$

(ii) *If*

$$[h(y)\overline{h(x)}] \leq [K^{(1)}(y, x) \cdot K^{(2)}(y, x)] \text{ on } X_0,$$

then there are sequences of functions $(f_n)_{n \geq 1} \subset \mathcal{H}_{K^{(1)}}(X)$ and $(g_n)_{n \geq 1} \subset \mathcal{H}_{K^{(2)}}(X)$ such that

$$\sum_1^\infty \|f_n\|_{K^{(1)}}^2 \|g_n\|_{K^{(2)}}^2 \leq 1 \text{ and } \sum_1^\infty f_n(x)g_n(x) = h(x) \text{ for } x \in X_0. \quad (2.21)$$

Proof. (i) For this part, consider the kernel $K(y, x) = K^{(1)}(y, x) + K^{(2)}(y, x)$ on X . We know that $K(y, x)$ is positive definite because $K^{(1)}$ and $K^{(2)}$ are positive definite .

Let $\mathcal{H}_{K^{(1)}}(X)$ and $\mathcal{H}_{K^{(2)}}(X)$ be associated reproducing kernel Hilbert spaces for $K^{(1)}$ and respectively $K^{(2)}$ their reproducing kernels. By Theorem 2.2.2, we have \mathcal{H}_K , the reproducing kernel Hilbert space admitting the reproducing kernel $K(y, x)$ such that $\mathcal{H}_K(X) = \mathcal{H}_{K^{(1)}}(X) + \mathcal{H}_{K^{(2)}}(X)$. By assumption, since

$$[h(y)\overline{h(x)}] \leq [K^{(1)}(y, x) + K^{(2)}(y, x)] = K(y, x) \text{ on } X_0,$$

applying previous theorem, there exists $\tilde{h} \in \mathcal{H}_K(x)$ such that $\|\tilde{h}\|_K \leq 1$ and $\tilde{h}(x) = h(x)$ for $x \in X_0$.

Returning to Theorem 2.2.2, there are functions $\tilde{f} \in \mathcal{H}_{K^{(1)}}(X)$ and $\tilde{g} \in \mathcal{H}_{K^{(2)}}(X)$ such that $\tilde{h} = \tilde{f} + \tilde{g}$ and

$$\|\tilde{h}\|_K^2 = \min \{ \|\tilde{f} + k\|_{K^{(1)}}^2 + \|\tilde{g} - k\|_{K^{(2)}}^2 : k \in \mathcal{H}_{K^{(1)}} \cap \mathcal{H}_{K^{(2)}} \} \leq 1.$$

Hence, we get

$$\|\tilde{h}\|_K^2 = \|\tilde{f}\|_{K^{(1)}}^2 + \|\tilde{g}\|_{K^{(2)}}^2 \leq 1 \quad \text{and} \quad \tilde{h}(x) = \tilde{f}(x) + \tilde{g}(x) \text{ for } x \in X_0.$$

(ii) Consider $K(y, x)$, the pointwise product of two kernels $K^{(1)}(y, x)$ and $K^{(2)}(y, x)$, that is $K(y, x) = K^{(1)}(y, x) \cdot K^{(2)}(y, x)$. Since both $K^{(1)}$ and $K^{(2)}$ are positive definite, then $K(y, x)$ is also positive definite, see Remark 2.2.4. Similarly as in (i), let $\mathcal{H}_{K^{(1)}}(X)$ and $\mathcal{H}_{K^{(2)}}(X)$ be the associated reproducing kernel Hilbert spaces for $K^{(1)}$ and $K^{(2)}$ their reproducing kernels, respectively. Denote by $\mathcal{H}_K(X)$ the reproducing kernel Hilbert space admitting the reproducing kernel $K(y, x)$.

Since, by assumption,

$$[h(y)\overline{h(x)}] \leq [K^{(1)}(y, x) \cdot K^{(2)}(y, x)] = K(y, x) \text{ on } X_0,$$

then there exists $\tilde{h} \in \mathcal{H}_K(X)$ such that $\|\tilde{h}\|_K \leq 1$ and $\tilde{h}(x) = h(x)$ for $x \in X_0$.

By Theorem 2.2.5, the reproducing kernel Hilbert space \mathcal{H}_K consists of all functions \tilde{h} on X such that there exist sequences $(f_n)_{n \geq 1} \subset \mathcal{H}_{K^{(1)}}(X)$ and $(g_n)_{n \geq 1} \subset \mathcal{H}_{K^{(2)}}(X)$ subject to the conditions:

$$\tilde{h}(x) = \sum_1^\infty f_n(x)g_n(x), \quad \text{for } x \in X_0$$

and

$$\sum_1^{\infty} \|f_n\|_{K^{(1)}}^2 \|g_n\|_{K^{(2)}}^2 < \infty$$

with

$$\|\tilde{h}\|_K^2 = \min \left\{ \sum_1^{\infty} \|f_n\|_{K^{(1)}}^2 \|g_n\|_{K^{(2)}}^2 \right\}.$$

But we know $\|\tilde{h}\|_K^2 \leq 1$. Hence we have

$$\sum_1^{\infty} \|f_n\|_{K^{(1)}}^2 \|g_n\|_{K^{(2)}}^2 \leq 1. \quad \square$$

As a result of the above theorem and Theorem 2.1.13, any positive definite kernel $K(y, x)$ on a subset X_0 of X can be extended to a positive definite kernel on the whole set X .

Let $K(y, x)$ be a positive definite kernel on X . For a continuous linear operator L on $\mathcal{H}_K(X)$, (i.e. $L : \mathcal{H}_K(X) \longrightarrow \mathcal{H}_K(X)$), we associate its kernel as

$$T(y, x) = \langle LK_x, K_y \rangle \text{ for } x, y \in X. \quad (2.22)$$

Recall that $\{K_x : x \in X\}$ is total in $\mathcal{H}_K(X)$. For any finitely supported $[\xi_x]$ and $[\eta_x]$, we have

$$\langle L(\sum_x \xi_x K_x), \sum_y \eta_y K_y \rangle = \sum_{x,y} \xi_x \bar{\eta}_y \langle LK_x, K_y \rangle = \sum_{x,y} \xi_x \bar{\eta}_y T(y, x).$$

Therefore, $\|L\| \leq 1$ is equivalent to the following condition:

$$\begin{aligned} \left| \sum_{x,y} \xi_x \bar{\eta}_y T(y, x) \right|^2 &= \left| \langle L(\sum_x \xi_x K_x), \sum_y \eta_y K_y \rangle \right|^2 \\ &\leq \left\| \sum_x \xi_x K_x \right\|^2 \left\| \sum_y \eta_y K_y \right\|^2 \\ &= \left(\sum_{x,y} \xi_x \bar{\xi}_y K(y, x) \right) \left(\sum_{x,y} \eta_x \bar{\eta}_y K(y, x) \right). \end{aligned}$$

Thus,

$$\left| \sum_{x,y} \xi_x \bar{\eta}_y T(y, x) \right|^2 \leq \left(\sum_{x,y} \xi_x \bar{\xi}_y K(y, x) \right) \left(\sum_{x,y} \eta_x \bar{\eta}_y K(y, x) \right). \quad (2.23)$$

This last condition (2.23) can be expressed as the positive definiteness of the kernel \tilde{K} on the disjoint union of X and its copy X' , where \tilde{K} is defined as

$$\tilde{K}(y, x) = \begin{cases} K(y, x) & \text{if } x, y \in X \\ T(y, x') & \text{if } x \in X', y \in X \\ \overline{T(x, y')} & \text{if } x \in X, y \in X' \\ K(y', x') & \text{if } x, y \in X' \end{cases}$$

where the canonical identification map $X' \rightarrow X$ is denoted by prime and the positive definiteness will be denoted by

$$\begin{bmatrix} [K(y, x)] & [T(y, x)] \\ [\overline{T(x, y)}] & [K(y, x)] \end{bmatrix} \geq 0 \text{ on } X. \quad (2.24)$$

Conversely, if a kernel $T(y, x)$ is given on X and satisfies one of the equations (2.23) or (2.24), then there is a continuous linear operator L on $\mathcal{H}_K(X)$ with $\|L\| \leq 1$ whose kernel coincides with T .

Lemma 2.3.3. *Let L be a bounded linear operator on \mathcal{H}_K and T the associated kernel as in (2.22). Then L is self-adjoint (respectively positive) if and only if the kernel $T(y, x)$ is Hermitian (respectively positive definite).*

Proof. L is self-adjoint if and only if $L = L^*$, equivalently for all $x, y \in X$,

$$T(y, x) = \langle LK_x, K_y \rangle = \langle L^*K_x, K_y \rangle = \overline{\langle K_y, L^*K_x \rangle} = \overline{T(x, y)}.$$

Moreover, L is positive if and only if

$$\langle Lf, f \rangle \geq 0, \text{ for all } f \in \mathcal{H}_K,$$

equivalently, for any finitely supported family of complex numbers $\{\epsilon_x\}_{x \in X}$,

$$\sum_{x, y} \bar{\epsilon}_y \epsilon_x L(y, x) \geq 0. \quad \square$$

Theorem 2.3.4. *If a kernel $T(y, x)$ on X_0 satisfies*

$$\begin{bmatrix} [K(y, x)] & [T(y, x)] \\ [\overline{T(y, x)}] & [K(y, x)] \end{bmatrix} \geq 0 \text{ on } X_0,$$

then there exists a kernel $\tilde{T}(y, x)$ on X such that $T(y, x) = \tilde{T}(y, x)$ for $x, y \in X_0$ and

$$\begin{bmatrix} [K(y, x)] & [T(y, x)] \\ [\overline{T(y, x)}] & [K(y, x)] \end{bmatrix} \text{ on } X.$$

If $T(y, x)$ is Hermitian (respectively positive definite), then $\tilde{T}(y, x)$ can be chosen Hermitian (respectively positive definite).

Proof. Let $K^{(0)}(y, x)$ be the restriction of $K(y, x)$ to X_0 . Consider the reproducing kernel Hilbert space $\mathcal{H}_{K^{(0)}}(X_0)$. Let $T(y, x)$ be a kernel on X_0 satisfying

$$\begin{bmatrix} [K(y, x)] & [T(y, x)] \\ [\overline{T(y, x)}] & [K(y, x)] \end{bmatrix} \geq 0 \text{ on } X_0.$$

This implies that there is a continuous linear operator L on $\mathcal{H}_{K^{(0)}}(X_0)$ such that $\|L\| \leq 1$ and $T(y, x) = \langle LK_x^{(0)}, K_y^{(0)} \rangle$ for $x, y \in X_0$. Then there is an isomorphism S from \mathcal{M} , the closed linear span of $\{K_x : x \in X_0\}$, onto $\mathcal{H}_{K^{(0)}}(X_0)$ such that $SK_x = K_x^{(0)}$ for $x \in X_0$. Denote by $P_{\mathcal{M}}$ the orthogonal projection onto \mathcal{M} . Define

$$\tilde{T}(y, x) := \langle LSP_{\mathcal{M}}K_x, SP_{\mathcal{M}}K_y \rangle \text{ for } x, y \in X.$$

Then

$$\tilde{T}(y, x) := \langle S^*LSP_{\mathcal{M}}K_x, P_{\mathcal{M}}K_y \rangle = \langle P_{\mathcal{M}}S^*LSP_{\mathcal{M}}K_x, K_y \rangle \text{ for all } x, y \in X.$$

This means that $\tilde{T}(y, x)$ becomes the kernel of the continuous linear operator $P_{\mathcal{M}}S^*LSP_{\mathcal{M}}$ on $\mathcal{H}_K(X)$. If we take the norm of this operator, by using the properties of orthogonal projections and isometric operators, we get

$$\begin{aligned} \|P_{\mathcal{M}}S^*LSP_{\mathcal{M}}\| &\leq \|P_{\mathcal{M}}\| \|S^*\| \|L\| \|S\| \|P_{\mathcal{M}}\| \\ &\leq \|S^*\| \|L\| \|S\| \\ &\leq \|L\| \leq 1. \end{aligned}$$

Then for any $x, y \in X_0$,

$$\tilde{T}(y, x) = \langle LSP_{\mathcal{M}}K_x, SP_{\mathcal{M}}K_y \rangle = \langle LSK_x, SK_y \rangle$$

and using $SK_x = K_x^{(0)}$ for $x \in X_0$, we get

$$\langle LSK_x, SK_y \rangle = \langle LK_x^{(0)}, K_y^{(0)} \rangle = T(y, x)$$

and hence

$$T(y, x) = \tilde{T}(y, x).$$

Moreover, if $T(y, x)$ is Hermitian (respectively positive definite), then L^* is self-adjoint (respectively positive definite) by previous remark and so the operator $P_{\mathcal{M}}S^*LSP_{\mathcal{M}}$ will be self-adjoint (respectively positive definite). Denote $P_{\mathcal{M}}S^*LSP_{\mathcal{M}} = N$. Then

$$\begin{aligned} \tilde{T}(y, x) &= \langle P_{\mathcal{M}}S^*LSP_{\mathcal{M}}K_x, K_y \rangle = \langle NK_x, K_y \rangle \\ &= \langle N^*K_x, K_y \rangle = \langle K_x, NK_y \rangle = \overline{\langle NK_y, K_x \rangle} = \overline{\langle \tilde{T}(x, y) \rangle}. \end{aligned}$$

So $\tilde{T}(y, x)$ is Hermitian (respectively positive definite). \square

When L is self-adjoint, then $\|L\| \leq 1$ if and only if

$$|\langle Lf, f \rangle| \leq \|L\| \|f\|_K^2 \leq \|f\|_K^2 \quad \text{for all } f \in \mathcal{H}_K(X). \quad (2.25)$$

This implies

$$|\langle L(\sum_x \xi_x K_x), \sum_y \xi_y K_y \rangle| \leq \sum_{x,y} \bar{\xi}_y \xi_x K(y, x),$$

which is equivalent to

$$|\sum_{x,y} \bar{\xi}_y \xi_x M(y, x)| \leq \sum_{x,y} \bar{\xi}_y \xi_x K(y, x), \quad \text{for all } [\xi_x]. \quad (2.26)$$

When L is not self adjoint, then either of the equations (2.25) or equivalently (2.26) only imply $L \leq 2$.

Theorem 2.3.5. *Let $K(y, x)$ be a positive definite kernel on X , and let $L(y, x)$ be a kernel on X_0 . If for any finitely supported family of complex numbers $\{\xi_x\}_{x \in X}$*

$$|\sum_{x,y \in X_0} \bar{\xi}_y \xi_x L(y, x)| \leq \sum_{x,y \in X_0} \bar{\xi}_y \xi_x K(y, x), \quad (2.27)$$

then there is a kernel $\tilde{L}(y, x)$ on X such that $L(y, x) = \tilde{L}(y, x)$ for all $x, y \in X_0$ and

$$|\sum_{x,y} \bar{\xi}_y \xi_x \tilde{L}(y, x)| \leq \sum_{x,y} \bar{\xi}_y \xi_x K(y, x) \text{ for all } [\xi_x].$$

If $L(y, x)$ is Hermitian (respectively positive definite), $\tilde{L}(y, x)$ can be chosen Hermitian (respectively positive definite).

Proof. Let equation (2.27) be satisfied for $K(y, x)$ a positive definite kernel on X and let $L(y, x)$ be a kernel on X_0 . So this is equivalent with Theorem 2.3.4. This implies that the linear operator L on $\mathcal{H}_K^{(0)}(X_0)$ satisfies

$$|\langle Lh, h \rangle_{K^{(0)}}| \leq \|h\|_{K^{(0)}}^2 \text{ for } h \in \mathcal{H}_K^{(0)}(X_0).$$

Taking S isometric and choosing $\tilde{L}(y, x) = \langle LSP_M K_x, SP_M K_y \rangle$, similarly as in the previous proof, we have

$$|\langle LSP_M f, SP_M f \rangle_{K^{(0)}}| \leq \|SP_M f\|_{K^{(0)}}^2 \leq \|f\|_K^2 \text{ for } f \in \mathcal{H}_K(X).$$

Hence, similarly as in the previous proof, we find that $L(y, x) = \tilde{L}(y, x)$ for $x, y \in X_0$. Therefore,

$$\begin{aligned} \left| \sum_{x,y} \bar{\xi}_y \xi_x \tilde{L}(y, x) \right| &= \left| \sum_{x,y} \bar{\xi}_y \xi_x \langle P_M S^* LSP_M K_x, K_y \rangle \right| \\ &= \left| \langle P_M S^* LSP_M \left(\sum_x \xi_x K_x \right), \sum_y \xi_y K_y \rangle \right| \\ &\leq \|P_M S^* LSP_M\| \left\| \sum_x \xi_x K_x \right\|^2 \\ &\leq \sum_{x,y} \bar{\xi}_y \xi_x \langle K_x, K_y \rangle \\ &\leq \sum_{x,y} \bar{\xi}_y \xi_x K(y, x). \end{aligned} \quad \square$$

Chapter 3

Spaces of Analytic Functions

3.1 Sesqui-analytic kernels

Definition 3.1.1. A two variable function on a domain Ω in the complex plane, is *sesqui-analytic* if it is analytic in the first variable and anti-analytic in the second variable.

For example, this holds if the kernel $K(w, z)$ is analytic in the first variable and Hermitian, that is,

$$K(w, z) = K_z(w) = \overline{K_w(z)} = \overline{K(z, w)} \quad \text{for all } w, z \in \Omega.$$

Definition 3.1.2. A function f defined on some topological space X with real or complex values is called *locally bounded*, if for any x_0 in X , there exists a neighborhood A of x_0 such that $f(A)$ is a bounded set, that is, for some number $M > 0$, $|f(x)| \leq M$ for all x in A .

We have the kernel $K(w, z)$ is locally bounded in the sense that it is bounded on $A \times B$ for every pair $\{A, B\}$ of compact subsets of a domain Ω .

Let us denote by Ω a connected domain of the complex plane.

Theorem 3.1.3. *The reproducing kernel Hilbert space $\mathcal{H}_K(\Omega)$ consists of analytic functions on Ω if and only if the positive definite kernel $K(w, z)$ on Ω is sesqui-analytic and locally bounded.*

Proof. Suppose that $\mathcal{H}_K(\Omega)$ consists of analytic functions on Ω . Let $K(w, z)$ be the reproducing kernel of $\mathcal{H}_K(\Omega)$ and note that $K_z \in \mathcal{H}_K$ is analytic. By the definition of a reproducing kernel which is positive definite, we have

$$K(w, z) = \overline{K(z, w)} \quad \text{for all } w, z \in \Omega.$$

So K is sesqui-analytic. To see the local boundedness, consider any pair $\{A, B\}$ of compact subsets of Ω . By assumption, every $f \in \mathcal{H}_K(\Omega)$ is analytic. Then

$$f(z) = \langle f, K_z \rangle, \quad z \in \Omega,$$

is analytic and hence continuous in z . Then the map $z \mapsto K_z$ is weakly continuous. This implies $\{K_z : z \in A\}$ is weakly compact, thus weakly bounded. Then by the Theorem A.3.6, weakly boundedness implies strong boundedness. That is, $\sup_{z \in A} \|K_z\| =: \gamma_A < \infty$. Now by Schwarz Inequality,

$$|K(w, z)| = |\langle K_z, K_w \rangle| \leq \|K_z\| \cdot \|K_w\| \leq \gamma_A \cdot \gamma_B,$$

for $w \in A, z \in B$. Hence K is locally bounded.

Conversely, suppose that a positive definite kernel $K(w, z)$ on Ω is sesqui-analytic and locally bounded. Recall that the set $\{K_z : z \in \Omega\}$ is total in $\mathcal{H}_K(\Omega)$. Then for each $f \in \mathcal{H}_K(\Omega)$, f is the strong limit of a sequence $(f_n)_{n \geq 1}$ in the linear span of $\{K_z : z \in \Omega\}$, that is $\|f_n - f\| \rightarrow 0$, as $n \rightarrow \infty$. By assumption, since $K(w, z)$ is sesqui-analytic, then $K(w, z) = K_z(w)$ is analytic in w . Then by reproducing property, since

$$f_n(w) = \langle f_n, K_w \rangle, \quad w \in \Omega,$$

it follows that f_n is analytic. Since $K(w, z)$ is locally bounded, we have $\sup_{z \in A} \|K_z\| \equiv \gamma_A < \infty$, where A is any compact subset of Ω . Then for $w \in A$,

$$\begin{aligned} |f_n(w) - f(w)| &= |\langle f_n, K_w \rangle - \langle f, K_w \rangle| = |\langle f_n - f, K_w \rangle| \\ &= \|f_n - f\| \|K_w\| \leq \gamma_A \|f_n - f\| \end{aligned}$$

and since $\|f_n - f\| \rightarrow 0$, (as $n \rightarrow \infty$), we have f_n converges to f uniformly on each compact subset A of Ω . Hence, $\mathcal{H}_K(\Omega)$ consists of analytic functions on Ω . \square

Definition 3.1.4. A subset Λ of Ω is called *determining subset* if every analytic function on Ω , equal to zero on Λ , vanishes identically on Ω . In particular, if Λ has a limit point in Ω , it is a determining subset.

If two analytic functions are equal on a determining subset Λ , then they coincide on the whole set Ω , i.e. $f_1|_\Lambda = f_2|_\Lambda$ implies $f_1|_\Omega = f_2|_\Omega$.

In particular, given two sesqui-analytic kernels $K^{(1)}(w, z)$ and $K^{(2)}(w, z)$ on Ω , if $K^{(1)}(w, z) = K^{(2)}(w, z)$ for $w, z \in \Lambda$ and Λ is determining subset, then $K^{(1)}(w, z) = K^{(2)}(w, z)$ on whole Ω .

Theorem 3.1.5. *Let $K(w, z)$ be a locally bounded, sesqui-analytic and positive definite kernel on Ω and Λ a determining subset of Ω .*

(i) *If a function h on Λ satisfies the condition*

$$[h(w)\overline{h(z)}] \leq [K(w, z)] \text{ on } \Lambda, \quad (3.1)$$

then there exists uniquely an analytic function \tilde{h} on Ω such that

$$\tilde{h}(z) = h(z) \text{ for } z \in \Lambda \text{ and } [\tilde{h}(w)\overline{\tilde{h}(z)}] \leq [K(w, z)]. \quad (3.2)$$

(ii) *If a positive definite kernel $L(w, z)$ satisfies*

$$[L(w, z)] \leq [K(w, z)] \text{ on } \Lambda, \quad (3.3)$$

then there exists uniquely a sesqui-analytic positive definite kernel $\tilde{L}(w, z)$ on Ω such that

$$\tilde{L}(w, z) = L(w, z) \text{ for } w, z \in \Lambda \text{ and } [\tilde{L}(w, z)] \leq [K(w, z)] \text{ on } \Omega. \quad (3.4)$$

Proof. Let $K(w, z)$ be a locally bounded, sesqui-analytic and positive definite kernel on Ω and Λ a determining subset of Ω .

(i) Suppose that for a function h on Ω (3.1) is satisfied. Then by Theorem 2.3.1, there exists $\tilde{h} \in \mathcal{H}_K(\Omega)$ such that $\|\tilde{h}\|_K \leq 1$ and $\tilde{h}(z) = h(z)$ for $z \in \Lambda$. By the same theorem, we can extend $h(w)\overline{h(z)}$ on Λ to a positive definite kernel $\tilde{h}(w)\overline{\tilde{h}(z)}$ on Ω . Then $\tilde{h}(w)\overline{\tilde{h}(z)} = h(w)\overline{h(z)}$ on Λ implies that

$$[\tilde{h}(w)\overline{\tilde{h}(z)}] \leq [K(w, z)].$$

Note that by Theorem 3.1.3, \mathcal{H}_K consists of analytic functions and hence \tilde{h} is analytic as well. The uniqueness part follows due to the assumption on the set Λ to be determining for Ω .

(ii) Suppose that a positive definite kernel $L(y, x)$ satisfies the condition (3.3). By Theorem 2.1.13 there exists a Hilbert space \mathcal{H} and a function $h: \Lambda \rightarrow \mathcal{H}$ such that $L(w, z) = \langle h_z, h_w \rangle_{\mathcal{H}}$ for all $z, w \in \Lambda$. Then we use part (i) for Hilbert space valued functions. \square

In the successive theorem, we will use the following lemma:

Lemma 3.1.6. *Let Ω be a domain in the complex plane and f either analytic or harmonic in Ω . Then for all $w \in \Omega$ and $\epsilon > 0$ such that $D(w; \epsilon) := \{z \in \mathbb{C} : |z - w| \leq \epsilon\} \subset \Omega$, we have*

$$f(w) = \frac{1}{\pi\epsilon^2} \iint_{D(w; \epsilon)} f(z) dm(z)$$

where $m(\cdot)$ is the planar Lebesgue measure in \mathbb{C} .

Proof. By writing $f = u + iv$, it follows that it is sufficient to prove the statement for f harmonic on Ω . Also recall that by the Cauchy integral formula for harmonic functions, for all $r \in [0, \epsilon]$,

$$f(w) = \frac{1}{2\pi} \int_0^{2\pi} f(w + re^{it}) dt.$$

Then by using the change of variables to polar coordinates

$$x = a + r \cos t, \quad y = b + r \sin t,$$

where $z = x + iy$ and $w = a + ib$, we have

$$\begin{aligned}
\frac{1}{\pi\epsilon^2} \iint_{D(w,\epsilon)} f(z) dm(z) &= \frac{1}{\pi\epsilon^2} \int_0^\epsilon \int_0^{2\pi} f(w + re^{it}) r dt dr \\
&= \frac{1}{\pi\epsilon^2} \int_0^\epsilon \left(\int_0^{2\pi} f(w + re^{it}) dt \right) r dr \\
&= \frac{1}{\pi\epsilon^2} \int_0^\epsilon 2\pi f(w) r dr = \frac{2\pi f(w)}{\pi\epsilon^2} \cdot \frac{r^2}{2} \Big|_0^\epsilon \\
&= f(w). \quad \square
\end{aligned}$$

Theorem 3.1.7. *Let $K(w, z)$ be a locally bounded, sesqui-analytic kernel on Ω . If $K(w, z)$ is positive definite on a determining subset Λ , then it is on the whole Ω .*

Proof. Let $K(w, z)$ be a locally bounded, sesqui-analytic kernel on Ω . Suppose that $K(w, z)$ is positive definite on Λ . Our aim is to show that $K(w, z)$ is positive definite on the whole Ω . The proof is divided in six steps:

Step 1: $K(w, z)$ is Hermitian on Ω .

Since $K(w, z)$ is positive definite on Λ , then $K(w, z) = \overline{K(z, w)}$ on Λ . Then $\overline{K(z, w)}$ will also be sesqui-analytic on Λ . This implies that $K(w, z)$ and $\overline{K(z, w)}$ are equal on Ω . Hence $K(w, z)$ is Hermitian.

Step 2: *There exists a positive Borel function $\rho(z)$ on Ω which satisfies the following conditions:*

- (i) $1/\rho(z)$ is locally bounded.
- (ii) $\int_\Omega |K(w, z)|^2 \rho(z) dm(z) < \infty$ for all $w \in \Omega$.
- (iii) $\int_\Omega \int_\Omega |K(w, z)|^2 \rho(z) \rho(w) dm(z) dm(w) < \infty$

where $m(\cdot)$ denotes the planar Lebesgue measure.

Let us write Ω as an increasing union of bounded subdomains $\{\Omega_n\}_{n \geq 1}$ such that $\overline{\Omega_n} \subset \Omega_{n+1}$.

Let $\sup_{w,z \in \Omega_n} |K(w,z)|^2 = \gamma_n$, ($n = 1, 2, 3, \dots$). Since $K(w,z)$ is locally bounded, we have $\gamma_n < \infty$ for each n .

We define ρ as follow,

$$\rho(z) := \frac{1}{2^n(\gamma_n + 1)m(\Omega_n \setminus \Omega_{n-1})} \quad \text{for } z \in \Omega_n \setminus \Omega_{n-1} \quad \text{with } \Omega_0 = \emptyset. \quad (3.5)$$

(i) We have $1/\rho(z)$ is bounded on each Ω_n and every compact subset of Ω is absorbed in some Ω_m . So we get $1/\rho(z)$ is locally bounded.

(ii) Let $w \in \Omega_n \setminus \Omega_{n-1}$. Then

$$\begin{aligned} & \int_{\Omega} |K(w,z)|^2 \rho(z) dm(z) \\ &= \sum_{k=1}^{\infty} \int_{\Omega_k \setminus \Omega_{k-1}} |K(w,z)|^2 \rho(z) dm(z) \\ &\leq \sum_{k=1}^{\infty} \int_{\Omega_k \setminus \Omega_{k-1}} \gamma_k \rho(z) dm(z) \\ &= \sum_{k=1}^{\infty} \int_{\Omega_k \setminus \Omega_{k-1}} \gamma_k \frac{1}{2^k(\gamma_k + 1)m(\Omega_k \setminus \Omega_{k-1})} dm(z) \\ &= \sum_{k=1}^{\infty} \frac{\gamma_k}{2^k(\gamma_k + 1)m(\Omega_k \setminus \Omega_{k-1})} \int_{\Omega_k \setminus \Omega_{k-1}} dm(z) \\ &\leq \sum_{k=1}^{\infty} 1/2^k = 1 < \infty. \end{aligned}$$

(iii) To see this we have the following estimations:

$$\begin{aligned} & \int_{\Omega} \int_{\Omega} |K(w,z)|^2 \rho(z) \rho(w) dm(z) dm(w) \\ &= \int_{\Omega} \rho(w) \left(\int_{\Omega} |K(w,z)|^2 \rho(z) dm(z) \right) dm(w) \\ &= \sum_{k=1}^{\infty} \int_{\Omega_k \setminus \Omega_{k-1}} \rho(w) \left(\int_{\Omega} |K(w,z)|^2 \rho(z) dm(z) \right) dm(w) \\ &\leq \sum_{k=1}^{\infty} \int_{\Omega_k \setminus \Omega_{k-1}} \rho(w) dm(w) \quad (\text{by (ii)}) \\ &= \sum_{k=1}^{\infty} \frac{1}{2^k(\gamma_k + 1)} \leq 1 < \infty. \end{aligned}$$

Step 3: Define a new measure $d\mu(z) := \rho(z)dm(z)$ on Ω and let $L^2(\Omega, \mu)$ be the associated Hilbert space. Let $A^2(\Omega, \mu)$ be the subspace of $L^2(\Omega, \mu)$ consisting of all analytic functions in $L^2(\Omega, \mu)$. In the following we show that $A^2(\Omega, \mu)$ is a reproducing kernel Hilbert space on Ω and it is closed in $L^2(\Omega, \mu)$.

Let us fix $w \in \Omega$ and take $\varepsilon > 0$ such that the open disk $D(w, \varepsilon) := \{z : |z - w| \leq \varepsilon\}$ is contained in Ω . For any analytic function $f \in A^2(\Omega, \mu)$, according to the previous lemma,

$$f(w) = \frac{1}{\pi\varepsilon^2} \int_{D(w, \varepsilon)} f(z)dm(z).$$

Then by the Schwarz Inequality in $L^2(\Omega, \mu)$,

$$\begin{aligned} |f(w)| &= \left| \frac{1}{\pi\varepsilon^2} \int_{D(w, \varepsilon)} f(z)dm(z) \right| \\ &= \frac{1}{\pi\varepsilon^2} \left(\int_{\Omega} |f(z)|^2 \rho(z)dm(z) \right)^{1/2} \left(\int_{D(w, \varepsilon)} 1/\rho(z)dm(z) \right)^{1/2} \\ &\leq \kappa \|f\| \end{aligned}$$

where κ is a finite constant depending on w and ε , but not on $f \in A^2(\Omega, \mu)$. Thus, the point evaluation functional $f \mapsto f(w)$ is continuous on $L^2(\Omega, \mu)$ and hence $A^2(\Omega, \mu)$ is a reproducing kernel Hilbert space on Ω .

To see that $A^2(\Omega, \mu)$ is closed, by the above discussion, since $|f(w)| \leq \kappa \|f\|$, the strong topology of $A^2(\Omega, \mu)$ is stronger than the topology of local uniform convergence. This implies that the closure of $A^2(\Omega, \mu)$ consists of analytic functions, that is, $A^2(\Omega, \mu)$ is closed in $L^2(\Omega, \mu)$.

Step 4: Define a linear operator \mathbf{K} in $L^2(\Omega, \mu)$ such that

$$(\mathbf{K}f)(w) := \langle f, K_w \rangle = \int_{\Omega} K(w, z)f(z)d\mu(z) \quad (3.6)$$

for all $f \in L^2(\Omega, \mu)$ and $w \in \Omega$. We claim that it is unique and bounded.

Note that by (ii), $K_w(z) = K(z, w) = \overline{K(w, z)} \in L^2(\Omega, \mu)$ and hence \mathbf{K} is

well-defined. Then by using (iii) and (3.6),

$$\begin{aligned}
\int_{\Omega} |(\mathbf{K}f)(w)|^2 d\mu(w) &= \int_{\Omega} |\langle f, K_w \rangle|^2 d\mu(w) \\
&\leq \int_{\Omega} \|f\|^2 \|K_w\|^2 d\mu(w) \\
&= \|f\|^2 \int_{\Omega} \langle K_w, K_w \rangle d\mu(w) \\
&= \|f\|^2 \int_{\Omega} K(w, z) K_w(z) \rho(z) dm(z) \rho(w) dm(w) \\
&= \|f\|^2 \int_{\Omega} \int_{\Omega} |K(w, z)|^2 \rho(z) \rho(w) dm(z) dm(w) \\
&\leq C \|f\|^2
\end{aligned}$$

where C is a finite constant. Hence \mathbf{K} is bounded.

Step 5 : \mathbf{K} maps $L^2(\Omega, \mu)$ in $A^2(\Omega, \mu)$.

Note that since the kernel K is Hermitian, it follows that the operator \mathbf{K} is self-adjoint. Since $A^2(\Omega, \mu)$ is a closed subspace and all analytic functions in $L^2(\Omega, \mu)$ are contained in $A^2(\Omega, \mu)$ and for all w , $K_w(z)$ is analytic in z , it follows that $K_w \in A^2(\Omega, \mu)$. By Step 4, we have

$$\{K_w : w \in \Omega\}^{\perp} \subseteq \ker(\mathbf{K}) = (\text{Ran}\mathbf{K}^*)^{\perp} = (\text{Ran}\mathbf{K})^{\perp}, \quad \text{as } \mathbf{K} = \mathbf{K}^*.$$

Then $(\text{Ran}\mathbf{K}) = (\text{Ran}\mathbf{K}^*) \subset A^2(\Omega, \mu)$.

Step 6: Let $L(w, z)$ be the reproducing kernel of $A^2(\Omega, \mu)$. Then we have $\langle \mathbf{K}L_z, L_w \rangle = K(w, z)$ for all $w, z \in \Omega$.

Let P be the orthogonal projection onto $A^2(\Omega, \mu)$. Then for $f \in L^2(\Omega, \mu)$, we have $\langle \mathbf{K}f, f \rangle = \langle \mathbf{K}Pf, Pf \rangle$ since $\text{Ran}\mathbf{K} \subset A^2(\Omega, \mu)$ and \mathbf{K} is self adjoint.

In Step 4, put L_z instead of f with the reproducing property,

$$(\mathbf{K}L_z)(u) = \langle L_z, K_u \rangle = \overline{K_u(z)} = \overline{K(z, u)} = K(u, z) = K_z(u) \quad (3.7)$$

and

$$\langle \mathbf{K}L_z, L_w \rangle = \langle K_z, L_w \rangle = \overline{\langle L_w, K_z \rangle} = \overline{K_w(z)} = K_z(w) = K(w, z) \quad (3.8)$$

for $w, z \in \Omega$.

Since Λ is a determining subset of Ω and $\{L_z : z \in \Omega\}$ is total in $A^2(\Omega, \mu)$, it follows that $\{L_z : z \in \Lambda\}$ is total in $A^2(\Omega, \mu)$. Therefore, by (3.8) and taking into account that the linear operator \mathbf{K} is bounded and that the kernel K is positive on Λ , it follows that the operator \mathbf{K} is positive and hence the kernel K is positive definite on Ω . \square

Theorem 3.1.8. *Suppose that $\Omega^{(1)} \cap \Omega^{(2)} \neq \emptyset$ where $\Omega^{(1)}$ and $\Omega^{(2)}$ are connected domains of the complex plane. If $K^{(1)}(w, z)$ and $K^{(2)}(w, z)$ are locally bounded, sesqui-analytic, positive definite kernels on $\Omega^{(1)}$ and $\Omega^{(2)}$ respectively, such that*

$$K^{(1)}(w, z) = K^{(2)}(w, z) \text{ for all } w, z \in \Omega^{(1)} \cap \Omega^{(2)}, \quad (3.9)$$

then there exists uniquely a locally bounded, sesqui-analytic, positive definite kernel $\tilde{K}(w, z)$ on $\Omega^{(1)} \cup \Omega^{(2)}$ such that

$$\tilde{K}(w, z) = K^{(i)}(w, z) \text{ for } w, z \in \Omega^{(i)}, \quad (i = 1, 2). \quad (3.10)$$

Proof. Let Ω be the intersection of $\Omega^{(1)}$ and $\Omega^{(2)}$ and $K(w, z)$ be the restriction of $K^{(1)}(w, z) = K^{(2)}(w, z)$ to this Ω for $w, z \in \Omega$. So we have

$$K(w, z) = K^{(1)}(w, z) = K^{(2)}(w, z) \text{ for } w, z \in \Omega.$$

Since Ω is open in connected $\Omega^{(i)}$, Ω is a determining subset of $\Omega^{(1)}$ and $\Omega^{(2)}$. Then, there exists an isometric operator $T^{(i)}$ from $\mathcal{H}_{K^{(i)}}(\Omega^{(i)})$ to $\mathcal{H}_K(\Omega)$ such that

$$T^{(i)}K_z^{(i)} = K_z \text{ for } z \in \Omega \quad (i = 1, 2).$$

Now we can define the kernel $\tilde{K}(w, z)$ on $\tilde{\Omega} := \Omega^{(1)} \cup \Omega^{(2)}$ by

$$\tilde{K}(w, z) := \langle T^{(i)}K_z^{(i)}, T^{(j)}K_w^{(j)} \rangle_K \text{ if } w \in \Omega^{(i)}, z \in \Omega^{(j)} \text{ and } i, j \in \{1, 2\}.$$

Then, since $K^{(1)}(w, z)$ and $K^{(2)}(w, z)$ are locally bounded and sesqui-analytic, so is $\tilde{K}(w, z)$.

Moreover by the isometric property of operator $T^{(i)}$, we get from the last equation that

$$\tilde{K}(w, z) = K^{(i)}(w, z) \text{ for } w, z \in \Omega^{(i)} \quad (i = 1, 2).$$

Finally, since $\tilde{K}(w, z)$ is locally bounded and sesqui-analytic on $\tilde{\Omega}$, and $\tilde{K}(w, z)$ is positive definite on the determining subset Ω of $\tilde{\Omega}$, by Theorem 3.1.7, $\tilde{K}(w, z)$ is positive definite on the whole $\tilde{\Omega}$. \square

3.2 Bergman Spaces

Definition 3.2.1. The space of all analytic functions $f(z)$ on Ω for which

$$\iint_{\Omega} |f(z)|^2 dx dy < \infty, \quad (z = x + iy)$$

is satisfied, is called the *Bergman space* on Ω and denoted by $A^2(\Omega)$.

Remark 3.2.2. $A^2(\Omega)$ is a reproducing kernel Hilbert space with respect to the inner product

$$\langle f, g \rangle \equiv \langle f, g \rangle_{\Omega} := \iint_{\Omega} f(z) \overline{g(z)} dx dy$$

and its kernel is called the *Bergman kernel* of Ω and denoted by $B^{(\Omega)}(w, z)$.

In the following we will calculate the Bergman kernel for any simply connected domain Ω . Consider the simplest case, that is $\Omega = \mathbb{D}$, where $\mathbb{D} := \{z : |z| < 1\}$. For this case the inner product is

$$\langle f, g \rangle = \int_0^1 \int_0^{2\pi} f(re^{i\theta}) \overline{g(re^{i\theta})} d\theta r dr, \quad (z = re^{i\theta}, r \leq 1).$$

Theorem 3.2.3. The Bergman kernel for the open unit disc \mathbb{D} is given by

$$B^{(\mathbb{D})}(w, z) = \frac{1}{\pi} \frac{1}{(1 - w\bar{z})^2} \quad \text{for } w, z \in \mathbb{D}. \quad (3.11)$$

Proof. We divide the proof in three steps.

Step 1: $f_n(z) = \sqrt{\frac{n+1}{\pi}} z^n$, ($n=0,1,2,\dots$) form an orthonormal sequence in $A^2(\mathbb{D})$.

Since

$$\langle z^n, \overline{z^m} \rangle = \int_0^1 \int_0^{2\pi} r^n e^{in\theta} r^m e^{-im\theta} d\theta r dr = \frac{2\pi}{n+m+2} \delta_{nm},$$

then

$$\langle f_n, f_m \rangle = \delta_{nm},$$

which means that $\{f_n(z)\}_{n \geq 0}$ forms an orthonormal sequence.

Step 2: $\{f_n(z)\}_{n \geq 0}$ is total in $A^2(\mathbb{D})$.

Consider any function $f \in A^2(\mathbb{D})$. By the definition of Bergman space, f is analytic in \mathbb{D} and so we have the Taylor series expansion of f as

$$f(z) = \sum_0^{\infty} a_n z^n.$$

Then we get

$$\langle f, f_n \rangle = \sqrt{\frac{\pi}{n+1}} a_n, \quad (n = 0, 1, 2, \dots).$$

Therefore, if f is orthogonal to all of f_n , that is,

$$\langle f, f_n \rangle = \sqrt{\frac{\pi}{n+1}} a_n = 0, \quad (n = 0, 1, 2, \dots)$$

then $a_n = 0$ for all $n \geq 0$. In other words, all Taylor coefficients of f vanish and this gives $f = 0$. Hence, $\{f_n\}_{n \geq 0}$ is total in $A^2(\mathbb{D})$.

Step 3: *The Bergman kernel for \mathbb{D} is*

$$B^{(\mathbb{D})}(w, z) = \sum_0^{\infty} f_n(w) \overline{f_n(z)}.$$

If this is true, then it has to satisfy the reproducing property. Let us check this fact,

$$\begin{aligned} \langle f, B_z^{(\mathbb{D})} \rangle &= \left\langle \sum_0^{\infty} a_n f_n(w), \sum_0^{\infty} f_n(w) \overline{f_n(z)} \right\rangle = \sum_0^{\infty} f_n(z) a_n \langle f_n, f_n \rangle \\ &= \sum_0^{\infty} f_n(z) a_n = f(z). \end{aligned}$$

Hence, by the uniqueness of the reproducing kernel we have

$$B^{(\mathbb{D})}(w, z) = \sum_0^{\infty} f_n(w) \overline{f_n(z)} \text{ for all } w, z \in \mathbb{D}.$$

It remains to insert the values of f_n and f_m into the above equation and get

$$\begin{aligned} B^{(\mathbb{D})}(w, z) &= \sum_{n=0}^{\infty} f_n(w) \overline{f_n(z)} = \sum_{n=0}^{\infty} \sqrt{\frac{n+1}{\pi}} w^n \sqrt{\frac{n+1}{\pi}} \overline{z^n} = \frac{1}{\pi} \sum_{n=0}^{\infty} (n+1) (w\overline{z})^n \\ &= \frac{1}{\pi} (1 + 2w\overline{z} + 3(w\overline{z})^2 + \dots) = \frac{1}{\pi} \frac{d}{d\xi} \left(\sum_{n=0}^{\infty} \xi^n \right) \Big|_{\xi=w\overline{z}} \\ &= \frac{1}{\pi} \frac{1}{(1-\xi)^2} \Big|_{\xi=w\overline{z}} = \frac{1}{\pi} \frac{1}{(1-w\overline{z})^2}. \end{aligned}$$

Hence the Bergman kernel for the open unit disc \mathbb{D} is

$$B^{(\mathbb{D})}(w, z) = \frac{1}{\pi} \frac{1}{(1-w\overline{z})^2} \text{ for } w, z \in \mathbb{D}. \quad \square$$

By the Riemann Mapping Theorem, each simply connected domain (which is not equal to \mathbb{C}) is mapped conformally onto the open unit disc. Hence we can find the Bergman kernel for an arbitrary simply connected domain Ω , in terms of the associated conformal mapping function. The proof of the following theorem includes the calculation of this kernel.

Theorem 3.2.4. *The Bergman kernel of a simply connected domain $\Omega (\neq \mathbb{C})$ is given by*

$$B^{(\Omega)}(w, z) = \frac{1}{\pi} \frac{\phi'(w) \overline{\phi'(z)}}{(1 - \phi(w) \overline{\phi(z)})^2} \text{ for } w, z \in \Omega, \quad (3.12)$$

where ϕ is any conformal mapping function from Ω onto \mathbb{D} .

Proof. Let $f \in A^2(\mathbb{D})$. Assign $f \mapsto Uf$, where U is the linear mapping on Ω defined by

$$(Uf)(z) = f(\phi(z)) \phi'(z) \text{ for } z \in \Omega. \quad (3.13)$$

Claim: $U : A^2(\mathbb{D}) \longrightarrow A^2(\Omega)$ is an isometric operator.

Since we have the Jacobian of ϕ as

$$|\phi'(z)|^2 = \det \begin{bmatrix} \frac{\partial}{\partial x} \operatorname{Re} \phi & \frac{\partial}{\partial y} \operatorname{Re} \phi \\ \frac{\partial}{\partial x} \operatorname{Im} \phi & \frac{\partial}{\partial y} \operatorname{Im} \phi \end{bmatrix} \text{ where } z = x + iy,$$

then by the formula of change of variables we have

$$\begin{aligned} \iint_{\Omega} |(Uf)(z)|^2 dx dy &= \iint_{\Omega} |f(\phi(z))|^2 |\phi'(z)|^2 dx dy, \quad (z = x + iy) \\ &= \iint_{\mathbb{D}} |f(w)|^2 du dv, \quad (w = u + iv, \phi(z) = w). \end{aligned}$$

Similarly, take $g \in A^2(\Omega)$ and assign $g \mapsto Vg$ where V is the linear mapping on Ω defined by

$$(Vg)(w) = g(\psi(w)) \cdot \psi'(w) \text{ for } w \in \mathbb{D} \quad (3.14)$$

and ψ is the inverse mapping of ϕ , i.e. $\psi(w) = z$. By similar arguments, we get that V is also isometric. Since $\phi(\psi(w)) = \phi(z) = w$ and $\psi(\phi(z)) = \psi(w) = z$, with (3.13) and (3.14) we get U and V are inverse to each other. Thus, since they are both isometric operators, so U is a unitary operator.

For the last part, let us fix $z \in \Omega$. Take $f \in A^2(\Omega)$. By using the reproducing property of Bergman kernels $B^{(\Omega)}$ and $B^{(\mathbb{D})}$,

$$f(\phi(z))\phi'(z) = (Uf)(z) = \langle Uf, B_z^{(\Omega)} \rangle_{\Omega} = \langle f, U^* B_z^{(\Omega)} \rangle_{\mathbb{D}}$$

and

$$f(\phi(z))\phi'(z) = \phi'(z) \langle f, B_{\phi(z)}^{(\mathbb{D})} \rangle_{\mathbb{D}} = \langle f, \overline{\phi'(z)} B_{\phi(z)}^{(\mathbb{D})} \rangle_{\mathbb{D}}.$$

Then, by combining these formulas we get

$$U^* B_z^{(\Omega)} = \overline{\phi'(z)} B_{\phi(z)}^{(\mathbb{D})},$$

or equivalently by using the property of U being unitary, we have

$$B_z^{(\Omega)} = U \overline{\phi'(z)} B_{\phi(z)}^{(\mathbb{D})}. \quad (3.15)$$

It remains to calculate $B^{(\Omega)}(w, z)$ by using (3.11) and (3.15) with unitarity of U ,

$$\begin{aligned} B^{(\Omega)}(w, z) &= \langle B_z^{(\Omega)}, B_w^{(\Omega)} \rangle_{\Omega} = \langle U \overline{\phi'(z)} B_{\phi(z)}^{(\mathbb{D})}, U \overline{\phi'(w)} B_{\phi(w)}^{(\mathbb{D})} \rangle_{\Omega} \\ &= \phi'(w) \cdot \overline{\phi'(z)} \cdot \langle B_{\phi(z)}^{(\mathbb{D})}, B_{\phi(w)}^{(\mathbb{D})} \rangle_{\mathbb{D}} \\ &= \frac{1}{\pi} \cdot \frac{\phi'(w) \overline{\phi'(z)}}{(1 - \phi(w)\phi(z))^2} \text{ for } w, z \in \Omega. \quad \square \end{aligned}$$

The following result is the converse of the previous theorem.

Lemma 3.2.5. *A conformal mapping from Ω to \mathbb{D} can be recaptured from the Bergman kernel of Ω .*

Proof. For fixed $z_0 \in \Omega$, by the Riemann Mapping Theorem, there is a unique analytic function $w = \phi(z)$ mapping Ω onto the unit disc \mathbb{D} such that

$$\phi(z_0) = 0 \text{ and } \phi'(z_0) > 0. \quad (3.16)$$

By (3.12), for $w \in \Omega$ and $z_0 \in \Omega$ satisfying (3.16), we find

$$\phi'(w) = \pi \frac{B^{(\Omega)}(w, z_0)}{\phi'(z_0)}$$

and letting z_0 instead of w in the last equation, we get

$$\phi'(z_0) = \sqrt{\pi B^{(\Omega)}(z_0, z_0)}.$$

Then if we integrate this equation, we find the conformal mapping function in terms of Bergman kernel, i.e.

$$\phi(z) = \sqrt{\frac{\pi}{B^{(\Omega)}(z_0, z_0)}} \int_{z_0}^z B^{(\Omega)}(w, z_0) dw. \quad (3.17)$$

This completes the proof. \square

Definition 3.2.6. A *Jordan curve* is a continuous one-to-one image of $\{|\xi| = 1\}$ in \mathbb{C} .

Definition 3.2.7. A *Green's function* $G(w, z)$ of Ω is a function harmonic in Ω except at z , where it has logarithmic singularity, and continuous in the closure $\bar{\Omega}$, with boundary values $G(w, z) = 0$ for all $w \in \partial\Omega$, where Ω is a finitely connected domain of complex plane.

Suppose now that Ω is a finitely connected domain in the complex plane, bounded by analytic Jordan curves. $G(w, z)$ has a logarithmic singularity at z means that $G(w, z) - \log \frac{1}{|w-z|}$ is harmonic in a neighborhood of z . The symmetry relation $G(w, z) = G(z, w)$ is satisfied for the Green's function. Moreover, the

Green function is conformally invariant. That is, if $\phi(z) = w$ maps a domain \mathbb{D} conformally onto Ω , and if $G(w, z_0)$ is the Green's function of Ω , then $H(z, \zeta) = G(\phi(z), \phi(\zeta))$ is the Green's function of \mathbb{D} . For a simply connected domain Ω , the Green's function is $G(w, z) = -\log |\phi(w)|$, where ϕ maps Ω conformally onto \mathbb{D} and $\phi(z) = 0$. In particular, the Green function of \mathbb{D} is

$$G(w, z) = -\log \left| \frac{w - z}{1 - w\bar{z}} \right|. \quad (3.18)$$

The following theorem gives the Bergman kernel in terms of the Green's function for the general case.

Theorem 3.2.8. *Let Ω be a finitely connected domain bounded by analytic Jordan curves, and let $G(w, z)$ be the Green's function of Ω . Then the Bergman kernel function is*

$$B^{(\Omega)}(w, z) = -\frac{2}{\pi} \frac{\partial^2 G}{\partial w \partial \bar{z}}(w, z), \quad w \neq z. \quad (3.19)$$

Proof. By the definition of the Green's function, we have

$$G(w, z) = \log \frac{1}{|w - z|} + H(w, z)$$

in some neighborhood of z , where $H(w, z)$ is a harmonic function of w . Taking partial derivative with respect to w , we get

$$\frac{\partial G}{\partial w}(w, z) = -\frac{1}{2} \frac{1}{|w - z|} + \frac{\partial H}{\partial w}(w, z)$$

and now taking partial derivative with respect to \bar{z} , since $\frac{\partial^2}{\partial w \partial \bar{z}} \left(-\frac{1}{2} \frac{1}{|w - z|} \right) = 0$, then we get

$$\frac{\partial^2 G}{\partial w \partial \bar{z}}(w, z) = \frac{\partial^2 H}{\partial w \partial \bar{z}}, \quad \text{for } w \neq z.$$

Since the boundary curves are analytic and the Green's function vanishes on the boundary, it has a harmonic extension across the boundary. Also, for each $z \in \Omega$, $\frac{\partial G}{\partial w}(w, z)$ is analytic in w . Then, $\frac{\partial^2 G}{\partial w \partial \bar{z}}(w, z)$ is bounded and analytic in $w \in \Omega$, so it belongs to the Bergman space $A^2(\Omega)$.

Recall that according to the *Cauchy – Green theorem* (see [5]), we have

$$\int_{\partial\Omega} F(z) dz = 2i \int_{\Omega} \frac{\partial F}{\partial \bar{z}} dA, \quad \text{for } F \in C^1(\bar{\Omega}).$$

Suppose f is analytic in Ω and continuous in $\bar{\Omega}$. Let Ω_ε be the domain inside Ω given by a small disc $|w - z| \leq \varepsilon$, and let \mathbb{T}_ε denote the boundary of this disc. Since the Green's function $G(w, z)$ vanishes on the boundary, then $\frac{\partial G}{\partial w}$ also vanishes for z on $\partial\Omega$, then by the Cauchy-Green formula

$$\frac{1}{2i} \int_{\mathbb{T}_\varepsilon} \frac{\partial G}{\partial w}(w, z) f(z) dz = - \iint_{\Omega_\varepsilon} \frac{\partial^2 G}{\partial w \partial \bar{z}}(w, z) f(z) dA(z), \quad (3.20)$$

where the orientation of \mathbb{T}_ε is counterclockwise. However, if we apply the Cauchy Theorem to the left hand side of the above equation,

$$\int_{\mathbb{T}_\varepsilon} \frac{\partial G}{\partial w}(w, z) f(z) dz = \int_{\mathbb{T}_\varepsilon} \left(-\frac{1}{2} \frac{1}{|w - z|} + \frac{\partial H}{\partial w} \right) f(z) dz \longrightarrow \pi i f(w) \text{ as } \varepsilon \rightarrow 0.$$

Hence, applying this result to (3.20) we obtain

$$\frac{1}{2i} \pi i f(w) = - \iint_{\Omega} \frac{\partial^2 G}{\partial w \partial \bar{z}}(w, z) f(z) dA(z).$$

Thus we get

$$f(w) = -\frac{2}{\pi} \iint_{\Omega} \frac{\partial^2 G}{\partial w \partial \bar{z}}(w, z) f(z) dA(z),$$

where f is analytic in Ω and continuous in $\bar{\Omega}$.

Finally, since Ω has analytic boundary, then the kernel function of Ω has an analytic continuation across the boundary. Applying this to the function $f(w) = B^{(\Omega)}(w, \zeta)$ we get,

$$-\frac{2}{\pi} \iint_{\Omega} \frac{\partial^2 G}{\partial w \partial \bar{z}}(w, z) B^{(\Omega)}(z, \zeta) dA(z) = B^{(\Omega)}(w, \zeta).$$

Therefore, by the reproducing property of the kernel function, we get the desired result, i.e.

$$-\frac{2}{\pi} \iint_{\Omega} \frac{\partial^2 G}{\partial w \partial \bar{z}}(w, z) B^{(\Omega)}(z, \zeta) dA(z) = -\frac{2}{\pi} \frac{\partial^2 G}{\partial w \partial \bar{\zeta}}(w, \zeta) = B^{(\Omega)}(w, \zeta). \quad \square$$

3.3 Szegő Kernel

Consider the kernels $K_\alpha(w, z)$ on the open unit disc \mathbb{D} , ($0 < \alpha < \infty$),

$$K_\alpha(w, z) := \frac{1}{(1 - w\bar{z})^\alpha} \text{ for } w, z \in \mathbb{D}, \quad (3.21)$$

where ξ^α is the analytic continuation of t^α on \mathbb{R}_+ to the open half plane $\{\xi : \operatorname{Re}\xi > 0\}$. By the above definition of kernels, if we have $\alpha = 2$, we get π times Bergman kernel for \mathbb{D} , i.e.

$$K_2(w, z) = \frac{1}{(1 - w\bar{z})^2} = \pi B^{(\mathbb{D})}(w, z).$$

Since $K_\alpha(w, z) = \overline{K_\alpha(z, w)}$, it is sesqui-analytic and moreover it is locally bounded.

Lemma 3.3.1. *The kernel K_α defined at (3.21) is positive definite.*

Proof. Since

$$\frac{1}{(1 - \xi)^2} = \frac{d}{d\xi} \left(\sum_{n=0}^{\infty} \xi^n \right), \quad |\xi| < 1,$$

if we take derivatives of both sides α times, it follows that

$$\begin{aligned} \frac{1}{(1 - \xi)^\alpha} &= \frac{1}{(\alpha - 1)!} \cdot \frac{d^{\alpha-1}}{d\xi^{\alpha-1}} \cdot \left(\sum_{n=0}^{\infty} \xi^n \right) \\ &= \frac{1}{(\alpha - 1)!} \sum_{n=0}^{\infty} n(n-1)(n-2)(n-3) \cdots (n - (\alpha - 2)) \xi^{n-(\alpha-1)} \\ &= \frac{1}{(\alpha - 1)!} \sum_{N=0}^{\infty} (N + \alpha - 1)(N + \alpha - 2) \cdots (N + 1) \xi^N. \end{aligned}$$

Let Γ be the *Gamma function*. For $(\alpha = 1, 2, \dots)$, we have $\Gamma(\alpha) = (\alpha - 1)!$ and

$$\frac{\Gamma(n + \alpha)}{\Gamma(n + 1)} = (n + \alpha - 1)(n + \alpha - 2) \cdots (n + 1).$$

If we put $\xi = w\bar{z}$ into the above equation we obtain

$$\sum_{n=0}^{\infty} \frac{\Gamma(n + \alpha)}{\Gamma(\alpha)\Gamma(n + 1)} (w\bar{z})^n = \frac{1}{(1 - w\bar{z})^\alpha} \quad \text{for } w, z \in \mathbb{D}, \quad (\alpha = 1, 2, \dots).$$

We can generalize this result for each $0 < \alpha < \infty$, so that

$$\sum_{n=0}^{\infty} \frac{\Gamma(n + \alpha)}{\Gamma(\alpha)\Gamma(n + 1)} (w\bar{z})^n = \frac{1}{(1 - w\bar{z})^\alpha} = K_\alpha(w, z) \quad \text{for } w, z \in \mathbb{D}. \quad (3.22)$$

Therefore, since $[w\bar{z}^n]$ is positive definite for each n and

$$\frac{\Gamma(n + \alpha)}{\Gamma(\alpha)\Gamma(n + 1)} > 0, \quad (n = 0, 1, 2, \dots),$$

then $K_\alpha(w, z)$ is positive definite. □

Theorem 3.3.2. (i) *The Hilbert space $\mathcal{H}_{K_\alpha}(\mathbb{D})$ coincides with the space of analytic functions $f(z) = \sum_0^\infty a_n z^n$ on \mathbb{D} such that*

$$\sum_{n=0}^{\infty} \frac{\Gamma(\alpha)\Gamma(n+1)}{\Gamma(n+\alpha)} |a_n|^2 < \infty,$$

equipped with the inner product

$$\langle g, h \rangle = \sum_0^{\infty} \frac{\Gamma(\alpha)\Gamma(n+1)}{\Gamma(n+\alpha)} b_n \bar{c}_n, \quad (3.23)$$

for $g(z) = \sum_0^\infty b_n z^n$ and $h(z) = \sum_0^\infty c_n z^n$.

(ii) *When $\alpha > 1$, $\mathcal{H}_{K_\alpha}(\mathbb{D})$ also coincides with the space of analytic functions $f(z) = \sum_{n=0}^\infty a_n z^n$ such that*

$$\iint_{\mathbb{D}} (1 - |z|^2)^{\alpha-2} |f(z)|^2 dx dy < \infty, \quad (z = x + iy)$$

equipped with the inner product

$$\langle g, h \rangle = \frac{\alpha-1}{\pi} \iint_{\mathbb{D}} (1 - |z|^2)^{\alpha-2} |g(z)| \overline{|h(z)|} dx dy. \quad (3.24)$$

Proof. (i) Since $f(z) = \sum_0^\infty a_n z^n$, then

$$\begin{aligned} |f(z)| &= \left| \sum_{n=0}^{\infty} a_n z^n \right| \leq \sum_{n=0}^{\infty} |a_n| |z|^n \\ &\leq \left(\sum_{n=0}^{\infty} \frac{\Gamma(\alpha)\Gamma(n+1)}{\Gamma(n+\alpha)} |a_n|^2 \right)^{1/2} \left(\sum_{n=0}^{\infty} \frac{\Gamma(n+\alpha)}{\Gamma(\alpha)\Gamma(n+1)} |z|^{2n} \right)^{1/2}, \end{aligned}$$

and with the inner product (3.23), strong topology is stronger than the topology of local uniform convergence on \mathbb{D} . Hence this space of analytic functions becomes a reproducing kernel Hilbert space on \mathbb{D} .

The construction of the kernel will be similar as the construction of Bergman kernel.

Step 1: $f_n(z) := \sqrt{\frac{\Gamma(\alpha+n)}{\Gamma(\alpha)\Gamma(n+1)}} z^n$, ($n = 0, 1, 2, \dots$), *form an orthonormal sequence.*

$$\begin{aligned} \langle f_n, f_m \rangle &= \left\langle \sqrt{\frac{\Gamma(\alpha+n)}{\Gamma(\alpha)\Gamma(n+1)}} z^n, \sqrt{\frac{\Gamma(\alpha+m)}{\Gamma(\alpha)\Gamma(m+1)}} z^m \right\rangle \\ &= \delta_{nm}, \quad \text{for } n, m = 0, 1, \dots \end{aligned}$$

Thus, $\{f_n\}_{n \geq 0}$ is an orthonormal sequence.

Step 2: $\{f_n\}_{n \geq 0}$ is complete.

Consider the inner product in (3.23) of $f_n(z)$ with $f = \sum_{n=0}^{\infty} a_n z^n$,

$$\langle f, f_n \rangle = a_n \sqrt{\frac{\Gamma(\alpha)\Gamma(n+1)}{\Gamma(n+\alpha)}}, \quad (n = 0, 1, 2, \dots).$$

Therefore, f is orthogonal to all of f_n , that is

$$\langle f, f_n \rangle = a_n \sqrt{\frac{\Gamma(\alpha)\Gamma(n+1)}{\Gamma(n+\alpha)}} = 0, \quad (n = 0, 1, 2, \dots)$$

only if $a_n = 0$ for each n . Thus $f = 0$ and f_n is total.

Step 3: $K_\alpha(w, z) = \sum_{n=0}^{\infty} f_n(w) \overline{f_n(z)}$.

To prove this, we verify the reproducing property:

$$\begin{aligned} \langle f, K_z \rangle &= \left\langle \sum_{n=0}^{\infty} a_n f_n(w), \sum_{n=0}^{\infty} f_n(w) f_n(z) \right\rangle \\ &= \sum_{n=0}^{\infty} a_n f_n(z) = f(z). \end{aligned}$$

Hence, $K_\alpha(w, z) = \sum_{n=0}^{\infty} f_n(w) \overline{f_n(z)}$.

Finally, inserting the values of $f_n(w)$ and $f_n(z)$ in this equation, we get

$$\begin{aligned} K_\alpha(w, z) &= \sum_{n=0}^{\infty} f_n(w) \overline{f_n(z)} = \sum_{n=0}^{\infty} \sqrt{\frac{\Gamma(\alpha+n)}{\Gamma(\alpha)\Gamma(n+1)}} w^n \sqrt{\frac{\Gamma(\alpha+n)}{\Gamma(\alpha)\Gamma(n+1)}} \bar{z}^n \\ &= \sum_{n=0}^{\infty} \frac{\Gamma(\alpha+n)}{\Gamma(\alpha)\Gamma(n+1)} (w\bar{z})^n = \frac{1}{(1-w\bar{z})^\alpha}. \end{aligned}$$

(ii) We change the coordinates to polar coordinates. For $f = \sum_{n=0}^{\infty} a_n z^n$,

consider the following integration,

$$\begin{aligned}
& \frac{\alpha - 1}{\pi} \iint_{\mathbb{D}} (1 - |z|^2)^{\alpha-2} |f(z)|^2 dx dy \\
&= \frac{\alpha - 1}{\pi} \int_0^1 \int_0^{2\pi} (1 - r^2)^{\alpha-2} \left| \sum_{n=0}^{\infty} a_n (r e^{i\theta})^n \right|^2 d\theta r dr \quad (z = r e^{i\theta}) \\
&= \frac{\alpha - 1}{\pi} \int_0^1 \int_0^{2\pi} (1 - r^2)^{\alpha-2} \left(\sum_{n,m \geq 0} a_m \bar{a}_n r^m r^n e^{i\theta(m-n)} \right) d\theta r dr \\
&= \frac{\alpha - 1}{\pi} \sum_{n,m \geq 0} a_m \bar{a}_n \int_0^1 \int_0^{2\pi} (1 - r^2)^{\alpha-2} r^{m+n+1} e^{i\theta(m-n)} d\theta r dr \\
&= \frac{\alpha - 1}{\pi} \sum_{n=0}^{\infty} |a_n|^2 \int_0^1 2\pi (1 - r^2)^{\alpha-2} r^{2n+1} d\theta r dr \\
&= 2(\alpha - 1) \sum_{n=0}^{\infty} |a_n|^2 \int_0^1 2\pi (1 - r^2)^{\alpha-2} r^{2n+1} d\theta r dr \quad (t = r^2, dt = 2r dr) \\
&= (\alpha - 1) \sum_{n=0}^{\infty} |a_n|^2 \int_0^1 (1 - t)^{\alpha-2} t^n dt.
\end{aligned}$$

Since

$$\int_0^1 (1 - t)^{\beta-1} t^{\gamma-1} dt = \frac{\Gamma(\beta)\Gamma(\gamma)}{\Gamma(\beta + \gamma)}, \quad \text{for } \beta, \gamma > 0,$$

then

$$(\alpha - 1) \int_0^1 (1 - t)^{\alpha-2} t^n dt = (\alpha - 1) \frac{\Gamma(\alpha - 1)\Gamma(n + 1)}{\Gamma(n + \alpha)} = \frac{\Gamma(\alpha)\Gamma(n + 1)}{\Gamma(n + \alpha)}.$$

Inserting this in the above integration, we get

$$(\alpha - 1) \sum_0^{\infty} |a_n|^2 \int_0^1 (1 - t)^{\alpha-2} t^n dt = \sum_0^{\infty} \frac{\Gamma(\alpha)\Gamma(n + 1)}{\Gamma(n + \alpha)} |a_n|^2$$

or equivalently,

$$\frac{\alpha - 1}{\pi} \iint_{\mathbb{D}} (1 - |z|^2)^{\alpha-2} |f(z)|^2 dx dy = \sum_0^{\infty} \frac{\Gamma(\alpha)\Gamma(n + 1)}{\Gamma(n + \alpha)} |a_n|^2.$$

Hence, the last equality shows that on the space of analytic functions, the two inner products (3.23) and (3.24) are the same. \square

According to the values that α takes, the behavior of $\mathcal{H}_{K_\alpha}(\mathbb{D})$ changes. If $\alpha > 1$, $\mathcal{H}_{K_\alpha}(\mathbb{D})$ becomes a Hilbert space of analytic functions $f(z)$ on \mathbb{D} such that

$$\int_{\mathbb{D}} |f(z)|^2 d\mu_\alpha(z) < \infty,$$

where μ_α is the measure on \mathbb{D} given by

$$d\mu_\alpha(z) = \frac{\alpha - 1}{\pi} (1 - |z|^2)^{\alpha-2} dx dy, \quad (z = x + iy).$$

Definition 3.3.3. For $\alpha > 1$, $\mathcal{H}_{K_\alpha}(\mathbb{D})$ is called *the weighted Bergman space* on \mathbb{D} (with weight $\frac{\alpha-1}{\pi}(1 - |z|^2)^{\alpha-2}$).

Lemma 3.3.4. *If $\alpha < 1$, $\mathcal{H}_{K_\alpha}(\mathbb{D})$ is not a reproducing kernel Hilbert space with respect to the inner product (3.24).*

Proof. Suppose that for $\alpha < 1$, $\mathcal{H}_{K_\alpha}(\mathbb{D})$ is a reproducing kernel space. Let $\phi(z)$ be an analytic function such that $|\phi(z)| \leq 1$ for $z \in \mathbb{D}$. Then $f \mapsto \phi \cdot f$ defines a continuous linear operator on $\mathcal{H}_{K_\alpha}(\mathbb{D})$ such that

$$\|\phi f\|_{K_\alpha} \leq \|f\|_{K_\alpha}, \quad \text{where } f \in \mathcal{H}_{K_\alpha}(\mathbb{D}). \quad (3.25)$$

Then, in terms of the kernel $K_\alpha(w, z)$, inequality (3.25) becomes

$$[\phi(w) K_\alpha(w, z) \overline{\phi(z)}] \leq [K_\alpha(w, z)] \text{ on } \mathbb{D},$$

and hence we get

$$[K_\alpha(w, z)(1 - \phi(w) \overline{\phi(z)})] \geq 0 \text{ on } \mathbb{D}.$$

This implies that, for any $z \in \mathbb{D}$, the following 2×2 matrix is positive definite

$$\begin{bmatrix} K_\alpha(0, 0)(1 - |\phi(0)|^2) & K_\alpha(0, z)(1 - \phi(0) \overline{\phi(z)}) \\ K_\alpha(z, 0)(1 - \phi(z) \overline{\phi(0)}) & K_\alpha(z, z)(1 - |\phi(z)|^2) \end{bmatrix} \geq 0. \quad (3.26)$$

If $\phi(0) = 0$, then (3.26) becomes

$$\begin{bmatrix} K_\alpha(0, 0) & K_\alpha(0, z) \\ K_\alpha(z, 0) & K_\alpha(z, z)(1 - |\phi(z)|^2) \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & \frac{1}{(1-|z|^2)^\alpha}(1 - |\phi(z)|^2) \end{bmatrix} \geq 0,$$

where $K_\alpha(w, z) = 1/(1 - w\bar{z})^\alpha$. This implies

$$\frac{1 - |\phi(z)|^2}{(1 - |z|^2)^\alpha} - 1 \geq 0 \quad \text{and} \quad 1 - (1 - |z|^2)^\alpha \geq |\phi(z)|^2.$$

Then, for $0 \neq z \in \mathbb{D}$,

$$\frac{1 - (1 - |z|^2)^\alpha}{|z|^2} \geq \frac{|\phi(z)|^2}{|z|^2}. \quad (3.27)$$

Now consider the analytic function $\phi_t(z) := z(z - t)/(1 - tz)$, ($0 < t < 1$), which satisfies $\phi_t(0) = 0$ and $|\phi_t(z)| \leq 1$ for $z \in \mathbb{D}$ and fix $z_0 \in \mathbb{D}$. Since we have $\alpha < 1$, then

$$(1 - |z_0|^2)^\alpha > 1 - |z_0|^2$$

and by this, we get the following

$$1 > \frac{1 - (1 - |z_0|^2)^\alpha}{|z_0|^2}. \quad (3.28)$$

Then, if we take the limit of $|\phi_t(z_0)/z_0|^2$ as $t \rightarrow 1$,

$$\lim_{t \rightarrow 1} |\phi_t(z_0)/z_0|^2 = \lim_{t \rightarrow 1} \left| \frac{z_0(z_0 - t)}{(1 - tz_0)z_0} \right|^2 = \lim_{t \rightarrow 1} \left| \frac{z_0 - t}{1 - tz_0} \right|^2 = 1.$$

But this contradicts with the inequality (3.27). Hence, for $\alpha < 1$, $\mathcal{H}_{K_\alpha}(\mathbb{D})$ is not a reproducing kernel Hilbert space with respect to the inner product (3.24). \square

It remains the case when $\alpha = 1$. Let us denote by \mathbb{T} the boundary of \mathbb{D} , i.e. $\mathbb{T} = \{\xi : |\xi| = 1\}$ and let σ be the normalized arc-length measure such that

$$d\sigma(\xi) := \frac{1}{2\pi} |d\xi| \equiv \frac{1}{2\pi} d\theta, \quad (\xi = e^{i\theta}).$$

Consider the Hilbert space $L^2(\mathbb{T}) \equiv L^2(\mathbb{T}, \sigma)$ of measurable functions on \mathbb{T} . We have the inner product

$$\langle f, g \rangle_{L^2} := \int_{\mathbb{T}} f(\xi) \overline{g(\xi)} d\sigma(\xi),$$

with respect to which the functions $\phi_n(\xi) := \xi^n$, ($n \in \mathbb{Z}$), which form a complete orthonormal sequence.

Definition 3.3.5. The closed linear span of $\{\phi_n : n = 0, 1, \dots\}$ is called the *Hardy space* on \mathbb{T} and is denoted by $H^2(\mathbb{T})$.

By Corollary A.2.3, $f \in L^2(\mathbb{T})$ belongs to Hardy space $H^2(\mathbb{T})$ if and only if it is orthonormal to all ϕ_n ($n < 0$), i.e. all Fourier coefficients of f which have negative indices vanish. Then we have

$$\langle f, g \rangle_{L^2} = \sum_{n=0}^{\infty} a_n \bar{b}_n \text{ for } f, g \in H^2(\mathbb{T}), \quad (3.29)$$

where

$$a_n = \langle f, \phi_n \rangle_{L^2} \text{ and } b_n = \langle g, \phi_n \rangle_{L^2} \quad (n = 0, 1, \dots).$$

Before stating the last theorem of this chapter, we will define the *Poisson kernel* and some important properties of this kernel.

Definition 3.3.6. For $z \in \mathbb{D}$ and $\xi \in \mathbb{T}$, $P_z(\xi)$ is called the *Poisson kernel* and defined by

$$P_z(\xi) := \frac{1 - |z|^2}{|1 - \xi \bar{z}|^2}.$$

Equivalently, for $z = re^{i\theta}$ ($0 \leq r < 1$) and $\xi = e^{it}$ ($-\infty < t < \infty$), the Poisson kernel can be written as

$$P_r(t) := \frac{1 - r^2}{1 - 2r \cos t + r^2} = \sum_{-\infty}^{\infty} r^{|n|} e^{int}.$$

We recall some properties of the Poisson kernel, (e.g. see [7]):

- (i) $P_r(t) \geq 0$, for all $r < 1$.
- (ii) $P_r(t + 2\pi) = P_r(t)$
- (iii) $\int_{-\pi}^{\pi} P_r(t) dt = 2\pi$, for all $r < 1$.
- (iv) Given $\delta > 0$, $P_r(t) \rightarrow 0$ uniformly for $\delta \leq |t| \leq \pi$ as $r \rightarrow 1$.

Theorem 3.3.7. *The correspondence*

$$f \longmapsto \tilde{f}(z) := \sum_{n=0}^{\infty} a_n z^n \quad \text{with } a_n = \langle f, \phi_n \rangle_{L^2}, \quad (n = 0, 1, \dots)$$

yields a unitary operator U from $H^2(\mathbb{T})$ onto $H_{K_1}(\mathbb{D})$. Conversely, f can be re-captured from $\tilde{f}(z)$ by the formula

$$f(\xi) = \lim_{r \rightarrow 1} \tilde{f}(r\xi) \text{ for almost all } \xi \in \mathbb{T}.$$

Proof. Put $\alpha = 1$ in $\frac{\Gamma(\alpha)\Gamma(n+1)}{\Gamma(n+\alpha)}$. We get $\frac{\Gamma(1)\Gamma(n+1)}{\Gamma(n+1)} = 1$.

Then (3.23) in Theorem 3.3.2 becomes

$$\langle g, h \rangle = \sum_0^{\infty} b_n \bar{c}_n, \text{ for } g(z) = \sum_0^{\infty} b_n z^n \text{ and } h(z) = \sum_0^{\infty} c_n z^n.$$

In other words, equation (3.23) becomes the same as the equation (3.29). This implies that the operator U from $H^2(\mathbb{T})$ onto $H_{K_1}(\mathbb{D})$, is isometric on the linear span of $\{\phi_n : n = 0, 1, \dots\}$. Since $\{\phi_n : n = 0, 1, \dots\}$ and $\{\tilde{\phi}_n : n = 0, 1, \dots\}$ are total in $H^2(\mathbb{T})$ and $H_{K_1}(\mathbb{D})$, respectively, then U is unitary. Now by (3.29),

$$\begin{aligned} \langle \tilde{f}(r\xi) - f(\xi), \tilde{f}(r\xi) - f(\xi) \rangle &= \int_{\mathbb{T}} |\tilde{f}(r\xi) - f(\xi)|^2 d\sigma(\xi) \\ &= \sum_0^{\infty} |a_n|^2 (1 - r^n)^2 \longrightarrow 0 \text{ as } r \rightarrow 1. \end{aligned}$$

Therefore, $\tilde{f}(r\xi)$ converges to $f(\xi)$ in measure. It remains to show that the convergence is almost everywhere. Let us define a kernel $S_z(\xi)$ as

$$S_z(\xi) := \frac{1}{1 - \xi \bar{z}} \text{ for } \xi \in \mathbb{T}. \quad (3.30)$$

Then we have

$$\tilde{f}(z) = \langle f, S_z \rangle_{L^2} \text{ for } z \in \mathbb{D}, f \in H^2(\mathbb{T}). \quad (3.31)$$

Consider the sum

$$\begin{aligned} S_z(\xi) + \overline{S_z(\xi)} - 1 &= \frac{1}{1 - \xi \bar{z}} + \frac{1}{1 - z \bar{\xi}} - 1, \quad (z = re^{i\theta}), (\xi = e^{it}) \\ &= \frac{1}{1 - e^{it} r e^{i\theta}} + \frac{1}{1 - r e^{i\theta} e^{it}} - 1 \\ &= \frac{1 - r^2}{1 - 2r \cos(t - \theta) + r^2} \\ &= P_r(t - \theta) \end{aligned}$$

and by

$$P_z(\xi) = \frac{1 - |z|^2}{|1 - \xi \bar{z}|^2} = \frac{1 - r^2}{1 - 2r \cos(t - \theta) + r^2} = P_r(t - \theta), \quad (z = re^{i\theta}), (\xi = e^{it})$$

we get

$$P_z(\xi) = S_z(\xi) + \overline{S_z(\xi)} - 1. \quad (3.32)$$

Then consider the inner product $\langle f, P_z \rangle$. Since $\overline{S_z(\xi)} - 1$ is orthogonal to f in $L^2(\mathbb{T})$, we get

$$\begin{aligned} \langle f, P_z \rangle_{L^2} &= \langle f, S_z(\xi) + \overline{S_z(\xi)} - 1 \rangle_{L^2} = \langle f, S_z(\xi) \rangle + \langle f, \overline{S_z(\xi)} - 1 \rangle_{L^2} \\ &= \langle f, S_z(\xi) \rangle_{L^2} = \tilde{f}(z). \end{aligned}$$

So, instead of (3.31), we can use $\tilde{f}(z) = \langle f, P_z \rangle_{L^2}$. Then, by the definition of the inner product in L^2 , we have

$$\tilde{f}(z) = \langle f, P_z \rangle_{L^2} = \int_{-\pi}^{\pi} f(\xi) \overline{P_z(\xi)} d\sigma(\xi). \quad (3.33)$$

Changing variables as $z = re^{i\theta}$ and $\xi = e^{i\theta}$, it follows

$$\tilde{f}(re^{i\theta}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(e^{i\theta}) P_r(t - \theta) dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(e^{i(t+\theta)}) P_r(t) dt.$$

We fix θ in $(-\pi, \pi)$ so that $F'(\theta)$ exists, where $F(\theta) := \int_{-\pi}^{\pi} f(e^{i\theta}) dt$. By using the property (iii) of the Poisson kernel,

$$\tilde{f}(re^{i\theta}) - F'(\theta) = \frac{1}{2\pi} \int_{-\pi}^{\pi} (\tilde{f}(re^{i\theta}) - F'(\theta)) P_r(t) dt.$$

We make change of variables and integrate by parts the right hand side of this equation, say $u = P_r(t)$ and $dv = (\tilde{f}(re^{i\theta}) - F'(\theta)) dt$ and get

$$\frac{1}{2\pi} \left[(F(\theta + t) - F'(\theta)t) P_r(t) \right]_{-\pi}^{\pi} - \frac{1}{2\pi} \int_{-\pi}^{\pi} (F(\theta + t) - F'(\theta)t) P'_r(t) dt.$$

Then the first term converges to 0 as $r \rightarrow 1$. So, we have

$$\lim_{r \rightarrow 1} \tilde{f}(re^{i\theta}) - F'(\theta) = \lim_{r \rightarrow 1} \frac{1}{2\pi} \int_{-\pi}^{\pi} (F(\theta + t) - F'(\theta)t) P'_r(t) dt. \quad (3.34)$$

For any $\varepsilon > 0$ there exists $\delta > 0$ such that

$$|F(\theta + t) - F(\theta - t) - 2F'(\theta)t| \leq \varepsilon t \quad \text{for } 0 \leq t \leq \delta. \quad (3.35)$$

On the other hand,

$$\begin{aligned} \tilde{f}(re^{i\theta}) - F'(\theta) &= \frac{1}{2\pi} \int_{-\pi}^{\pi} (F(\theta + t) - F'(\theta)t) P'_r(t) dt \\ &\leq \frac{1}{2\pi} \int_0^{\delta} (F(\theta + t) - F(\theta - t) - 2F'(\theta)t) P'_r(t) dt \\ &\quad + \frac{1}{2\pi} \int_{\delta \leq |t| \leq \pi} (F(\theta + t) - F'(\theta)t) P'_r(t) dt, \end{aligned}$$

and then by using (3.35),

$$\left| \frac{1}{2\pi} \int_{-\pi}^{\pi} (F(\theta + t) - F'(\theta) t) P_r'(t) dt \right| \leq \varepsilon \int_0^{\pi} t |P_r'(t)| dt + \gamma \sup_{\delta \leq |t| \leq \pi} |P_r'(t)|,$$

where

$$\gamma = \sup_{|t| \leq \pi} |F(\theta + t) - F'(\theta) t|.$$

Now, since we have the derivative of $P_r(t)$ as

$$P_r'(t) = \frac{2r \sin t (1 - r^2)}{(1 - 2r \cos t + r^2)^2},$$

taking modulus of this derivative we get

$$|P_r'(t)| = \left| \frac{2r \sin t (1 - r^2)}{(1 - 2r \cos t + r^2)^2} \right| \leq \frac{2r(1 - r^2)}{(1 - 2r \cos t + r^2)^2},$$

and this implies $\sup_{\delta \leq |t| \leq \pi} |P_r'(t)| \rightarrow 0$, as $r \rightarrow 1$. Also, $t |P_r'(t)|$ will be clearly integrable on $[0, \pi]$. Consequently, taking the lim sup of (3.34), it converges to 0 as $r \rightarrow 1$. That is

$$\limsup_{r \rightarrow 1} |\tilde{f}(re^{i\theta}) - F'(\theta)| = 0.$$

Hence $\tilde{f}(re^{i\theta})$ converges as $r \rightarrow 1$ at all θ for which $F'(\theta)$ exists. \square

Since $\tilde{f}(z)$ is analytic on \mathbb{D} and $f(\xi)$ is its boundary function on \mathbb{T} , then $\tilde{f}(z)$ is the analytic extension of f to \mathbb{D} .

Definition 3.3.8. The kernel

$$S(\xi, z) := \frac{1}{1 - \xi \bar{z}} \quad \text{for } \xi \in \mathbb{T} \text{ and } z \in \mathbb{D}, \quad (3.36)$$

or its analytic extension

$$\tilde{S}(w, z) := \frac{1}{1 - w \bar{z}} \quad \text{for } w, z \in \mathbb{D}. \quad (3.37)$$

is called the *Szegö kernel*.

Since $K_{\alpha}(w, z) = \frac{1}{(1 - w \bar{z})^{\alpha}}$, note that it is equal to the Szegö kernel for $\alpha = 1$, i.e. $\tilde{S}(w, z) = K_1(w, z)$.

We will end this chapter by a remark.

Remark 3.3.9. The orthogonal projection $P_{H^2(\mathbb{T})}$ from $L^2(\mathbb{T})$ to the Hardy space $H^2(\mathbb{T})$ can be expressed in terms of the Szegő kernel as follows:

$$(P_{H^2}g)(z) = \langle g, S_z \rangle_{L^2} \quad \text{for } z \in \mathbb{D} \text{ and } g \in L^2(\mathbb{T}). \quad (3.38)$$

Moreover, by Theorem 3.3.7, we have

$$(P_{H^2}g)(\xi) = \lim_{r \rightarrow 1} \langle g, S_{r\xi} \rangle_{L^2} \quad \text{for almost all } \xi \in \mathbb{T}. \quad (3.39)$$

Chapter 4

Interpolation Theorems of Nevanlinna-Pick Type

4.1 General definition of Hardy spaces

In the previous chapter we have defined H^p spaces for the case $p = 2$. Before giving a more general definition and some properties of H^p spaces, we first define the *integral means*.

Definition 4.1.1. For a function analytic in the unit disc \mathbb{D} , the *integral means* are defined by

$$M_p(r, \psi) = \left(\frac{1}{2\pi} \int_0^{2\pi} |\psi(re^{i\theta})|^p d\theta \right)^{1/p}, \quad (0 < p < \infty)$$

and

$$M_\infty(r, \psi) = \max_{0 \leq \theta < 2\pi} |\psi(re^{i\theta})|.$$

Definition 4.1.2. The class H^p is the set of all functions analytic in \mathbb{D} for which the integral means $M_p(r, \psi)$ are bounded for $0 \leq r < 1$.

By the above definitions and the maximum modulus principle, if $M_p(r, \psi)$ stays bounded as $r \rightarrow 1$, then ψ is said to belong to the H^p space.

Consider the case $p = 2$. If $\psi(z) = \sum a_n z^n$ is analytic in \mathbb{D} , then by Parseval's identity, we have

$$M_2^2(r, \psi) = \sum_{n=0}^{\infty} |a_n|^2 r^{2n}.$$

This shows that $M_2(r, \psi)$ increases with r , and that $\psi \in H^2$ if and only if $\sum |a_n|^2 < \infty$. Similarly, it follows from the maximum modulus principle that M_∞ increases with r . This situation is more complicated for the other values of p , but $M_p(r, \psi)$ is always a non-decreasing function.

The H^p class is a linear space under addition and scalar multiplication. The norm $\|\psi\|_{H^p}$ of a function $\psi \in H^p$ is defined as the limit of $M_p(r, \psi)$ as $r \rightarrow 1$ and it is a *true norm* if $p \geq 1$.

Another property of H^p spaces is that, if $0 < p < q \leq \infty$, then $H^p \supset H^q$.

Moreover, by the above relations, $H^\infty(\mathbb{D})$ is the class of all bounded analytic functions $\psi(z)$ on \mathbb{D} with norm

$$\|\psi\|_\infty = \sup_{z \in \mathbb{D}} |\psi(z)|. \quad (4.1)$$

According to Theorem 3.3.7, each $\psi \in H^\infty(\mathbb{D})$ admits the boundary value for σ -almost all $\zeta \in \mathbb{T}$, (\mathbb{T} is the unit circle),

$$\psi(\zeta) := \lim_{r \rightarrow 1} \psi(r\zeta). \quad (4.2)$$

From now on, norm $\|\cdot\|$ and inner product $\langle \cdot, \cdot \rangle$ will be used for the space $H^2(\mathbb{T})$ and the reproducing kernel, i.e. the Szegő kernel $S(w, z)$ for $H^2(\mathbb{T})$ will be denoted by $K(w, z)$, to avoid any confusion:

$$K(w, z) = K_z(w) := \frac{1}{1 - w\bar{z}} \text{ for } z, w \in \mathbb{D}.$$

In order to see the connection between $H^2(\mathbb{T})$ and $H^\infty(\mathbb{D})$, let us consider any function $\psi \in H^\infty(\mathbb{D})$. For each ψ , there is a *multiplication operator* M_ψ on $H^2(\mathbb{T})$ defined by

$$(M_\psi h)(\zeta) := \psi(\zeta)h(\zeta) \text{ for } \zeta \in \mathbb{T} \text{ and } h \in H^2(\mathbb{T}),$$

or, with respect to analytic extension,

$$(M_\psi h)(z) = \psi(z)h(z) \quad \text{for } z \in \mathbb{D}. \quad (4.3)$$

Lemma 4.1.3. *Let M_ψ be a multiplication operator on $H^2(\mathbb{T})$ defined as in (4.3). Then $\|M_\psi\| = \|\psi\|$ for all $\psi \in H^\infty(\mathbb{D})$.*

Proof. Since

$$\|\psi\|_\infty = \sup_{z \in \mathbb{D}} |\psi(z)|,$$

it is clear that $\|M_\psi\| \leq \|\psi\|_\infty$.

For the converse inequality, since

$$K(z, z) = K_z(z) = \frac{1}{1 - z\bar{z}} = \frac{1}{1 - |z|^2}, \quad z \in \mathbb{D},$$

then

$$\begin{aligned} \frac{|\psi(z)|}{1 - |z|^2} &= |\psi(z)| |K(z, z)| = |\psi(z)| |\langle K_z, K_z \rangle| \\ &= |\psi(z)| \langle K_z, K_z \rangle = |\langle \psi(z) K_z, K_z \rangle| \\ &= |\langle M_\psi K_z, K_z \rangle| \quad \text{since } \psi K_z = M_\psi K_z \\ &\leq \|M_\psi\| \|K_z\|^2 \\ &= \|M_\psi\| \left(\frac{1}{1 - |z|^2} \right), \quad z \in \mathbb{D}. \end{aligned}$$

So, we get

$$\frac{|\psi(z)|}{1 - |z|^2} \leq \|M_\psi\| \frac{1}{1 - |z|^2},$$

which yields

$$\sup_{z \in \mathbb{D}} |\psi(z)| = \|\psi\|_\infty \leq \|M_\psi\|.$$

Hence,

$$\|\psi\|_\infty = \|M_\psi\| \quad \text{for } \psi \in H^\infty(\mathbb{D}). \quad \square$$

If $\psi, \phi \in H^\infty(\mathbb{D})$, we have $M_{\phi\psi} = M_\phi M_\psi$ for the map $\psi \mapsto M_\psi$ from $H^\infty(\mathbb{D})$ to $B(H^2(\mathbb{T}))$ and ψ is linear and multiplicative.

Definition 4.1.4. The multiplication operator corresponding to the function $\phi(z) := z$, which is defined by

$$S(\phi(z)) = z\phi(z),$$

is called *the shift operator* and is denoted by S .

It is easy to see that, for multiplication operator M_ψ on $H^2(\mathbb{T})$ and shift operator S on $B(H^2(\mathbb{T}))$, we have

$$M_\psi S = SM_\psi. \quad (4.4)$$

Theorem 4.1.5. *If a continuous linear operator T on $H^2(\mathbb{T})$ commutes with the shift operator S , i.e. $TS = ST$, then there exists uniquely $\psi \in H^\infty(\mathbb{D})$ for which $T = M_\psi$.*

Proof. Let l be the constant function with value 1 and let $\psi = Tl$. Since

$$K_z(w) = \frac{1}{1 - w\bar{z}} = \sum_{n=0}^{\infty} (w\bar{z})^n, \quad z, w \in \mathbb{D},$$

then

$$K_z(z) = \sum_{n=0}^{\infty} \bar{z}^n z^n = \sum_{n=0}^{\infty} \bar{z}^n S^n l.$$

Since by assumption T is continuous on $H^2(\mathbb{T})$ and commutes with shift operator S , the above relation implies

$$\begin{aligned} TK_z &= \sum_{n=0}^{\infty} \bar{z}^n T S^n l = \sum_{n=0}^{\infty} \bar{z}^n S^n T l = \sum_{n=0}^{\infty} \bar{z}^n S^n \psi \quad \text{since } \psi = Tl, \\ &= \psi \sum_{n=0}^{\infty} \bar{z}^n z^n = \psi K_z = M_\psi K_z, \quad z \in \mathbb{D}. \end{aligned}$$

So we get $TK_z = M_\psi K_z$ for $z \in \mathbb{D}$. Thus $\psi \in H^\infty(\mathbb{D})$. Moreover, since T and M_ψ are continuous and $\{K_z : z \in \mathbb{D}\}$ is total in $H^2(\mathbb{T})$, then we have $T = M_\psi$.

For the uniqueness part, suppose that there exists ψ_1 and ψ_2 such that $T = M_{\psi_1}$ and $T = M_{\psi_2}$. It follows by (4.3) that

$$TK_z = M_{\psi_1} K_z = \psi_1 K_z \quad \text{and} \quad TK_z = M_{\psi_2} K_z = \psi_2 K_z,$$

which yields

$$\psi_1 K_z = M_{\psi_1} K_z = M_{\psi_2} K_z = \psi_2 K_z.$$

Thus $\psi_1 = \psi_2$. □

Corollary 4.1.6. *Let \mathcal{M} be a closed non-zero subspace of $H^2(\mathbb{T})$ and S be the shift operator. If \mathcal{M} is invariant for the adjoint S^* , i.e. $S^*h \in \mathcal{M}$ for all $h \in \mathcal{M}$, then the closed linear span of $\{S^n(\mathcal{M}) : n = 0, 1, \dots\}$ coincides with $H^2(\mathbb{T})$.*

Proof. Let \mathcal{M} be a closed non-zero subspace of $H^2(\mathbb{T})$. Denote by \mathcal{C} the closed linear span of $\{S^n(\mathcal{M}) : n = 0, 1, \dots\}$. Since $S^*S = I$ and by assumption $S^*(\mathcal{M}) \subseteq \mathcal{M}$, then \mathcal{C} is invariant for S^* , i.e. $S^*(S^n(g)) \subseteq \mathcal{C}$ for all $g \in \mathcal{M}$. In fact $S^*(S^n(g)) = S^{n-1}(g) \subseteq \mathcal{C}$ as $SS^* = I$. Clearly \mathcal{C} is also invariant for S .

Let P be the orthogonal projection to the closed linear span \mathcal{C} .

Claim: $PS = SP$.

By using the definition of orthogonal projection, for $g = g_1 + g_2$ where $g_1 \in \mathcal{M}$, and $g_2 \in \mathcal{M}^\perp$, we have $P(g) = g_1$. Taking $h \in \mathcal{M}$, $\langle h, Sg_2 \rangle = \langle S^*h, g_2 \rangle = 0$ since, by assumption, $S^*h \in \mathcal{M}$. In other words, $Sg_2 \in \mathcal{M}^\perp$. Then

$$(PS)(g) = (PS)(g_1 + g_2) = P(S(g_1) + S(g_2)) = S(g_1) = (SP)(g),$$

that is, $PS = SP$ as claimed.

By previous theorem, $P = M_\psi$ for some $\psi \in H^\infty(\mathbb{D})$. Since $P^2 = P$, we have $(M_\psi)^2 = M_\psi$ which implies that $\psi(z)^2 = \psi(z)$. But since $\psi \in H^\infty(\mathbb{D})$, ψ is analytic. This means, $\psi^2 = \psi$ only if $\psi(z) \equiv 0$ or $\psi(z) \equiv 1$. By assumption since $\mathcal{M} \neq 0$, then $\psi(z) \equiv 1$. That is,

$$M_\psi h = h \text{ for all } h \in H^2.$$

Hence the closed linear span of $\{S^n(\mathcal{M}) : n = 0, 1, \dots\}$ coincides with $H^2(\mathbb{T})$. □

The following theorem states that two functions in $H^2(\mathbb{T})$ can coincide on a set of positive σ -measure only when they represent one and the same element of $H^2(\mathbb{T})$.

Theorem 4.1.7. *A function $h \in H^2(\mathbb{T})$ can vanish on a measurable subset Λ of \mathbb{T} with positive σ -measure only if $h = 0$.*

Proof. Let Λ be a measurable subset \mathbb{T} with positive σ -measure. Consider the subspace $\mathcal{M} := \{f \in H^2(\mathbb{T}) : f(\zeta) = 0 \text{ on } \Lambda \text{ } \sigma - a.e \text{ (almost everywhere)}\}$.

Claim: $\mathcal{M} = \{0\}$.

\mathcal{M} is a closed subspace since strong convergence in $H^2(\mathbb{T})$ implies convergence in measure on \mathbb{T} . Let S be the shift operator. Since, by definition, $S(h) \in \mathcal{M}$ for all $h \in \mathcal{M}$, then \mathcal{M} is invariant for S . Let P be the orthogonal projection onto \mathcal{M} and l be the constant function with value 1. Consider $g := Pl$. Then $l - g = l - Pl = (I - P)l$ is orthogonal to \mathcal{M} . As \mathcal{M} is invariant for S , it follows that $l - g$ is also orthogonal to $S^n g$ for $n = 0, 1, \dots$

Since

$$\langle S^n g, l \rangle = (S^n g)(0) = 0 \quad \text{for } n = 1, 2, \dots,$$

we have

$$\begin{aligned} 0 &= \langle S^n g, l - g \rangle = \int_{\mathbb{T}} (S^n g)(\zeta) \overline{(l - g)(\zeta)} d\sigma(\zeta) \\ &= \int_{\mathbb{T}} S^n(\zeta) g(\zeta) \overline{l(\zeta)} d\sigma(\zeta) - \int_{\mathbb{T}} (S^n g)(\zeta) \overline{g(\zeta)} d\sigma(\zeta) \\ &= - \int_{\mathbb{T}} \zeta^n |g(\zeta)|^2 d\sigma(\zeta) \quad \text{as } \langle S^n g, l \rangle = 0. \end{aligned}$$

This means that all Fourier coefficients of $|g|^2$, except the constant term, vanish, thus $|g(\zeta)|^2$ is constant on \mathbb{T} $\sigma - a.e$. Since $g = Pl \in \mathcal{M}$, $g(\zeta)$ vanishes on Λ of positive σ -measure which implies that $g = 0$, i.e. $g = Pl = 0$. This means l is orthogonal to \mathcal{M} . Then for each $f \in \mathcal{M}$, as $Sf \in \mathcal{M}$, we have $(S^* f)(\zeta) = \bar{\zeta} f(\zeta)$, hence $S^* f$ vanishes on Λ $\sigma - a.e$., which means that \mathcal{M} is invariant for S^* . Hence, since \mathcal{M} is invariant for both S and S^* and l is not in \mathcal{M} , by previous corollary, $\mathcal{M} = \{0\}$. \square

Corollary 4.1.8. *Let P_+ be the orthogonal projection from the Hilbert space $L^2(\mathbb{T})$ to $H^2(\mathbb{T})$. If Λ is a measurable subset of \mathbb{T} with positive σ -measure, then $\{P_+(\chi_\Lambda f) : f \in L^2(\mathbb{T})\}$ is a dense subspace of $H^2(\mathbb{T})$, where $\chi_\Lambda(\zeta) = 1$ is the characteristic function of Λ .*

Proof. We show that, if $H^2(\mathbb{T}) \ni h \perp \chi_\Lambda f$ for all $f \in L^2(\mathbb{T})$, then $h = 0$. To see this, consider the previous theorem,

$$\begin{aligned} 0 &= \langle \chi_\Lambda h, h \rangle = \int_\Lambda (\chi_\Lambda h)(\zeta) \overline{h(\zeta)} d\sigma(\zeta) \\ &= \int_\Lambda h(\zeta) \overline{h(\zeta)} d\sigma(\zeta) = \int_\Lambda |h(\zeta)|^2 d\sigma(\zeta), \end{aligned}$$

implies that h vanishes on Λ σ -a.e. Hence $\{P_+(\chi_\Lambda f) : f \in L^2(\mathbb{T})\}$ is a dense subspace of $H^2(\mathbb{T})$. \square

A consequence of the above proof is that, $\{P_+(\chi_\Lambda g) : g \in H^2(\mathbb{T})\}$ is dense in $H^2(\mathbb{T})$.

4.2 Interpolation Inside Unit Disc

In this section we consider interpolation on finite and infinite subsets of the unit disc. In the following theorem we consider interpolation on a finite subset.

Theorem 4.2.1. (*Nevanlinna-Pick*) *Let $X = \{z_1, \dots, z_n\}$ be a finite subset of \mathbb{D} and ϕ be a function on X . Then, in order for there to exist an analytic function $\psi \in H^\infty(\mathbb{D})$ such that*

$$\|\psi\|_\infty \leq 1 \quad \text{and} \quad \psi(z_i) = \phi(z_i), \quad (i = 1, 2, \dots, n) \quad (4.5)$$

it is necessary and sufficient that the kernel $\left[\frac{1 - \phi(z_i) \overline{\phi(z_j)}}{1 - z_i \overline{z_j}} \right]$ on X is positive definite, that is,

$$\sum_{i,j=1}^n \bar{\xi}_i \xi_j \frac{1 - \phi(z_i) \overline{\phi(z_j)}}{1 - z_i \overline{z_j}} \geq 0 \quad \text{for all} \quad \{\xi_i\}_{i=1}^n \subset \mathbb{C}. \quad (4.6)$$

Proof. Necessity: Suppose that there exists $\psi \in H^\infty(\mathbb{D})$ satisfying (4.5).

Claim: $M_\psi^* K_z = \overline{\psi(z)} K_z$ where $K_z(w) = K(w, z) = \frac{1}{1 - w\bar{z}}$ for $w, z \in \mathbb{D}$.

By using the properties of multiplication operators and reproducing kernels,

we get

$$\begin{aligned}
(M_\psi^* K_z)(w) &= \langle M_\psi^* K_z, K_w \rangle = \langle K_z, M_\psi K_w \rangle = \langle K_z, \psi(z) K_w \rangle \\
&= \overline{\psi(z)} \langle K_z, K_w \rangle = \overline{\psi(z)} K(w, z) = \overline{\psi(z)} \overline{K(z, w)} \\
&= \overline{\psi(z)} K_z(w).
\end{aligned}$$

Hence $M_\psi^* K_z = \overline{\psi(z)} K_z$ for $z \in \mathbb{D}$, as claimed.

Since, by assumption, $\|\psi\|_\infty \leq 1$, and by Lemma 4.1.3 we have $\|M_\psi\| = \|\psi\|_\infty$, combining these we get $\|M_\psi\| = \|M_\psi^*\| \leq 1$. Then using Claim and assumption,

$$\begin{aligned}
0 &\leq \left\| \sum_{j=1}^n \xi_j K_{z_j} \right\|^2 - \left\| M_\psi^* \left(\sum_{j=1}^n \xi_j K_{z_j} \right) \right\|^2 \quad \text{as } \|M_\psi^*\| \leq 1 \\
&= \left\| \sum_{j=1}^n \xi_j K_{z_j} \right\|^2 - \left\| \sum_{j=1}^n \xi_j M_\psi^* K_{z_j} \right\|^2 \\
&= \left\| \sum_{j=1}^n \xi_j K_{z_j} \right\|^2 - \left\| \sum_{j=1}^n \xi_j \overline{\psi(z_j)} K_{z_j} \right\|^2 \quad \text{as } M_\psi^* K_z = \overline{\psi(z)} K_z \\
&= \left\langle \sum_{j=1}^n \xi_j K_{z_j}, \sum_{j=1}^n \xi_j K_{z_j} \right\rangle - \left\langle \sum_{j=1}^n \xi_j \overline{\psi(z_j)} K_{z_j}, \sum_{j=1}^n \xi_j \overline{\psi(z_j)} K_{z_j} \right\rangle \\
&= \sum_{i,j=1}^n \bar{\xi}_i \xi_j (1 - \psi(z_i) \overline{\psi(z_j)}) \langle K_{z_j}, K_{z_i} \rangle \\
&= \sum_{i,j=1}^n \bar{\xi}_i \xi_j \frac{1 - \phi(z_i) \overline{\phi(z_j)}}{1 - z_i \bar{z}_j} \quad \text{for all } \{\xi_i\}_{i=1}^n \subset \mathbb{C},
\end{aligned}$$

which is the desired result.

Sufficiency: Suppose that the kernel $\left[\frac{1 - \phi(z_i) \overline{\phi(z_j)}}{1 - z_i \bar{z}_j} \right]$ on X is positive definite, that is,

$$\sum_{i,j=1}^n \bar{\xi}_i \xi_j \frac{1 - \phi(z_i) \overline{\phi(z_j)}}{1 - z_i \bar{z}_j} \geq 0 \quad \text{for all } [\xi_i].$$

Denote by \mathcal{M} the closed linear span of K_{z_j} , ($j = 1, 2, \dots, n$), and S shift operator. By the Claim in the necessity part, we have $S^* K_z = \bar{z} K_z$ for any $z \in \mathbb{D}$. This implies that \mathcal{M} is invariant for S^* . Then consider the following function

$$\tilde{\psi}(z) := \sum_{i=1}^n \frac{\prod_{j \neq i} \frac{z - z_j}{1 - z \bar{z}_j}}{\prod_{j \neq i} \frac{z_i - z_j}{1 - z_i \bar{z}_j}} \phi(z_i).$$

Note that this function $\tilde{\psi} \in H^\infty(\mathbb{D})$ satisfies the following

$$\tilde{\psi}(z_i) = \phi(z_i) \quad (i = 1, 2, \dots, n). \quad (4.7)$$

Let P be the orthogonal projection onto \mathcal{M} . Now define the linear operator T on \mathcal{M} by

$$T = PM_{\tilde{\psi}}|_{\mathcal{M}}. \quad (4.8)$$

Since (4.7) is satisfied, then (4.6) becomes

$$\sum_{i,j=1}^n \bar{\xi}_i \xi_j \frac{1 - \tilde{\psi}(z_i) \overline{\tilde{\psi}(z_j)}}{1 - z_i \bar{z}_j} \geq 0 \quad \text{for } \{\xi_i\}_{i=1}^n \subset \mathbb{C}.$$

It follows that $\|T^*h\| \leq \|h\|$ for $h \in \mathcal{M}$ and hence $\|T\| \leq 1$. To be able to finish this proof we need the following lemma.

Lemma 4.2.2. *Suppose that \mathcal{M}_0 is a non-zero closed subspace of $H^2(\mathbb{T})$, invariant for the adjoint S^* , and that T_0 is a continuous linear operator on \mathcal{M}_0 , commuting with the restricted shift $S_0 := P_0 S|_{\mathcal{M}_0}$ where P_0 is the orthogonal projection onto \mathcal{M}_0 , that is,*

$$T_0 S_0 = S_0 T_0. \quad (4.9)$$

Then the closed linear span \mathcal{M}_1 of \mathcal{M}_0 and $S\mathcal{M}_0$ is invariant for S^ and there is a continuous linear operator T_1 on \mathcal{M}_1 such that*

$$P_0 T_1 = T_0 P_0, \quad \|T_1\| = \|T_0\|, \quad \text{and} \quad T_1 S_1 = S_1 T_1, \quad (4.10)$$

where $S_1 = P_1 S|_{\mathcal{M}_1}$, P_1 being the orthogonal projection onto \mathcal{M}_1 .

Proof. We prove this lemma in five steps.

Step 1: \mathcal{M}_1 is invariant for S^* .

By definition, \mathcal{M}_0 is invariant for S^* means $S^*(\mathcal{M}_0) \subseteq \mathcal{M}_0$. Since $S^*S = I$, we have $S^*(S(\mathcal{M}_0)) = \mathcal{M}_0$. Then, since \mathcal{M}_1 is defined as closed linear span of \mathcal{M}_0 and $S(\mathcal{M}_0)$, we have \mathcal{M}_1 is invariant for S^* .

Step 2: $S_1 = SP_0$ on \mathcal{M}_1 .

Let $\mathcal{M}_1 \ni g$ be orthogonal on \mathcal{M}_0 . Then, since by Step 1, $S^*h \in \mathcal{M}_1$ for $h \in \mathcal{M}_0$, it follows that

$$\langle Sg, h \rangle = \langle g, S^*h \rangle = 0 \text{ for } h \in \mathcal{M}_0,$$

that is, Sg is orthogonal to \mathcal{M}_0 and

$$\langle Sg, Sh \rangle = \langle g, h \rangle = 0 \quad \text{for } g \in \mathcal{M}_1, h \in \mathcal{M}_0.$$

This implies that Sg is also orthogonal to $S(\mathcal{M}_0)$. Now, since $S_1 = P_1S$ on \mathcal{M}_1 , where P_1 is the orthogonal projection onto \mathcal{M}_1 , then we get

$$S_1g = P_1Sg = P_1(S(g)) = 0 \quad \text{because } S(g) \perp \mathcal{M}_0.$$

It follows that $S_1 = S_1P_0$ on \mathcal{M}_1 . Since $S(\mathcal{M}_0) \subseteq \mathcal{M}_1$, we obtain

$$S_1 = S_1P_0 = SP_0.$$

Before the third step, without loss of generality we assume that we have $\|T_0\| = 1$.

Step 3 : $\|(S_1 - S_0P_0)T_0P_0h\| \leq \|(I_1 - (T_0P_0)^*(T_0P_0))^{1/2} \cdot S_1h\|$ for $h \in \mathcal{M}_1$, and I_1 is the identity operator of \mathcal{M}_1 .

By assumptions and previous steps, we have the following,

$$\begin{aligned} \|(S_1 - S_0P_0)T_0P_0h\|^2 &= \|(SP_0 - S_0P_0)T_0P_0h\|^2 = \|(S - S_0)P_0T_0P_0h\|^2 \\ &= \|(S - SP_0)P_0T_0P_0h\|^2 = \|(I - P_0)SP_0T_0P_0h\|^2 \\ &= \|SP_0T_0P_0h\|^2 - \|S_0T_0P_0h\|^2 \\ &\leq \|P_0h\|^2 - \|S_0T_0P_0h\|^2 = \|SP_0h\|^2 - \|T_0S_0P_0h\|^2 \\ &= \|S_1h\|^2 - \|T_0P_0S_1h\|^2 \\ &= \langle S_1h, S_1h \rangle - \langle T_0P_0S_1h, T_0P_0S_1h \rangle \\ &= \langle S_1h, S_1h \rangle - \langle (T_0P_0)^*(T_0P_0)S_1h, S_1h \rangle \\ &= \langle S_1h - (T_0P_0)^*(T_0P_0)S_1h, S_1h \rangle \\ &= \langle (I_1 - (T_0P_0)^*(T_0P_0))S_1h, S_1h \rangle. \end{aligned}$$

Considering the identity operator I_1 of \mathcal{M}_1 , since $\|T_0P_0\| \leq 0$ implies $I_1 - (T_0P_0)^*(T_0P_0) \geq 1$, we can consider the positive definite square root

$$(I_1 - (T_0P_0)^*(T_0P_0))^{1/2}.$$

So from the above inequality it follows as

$$\|(S_1 - S_0P_0)T_0P_0h\| \leq \|(I_1 - (T_0P_0)^*(T_0P_0))^{1/2} \cdot S_1h\| \quad \text{for } h \in \mathcal{M}_1.$$

This means that we have a well-defined linear operator from the range of $(I_1 - (T_0P_0)^*(T_0P_0))^{1/2} \cdot S_1$ to the range of $(S_1 - S_0P_0)T_0P_0$ and of norm ≤ 1 . Now consider its continuous extension, composed with the orthogonal projection to the closure of the range of $(I_1 - (T_0P_0)^*(T_0P_0))^{1/2} \cdot S_1$, then we can conclude that there is a continuous linear operator L from \mathcal{M}_1 to the closure of the range of $S_1 - S_0P_0$ such that $\|L\| \leq 1$ and

$$(S_1 - S_0P_0)T_0P_0 = L(I_1 - (T_0P_0)^*(T_0P_0))^{1/2} \cdot S_1. \quad (4.11)$$

Define T_1 by

$$T_1 = T_0P_0 + L(I_1 - (T_0P_0)^*(T_0P_0))^{1/2}. \quad (4.12)$$

Step 4: $P_0T_1 = T_0P_0$ and $T_1S_1 = S_1T_1$.

Since the range of L is contained in the closure of $S_1 - S_0P_0$, applying P_0 to both sides of (4.12), we get

$$\begin{aligned} P_0T_1 &= P_0T_0P_0 + P_0L \left(I_1 - (T_0P_0)^*(T_0P_0) \right)^{1/2} \\ &= P_0T_0P_0 + P_0(S_1 - S_0P_0)T_0P_0 \\ &= P_0T_0P_0 \quad \text{since } P_0(S_1 - S_0P_0) = 0. \end{aligned}$$

Similarly,

$$\begin{aligned} T_1S_1 &= T_0P_0S_1 + L(I_1 - (T_0P_0)^*(T_0P_0))^{1/2} S_1 \\ &= T_0S_0P_0 + (S_1 - S_0P_0)T_0P_0 \\ &= S_1T_0P_0 \\ &= S_1T_1. \end{aligned}$$

Hence

$$P_0T_1 = T_0P_0 \quad \text{and} \quad T_1S_1 = S_1T_1.$$

Step 5: $\|T_1\| = \|T_0\|$.

Since $P_0L = 0$, we have

$$\begin{aligned} \|T_1h\|^2 &= \|T_0P_0h\|^2 + \|L(I_1 - (T_0P_0)^*(T_0P_0))^{1/2}h\|^2 \\ &\leq \|T_0P_0h\|^2 + \|h\|^2 - \|T_0P_0h\|^2 \quad \text{by } \|L\| \leq 1 \\ &= \|h\|^2, \end{aligned}$$

which yields $\|T_1\| \leq 1 = \|T_0\|$.

Conversely,

$$\|T_0\| = \|T_0P_0\| = \|P_0T_1\| \leq \|T_1\|.$$

So we have $\|T_1\| = \|T_0\|$.

Therefore T_1 satisfies all the requirements in (4.10) of Lemma. \square

Let us return to the proof of *sufficiency* part of previous theorem. To avoid confusion, let us take $\mathcal{M}_0 := \mathcal{M}$ and $T_0 := T$ in Lemma. We have that \mathcal{M}_0 is invariant under S^* and $\|T_0\| \leq 1$. Since $T_0 = P_0M_{\tilde{\psi}}|_{\mathcal{M}_0}$ and $S_0 = P_0S|_{\mathcal{M}_0}$ and \mathcal{M}_0^\perp is invariant under S and $M_{\tilde{\psi}}$, we have $P_0M_{\tilde{\psi}}P_0 = P_0M_{\tilde{\psi}}$ and $P_0SP_0 = P_0S$ such that

$$T_0S_0 = P_0M_{\tilde{\psi}}P_0S|_{\mathcal{M}_0} = P_0M_{\tilde{\psi}}S|_{\mathcal{M}_0} = P_0SM_{\tilde{\psi}}|_{\mathcal{M}_0} = S_0T_0.$$

Now we can change our theorem into the lemma in such a way that, we have to find a sequence of linear operators T_n on \mathcal{M}_n , the closed linear span of \mathcal{M}_{n-1} and $S(\mathcal{M}_{n-1})$, ($n = 1, 2, \dots$), such that

$$P_{n-1}T_n = T_{n-1}P_{n-1}, \quad \|T_n\| = \|T_{n-1}\| \quad \text{and} \quad T_nS_n = S_nT_n, \quad (4.13)$$

where, for each n , P_n is the orthogonal projection onto \mathcal{M}_n and $S_n = P_nS|_{\mathcal{M}_n}$. By Corollary 4.1.6, the closed linear span of $\{\mathcal{M}_n : n = 0, 1, \dots\}$ which is equal to that of $\{S^n(\mathcal{M}_0) : n = 0, 1, \dots\}$ must coincide with $H^2(\mathbb{T})$. This means that, P_n strongly converges to I as $n \rightarrow \infty$. By the equation (4.13), we reach

$$\|T_nP_n\| \leq 1 \quad \text{and} \quad P_nT_mP_m = T_nP_n \quad \text{for } n \leq m, \quad (4.14)$$

and hence it follows that there is a continuous linear operator T_∞ on $\mathcal{H}^2(\mathbb{T})$ such that

$$\|T_\infty\| \leq 1 \quad \text{and} \quad P_n T_\infty = T_n P_n \quad \text{for} \quad n = 0, 1, \dots \quad (4.15)$$

Claim: $T_\infty S = S T_\infty$.

Since P_n strongly converges to I as $n \rightarrow \infty$, then $S_n P_n$ strongly converges to S . By (4.15), we have $T_\infty = \lim_{n \rightarrow \infty} P_n T_n P_n$. These imply

$$\begin{aligned} S T_\infty &= S \lim_{n \rightarrow \infty} P_n T_n P_n = \lim_{n \rightarrow \infty} S_n T_n P_n \\ &= \lim_{n \rightarrow \infty} T_n S_n P_n \quad \text{by (4.13)} \\ &= T_\infty \left(\lim_{n \rightarrow \infty} S_n P_n \right) = T_\infty S. \end{aligned}$$

Then by Theorem 4.1.5, there exists $\psi \in H^\infty(\mathbb{D})$ such that

$$\|\psi\|_\infty \leq 1 \quad \text{and} \quad M_\psi = T_\infty. \quad (4.16)$$

Then by (4.15) and (4.16) we reach to

$$P_0 M_\psi|_{\mathcal{M}_0} = P_0 T_\infty|_{\mathcal{M}_0} = T_0.$$

Finally, by (4.7) and (4.16),

$$\begin{aligned} \frac{\phi(z_i)}{1 - |z_i|^2} &= \phi(z_i) \langle K_{z_i}, K_{z_i} \rangle = \langle T_0 K_{z_i}, K_{z_i} \rangle \\ &= \langle T_\infty K_{z_i}, K_{z_i} \rangle = \langle M_\psi K_{z_i}, K_{z_i} \rangle \\ &= \frac{\psi(z_i)}{1 - |z_i|^2}. \end{aligned}$$

Hence ψ satisfies all the requirements in (4.5) of Theorem 4.2.1. \square

In the following corollary we consider interpolation on an infinite subset of the unit disc.

Corollary 4.2.3. *Let X be an infinite subset of \mathbb{D} , and ϕ a function on X . In order for there to exist a function $\psi \in H^\infty(\mathbb{D})$ such that*

$$\|\psi\|_\infty \leq 1 \quad \text{and} \quad \psi(z) = \phi(z) \quad \text{for} \quad z \in X, \quad (4.17)$$

it is necessary and sufficient that the kernel $\left[\frac{1 - \phi(z_i) \overline{\phi(z_j)}}{1 - z_i \bar{z}_j} \right]$ on X is positive definite.

Proof. This follows by an application of Zorn's Lemma and the previous theorem.

□

4.3 Interpolation on the Boundary

We will end this chapter by a theorem of interpolation on the boundary of unit disk \mathbb{D} . Before this we need the following lemma.

Lemma 4.3.1. *Let Λ be a measurable subset of \mathbb{T} and ϕ a bounded measurable function on Λ . Then in order for there to exist $\psi \in H^\infty(\mathbb{D})$ such that*

$$\|\psi\|_\infty \leq 1 \quad \text{and} \quad \psi(\zeta) = \phi(\zeta) \quad \text{for } \zeta \in \Lambda \quad \sigma - a.e., \quad (4.18)$$

it is necessary and sufficient that

$$\|P_+(\bar{\phi}\chi_\Lambda f)\| \leq \|P_+(\chi_\Lambda f)\| \quad \text{for } f \in L^2(\mathbb{T}). \quad (4.19)$$

Proof. Suppose that (4.18) is satisfied for $\psi \in H^\infty(\mathbb{D})$. Then

$$\begin{aligned} \|P_+(\bar{\phi}\chi_\Lambda f)\| &= \sup_{g \in H^2(\mathbb{D}), \|g\| \leq 1} |\langle \bar{\phi}\chi_\Lambda f, g \rangle| = \sup_{g \in H^2(\mathbb{D}), \|g\| \leq 1} |\langle \chi_\Lambda f, \psi g \rangle| \\ &\leq \|\psi\|_\infty \|P_+(\chi_\Lambda f)\| \\ &\leq \|P_+(\chi_\Lambda f)\| \quad \text{since } \|\psi\|_\infty \leq 1. \end{aligned}$$

Hence (4.19) is satisfied.

Conversely, suppose that (4.19) is satisfied. Since by Corollary 4.1.8 $\{P_+(\chi_\Lambda f) : f \in L^2(\mathbb{T})\}$ is a dense subspace of $H^2(\mathbb{T})$, then there exists an operator T such that

$$\|T\| \leq 1 \quad \text{and} \quad TP_+(\chi_\Lambda f) = P_+(\bar{\phi}\chi_\Lambda f) \quad \text{for } f \in L^2(\mathbb{T}).$$

Claim: $TS^* = S^*T$.

Denote by \tilde{S} the multiplication by the function $k(\zeta) := \zeta$ in $L^2(\mathbb{T})$. Then we have $S^*P_+ = P_+S^*$. It follows that

$$\begin{aligned} S^*TP_+(\chi_\Lambda f) &= S^*P_+(\bar{\phi}\chi_\Lambda f) = P_+\tilde{S}^*(\bar{\phi}\chi_\Lambda f) \\ &= P_+(\bar{\phi}\chi_\Lambda \tilde{S}^* f) = TP_+(\chi_\Lambda \tilde{S}^* f) \\ &= TP_+(\tilde{S}^*(\chi_\Lambda f)) = TS^*P_+(\chi_\Lambda f). \end{aligned}$$

Since $\{P_+(\chi_\Lambda f) : f \in L^2(\mathbb{T})\}$ is dense in $H^2(\mathbb{T})$, the above equation implies

$$S^*T = TS^*.$$

By Theorem 4.1.5 there exists $\psi \in H^\infty(\mathbb{D})$ such that

$$\|\psi\|_\infty = \|T^*\| \leq 1 \quad \text{and} \quad T^* = M_\psi.$$

Finally, it remains to show $\psi(\zeta) = \phi(\zeta)$ for $\zeta \in \Lambda$ σ -a.e. Take $f \in L^2(\mathbb{T})$. Then

$$\begin{aligned} \int_\Lambda \psi(\zeta) \overline{f(\zeta)} d\sigma(\zeta) &= \langle T^*l, \chi_\Lambda f \rangle = \langle T^*l, P_+(\chi_\Lambda f) \rangle \\ &= \langle l, TP_+(\chi_\Lambda f) \rangle = \langle l, P_+(\bar{\phi}\chi_\Lambda f) \rangle \\ &= \langle l, \bar{\phi}\chi_\Lambda f \rangle = \int_\Lambda \phi(\zeta) f(\zeta) d\sigma(\zeta) \end{aligned}$$

which yields that

$$\psi(\zeta) = \phi(\zeta) \quad \text{for } \zeta \in \Lambda, \sigma\text{-a.e.} \quad \square$$

Theorem 4.3.2. *Let Λ be a measurable subset of \mathbb{T} with positive σ -measure and ϕ a bounded measurable function on Λ . Then in order for there to exist $\psi \in H^\infty(\mathbb{D})$ such that*

$$\|\psi\|_\infty \leq 1 \quad \text{and} \quad \psi(\zeta) = \phi(\zeta) \quad \text{for } \zeta \in \Lambda \text{ } \sigma\text{-a.e.}, \quad (4.20)$$

it is necessary and sufficient that

$$\lim_{r \rightarrow 1} \int_\Lambda \int_\Lambda \left(\frac{1 - \phi(\xi)\overline{\phi(\zeta)}}{1 - r^2\xi\bar{\zeta}} \right) \overline{f(\xi)} f(\zeta) d\sigma(\xi) d\sigma(\zeta) \geq 0 \quad \text{for all } f \in L^2(\mathbb{T}). \quad (4.21)$$

Proof. Recall that in Chapter 2, the orthogonal projection P_{H^2} was expressed in terms of the Szegő kernel;

$$(P_{H^2}g)(z) = \langle g, K_z \rangle_{L^2}, \quad \text{for } z \in \mathbb{D} \text{ and } g \in L^2(\mathbb{T}),$$

and

$$(P_{H^2}g)(\zeta) = \lim_{r \rightarrow 1} \langle g, K_{r\zeta} \rangle \text{ for almost all } \zeta \in \mathbb{T}.$$

Then for $0 < r < 1$ and $\eta \in \mathbb{T}$, by above equations we get the followings;

$$(P_+g)(r\eta) = \int_{\mathbb{T}} g(\zeta) \frac{1}{1 - r\eta\bar{\zeta}} d\sigma(\zeta)$$

and

$$\begin{aligned} & \lim_{r \rightarrow 1} \int_{\mathbb{T}} |(P_+g)(r\eta) - (P_+g)(\eta)|^2 d\sigma(\eta) \\ &= \lim_{r \rightarrow 1} \int_{\mathbb{T}} \left| \int_{\mathbb{T}} g(\zeta) \frac{1}{1 - r\eta\bar{\zeta}} d\sigma(\zeta) - \int_{\mathbb{T}} g(\zeta) \frac{1}{1 - \eta\bar{\zeta}} d\sigma(\zeta) \right|^2 d\sigma(\eta) \\ &= \lim_{r \rightarrow 1} \int_{\mathbb{T}} \left| \int_{\mathbb{T}} g(\zeta) \left(\frac{1}{1 - r\eta\bar{\zeta}} - \frac{1}{1 - \eta\bar{\zeta}} \right) d\sigma(\zeta) \right|^2 d\sigma(\eta) \\ &= \lim_{r \rightarrow 1} \int_{\mathbb{T}} \left| \int_{\mathbb{T}} g(\zeta) \left(\frac{r\eta\bar{\zeta} - \eta\bar{\zeta}}{1 - \eta\bar{\zeta} - r\eta\bar{\zeta} + r\eta|\bar{\zeta}|^2} \right) d\sigma(\zeta) \right|^2 d\sigma(\eta) \\ &= 0. \end{aligned}$$

It follows that

$$\begin{aligned} \|P_+g\|^2 &= \langle P_+g, P_+g \rangle = \lim_{r \rightarrow 1} \int_{\mathbb{T}} |(P_+g)(r\eta)|^2 d\sigma(\eta) \\ &= \lim_{r \rightarrow 1} \int_{\mathbb{T}} \left(\int_{\mathbb{T}} \int_{\mathbb{T}} \frac{g(\zeta)}{1 - r\eta\bar{\zeta}} \frac{\overline{g(\varepsilon)}}{1 - r\eta\bar{\varepsilon}} d\sigma(\varepsilon) d\sigma(\zeta) \right) d\sigma(\eta) \\ &= \lim_{r \rightarrow 1} \int_{\mathbb{T}} \int_{\mathbb{T}} \left(\int_{\mathbb{T}} \frac{1}{1 - r\eta\bar{\zeta}} \frac{1}{1 - r\eta\bar{\varepsilon}} d\sigma(\eta) \right) \overline{g(\varepsilon)} g(\zeta) d\sigma(\varepsilon) d\sigma(\zeta) \\ &= \lim_{r \rightarrow 1} \int_{\mathbb{T}} \int_{\mathbb{T}} \langle K_{r\eta}, K_{r\varepsilon} \rangle \overline{g(\varepsilon)} g(\zeta) d\sigma(\varepsilon) d\sigma(\zeta) \\ &= \lim_{r \rightarrow 1} \int_{\mathbb{T}} \int_{\mathbb{T}} \frac{1}{1 - r^2\varepsilon\bar{\zeta}} \overline{g(\varepsilon)} g(\zeta) d\sigma(\varepsilon) d\sigma(\zeta). \end{aligned}$$

Therefore for all $g \in \mathcal{L}^2(\mathbb{T})$, we have

$$\|P_+g\|^2 = \lim_{r \rightarrow 1} \int_{\mathbb{T}} \int_{\mathbb{T}} \frac{1}{1 - r^2\varepsilon\bar{\zeta}} \overline{g(\varepsilon)} g(\zeta) d\sigma(\varepsilon) d\sigma(\zeta). \quad (4.22)$$

Since by previous lemma we have (4.19) by (4.18), these imply

$$\|P_+(\chi_\Lambda g)\|^2 - \|P_+(\bar{\phi} \chi_\Lambda g)\|^2 \geq 0.$$

Then it follows by (4.22) that

$$\lim_{r \rightarrow 1} \int_{\Lambda} \int_{\Lambda} \left(\frac{1 - \phi(\xi)\bar{\phi}(\zeta)}{1 - r^2\xi\bar{\zeta}} \right) \overline{g(\xi)} g(\zeta) d\sigma(\xi) d\sigma(\zeta) \geq 0 \quad \text{for all } g \in L^2(\mathbb{T}).$$

□

Appendix A

Hilbert Spaces

A.1 Definitions

Definition A.1.1. Let \mathcal{H} be a linear space over the complex field \mathbb{C} . An *inner product* on \mathcal{H} is a function $\langle \cdot, \cdot \rangle : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{C}$ which satisfies the following conditions:

- (i) $\langle f, g \rangle = \overline{\langle g, f \rangle}$ for $f, g \in \mathcal{H}$.
- (ii) $\langle \lambda f + \mu g, h \rangle = \lambda \langle f, h \rangle + \mu \langle g, h \rangle$ and
 $\langle f, \lambda g + \mu h \rangle = \bar{\lambda} \langle f, g \rangle + \bar{\mu} \langle f, h \rangle$ for $\lambda, \mu \in \mathbb{C}$ and f, g and $h \in \mathcal{H}$.
- (iii) $\langle f, f \rangle \geq 0$ for $f \in \mathcal{H}$ and $\langle f, f \rangle = 0$ if and only if $f = 0$.

Definition A.1.2. A *pre-Hilbert space* (*inner product space*) \mathcal{H} is a linear space over \mathbb{C} with an inner product defined on it.

If \mathcal{H} is an inner product space, $\| \cdot \|$ defines a norm on \mathcal{H} by

$$\|f\| = \langle f, f \rangle^{1/2}.$$

Sometimes we use the notations $\| \cdot \|_{\mathcal{H}}$ and $\langle \cdot, \cdot \rangle_{\mathcal{H}}$ for norm and inner product, respectively.

Some important properties:

$$\|f\| \geq 0 \text{ and equality occurs only when } f = 0. \quad (\text{A.1})$$

$$\|\xi f\| = |\xi| \|f\| \text{ for } f \in \mathcal{H}, \xi \in \mathbb{C}. \quad (\text{A.2})$$

$$|\langle f, g \rangle| \leq \|f\| \|g\| \text{ for } f, g \in \mathcal{H} \quad (\text{Schwarz Inequality}), \quad (\text{A.3})$$

and equality occurs when f and g are linearly dependent.

$$\|f + g\| \leq \|f\| + \|g\| \text{ for } f, g \in \mathcal{H} \quad (\text{Triangular Inequality}), \quad (\text{A.4})$$

and equality occurs when f and g are linearly dependent.

$$\langle f, g \rangle = \frac{1}{4} (\|f + g\|^2 - \|f - g\|^2 + i\|f + ig\|^2 - \|f - ig\|^2) \text{ for } f, g \in \mathcal{H} \quad (\text{A.5})$$

(polarization identity).

$$\|f + g\|^2 + \|f - g\|^2 = 2\|f\|^2 + 2\|g\|^2 \text{ for } f, g \in \mathcal{H} \quad (\text{parallelogram identity}). \quad (\text{A.6})$$

Define $d(f, g) := \|f - g\|$ the binary function. Observe that for f, g and $h \in \mathcal{H}$ and for all $\xi \in \mathbb{C}$, d satisfies all requirements to be a metric:

$$(i) \quad d(f, g) \geq 0 \text{ and equality occurs if and only if } f = g.$$

$$(ii) \quad d(f, g) = d(g, f).$$

$$(iii) \quad d(f, g) \leq d(f, h) + d(h, g).$$

$$(iv) \quad d(f - h, g - h) = d(f, g) \quad (\text{translation invariant}).$$

$$(v) \quad d(\xi f, \xi g) = |\xi| \cdot d(f, g).$$

This binary function introduces metric topology in \mathcal{H} which is called *strong topology* or *norm topology*.

In other words a sequence $(f)_{n \geq 0}$ in \mathcal{H} converges strongly to f if $\|f_n - f\| \rightarrow 0$ as $n \rightarrow \infty$.

Definition A.1.3. A map L from a linear space to another is called *linear operator* if it satisfies

$$L(\eta f + \zeta g) = \eta Lf + \zeta Lg$$

for $\eta, \zeta \in \mathbb{C}$ and $f, g \in \mathcal{H}$.

The operator L is continuous if L is continuous at each point of domain of L . To be able to speak of continuity of a linear operator, the domain and range spaces have to be equipped with respective topologies.

Let L be a linear operator from \mathcal{H} to \mathcal{G} with \mathcal{H} and \mathcal{G} are pre-Hilbert spaces. The *Lipschitz constant* for a linear operator L is called its norm and denoted by $\|L\|$ where

$$\|L\| = \sup\{\|Lf\|_{\mathcal{G}}/\|f\|_{\mathcal{H}} : 0 \neq f \in \mathcal{H}\}. \quad (\text{A.7})$$

Theorem A.1.4. *Let L be a linear operator from \mathcal{H} to \mathcal{G} where \mathcal{H} and \mathcal{G} are pre-Hilbert spaces. The followings are mutually equivalent:*

(i) L is continuous

(ii) L is bounded in the sense that

$$\sup\{\|Lf\|_{\mathcal{G}} : \|f\|_{\mathcal{H}} \leq \sigma\} < \infty$$

for $0 \leq \sigma < \infty$.

(iii) L is Lipschitz continuous in the sense that there is $0 \leq \gamma < \infty$ such that

$$\|Lf - Lg\|_{\mathcal{G}} \leq \gamma \|f - g\|_{\mathcal{H}},$$

for $f, g \in \mathcal{H}$.

Proof. (i) \implies (ii)

Let L be continuous. Define a neighborhood of 0 in \mathcal{G} as $v = \{g : \|g\|_{\mathcal{G}} \leq 1\}$. Continuity of L implies $L0 = 0$ and there is a neighborhood (of 0 in \mathcal{H}) of the form $u = \{f : \|f\|_{\mathcal{H}} \leq \delta\}$ which is mapped into v by L , where $\delta > 0$. Now let

us take a nonzero $f \in \mathcal{H}$ such that $\|f\|_{\mathcal{H}} \leq \rho$, for $\rho > 0$. Since $\|\delta f/\rho\|_{\mathcal{H}} \leq \delta$, then $\frac{\delta f}{\rho} \in u$. As L is linear,

$$\delta \|Lf\|_{\mathcal{G}}/\rho = \|L(\delta f/\rho)\|_{\mathcal{G}} \leq 1$$

and

$$\sup\{\|Lf\|_{\mathcal{G}} : \|f\|_{\mathcal{H}} \leq \rho\} \leq \frac{\rho}{\delta}.$$

(ii) \implies (iii)

Let L satisfies (ii). Consider the norm of $Lf - Lg$. By linearity property of L ,

$$\|Lf - Lg\|_{\mathcal{G}} = \|L(f - g)\|_{\mathcal{G}} \leq \gamma \|f - g\|_{\mathcal{H}} \quad \text{for } f, g \in \mathcal{H},$$

where γ is the supremum in (ii) with $\rho = 1$.

(iii) \implies (i)

This is obvious since boundedness implies continuity. □

Denote by $B(\mathcal{H}, \mathcal{G})$, all continuous linear operators from \mathcal{H} to \mathcal{G} , \mathcal{H} and \mathcal{G} are pre-Hilbert spaces. $B(\mathcal{H}, \mathcal{G})$ becomes a linear space with respect to the natural addition and scalar multiplication:

$$(\xi L + \zeta M)f = \xi Lf + \zeta Mf,$$

where L and M are linear operators, $f \in \mathcal{H}$ and $\xi \in \mathbb{C}$.

When $\mathcal{H} = \mathcal{G}$, $B(\mathcal{H}, \mathcal{G})$ is denoted by $B(\mathcal{H})$.

Let \mathcal{K} be another pre-Hilbert space. If $L \in B(\mathcal{H}, \mathcal{G})$ and $K \in B(\mathcal{G}, \mathcal{K})$, then their product will be $KL \in B(\mathcal{H}, \mathcal{K})$ and defined by

$$(KL)f = K(Lf) \quad \text{for } f \in \mathcal{H}.$$

Moreover we have the following properties:

(i) $K(\xi L + \zeta M) = \xi KL + \zeta KM$

$$(ii) \quad \|\xi L\| = |\xi| \cdot \|L\|$$

$$(iii) \quad \|L + M\| \leq \|L\| + \|M\| \quad \text{and}$$

$$(iv) \quad \|KL\| \leq \|K\| \|L\|.$$

For \mathcal{H}, \mathcal{G} are pre-Hilbert spaces and L, M are linear operators, $B(\mathcal{H}, \mathcal{G})$ becomes a metric space with respect to the translation invariant, positively homogeneous distance function

$$d(L, M) := \|L - M\|.$$

For each $K \in B(\mathcal{G}, \mathcal{K})$, the map $L \mapsto KL$ becomes a continuous linear operator from $B(\mathcal{H}, \mathcal{G})$ to $B(\mathcal{H}, \mathcal{K})$.

Definition A.1.5. A linear operator from \mathcal{H} to the scalar field \mathbb{C} is called a *linear functional* (or *linear form*).

Definition A.1.6. A pre-Hilbert space \mathcal{H} is called a *Hilbert space* if it is complete in metric. In other words, if f_n is a Cauchy sequence in \mathcal{H} , i.e. $\|f_n - f_m\| \rightarrow 0$ when $n, m \rightarrow \infty$, then there is $f \in \mathcal{H}$ such that $\|f_n - f\| \rightarrow 0$ when $n \rightarrow \infty$.

Every subspace of a pre-Hilbert space is again a pre-Hilbert space with respect to the induced inner product; but for a subspace of a Hilbert space to be again a Hilbert space, it has to be closed.

Let \mathcal{H} be a pre-Hilbert space. We can imbed \mathcal{H} as a dense subspace of a Hilbert space $\tilde{\mathcal{H}}$ in such a way that

$$\langle f, g \rangle_{\tilde{\mathcal{H}}} = \langle f, g \rangle_{\mathcal{H}} \quad \text{for } f, g \in \mathcal{H}.$$

This canonical method is called *completion*. If we have a continuous linear operator L from a dense subspace \mathcal{M} of a Hilbert space \mathcal{H} to a Hilbert space \mathcal{G} , we can extend L uniquely to a continuous linear operator from \mathcal{H} to \mathcal{G} with preserving norm.

Theorem A.1.7. *Let \mathcal{M} be a dense subspace of a Hilbert space \mathcal{H} , and \mathcal{N} a dense subspace of a Hilbert space \mathcal{G} . If a linear operator L from \mathcal{M} to \mathcal{G} satisfies*

$$|\langle Lf, g \rangle_{\mathcal{G}}| \leq |\gamma| \|f\|_{\mathcal{H}} \|g\|_{\mathcal{G}} \quad \text{for } f \in \mathcal{H}, g \in \mathcal{M} \quad \text{with some } 0 \leq \gamma < \infty, \quad (\text{A.8})$$

then it is uniquely extended to a continuous linear operator from \mathcal{M} to \mathcal{G} with norm $\leq \gamma$. The minimum of such γ coincides with norm.

Proof. Let L be a linear operator from M to \mathcal{G} and satisfies (A.8). By fixing f , since $g \mapsto \langle Lf, g \rangle_{\mathcal{G}}$ and $g \mapsto \|g\|_{\mathcal{G}}$ are continuous, we can assume that (A.8) is satisfied for all $g \in \mathcal{G}$.

Now by using (A.8) and Schwarz inequality, we get

$$\|Lf\|_{\mathcal{G}}^2 = \langle Lf, Lf \rangle_{\mathcal{G}} \leq \gamma \|f\|_{\mathcal{H}} \|Lf\|_{\mathcal{G}}$$

which implies

$$\|Lf\|_{\mathcal{G}} \leq \gamma \|f\|_{\mathcal{H}} \quad \text{for all } f \in M.$$

Hence L is continuous operator from M to \mathcal{G} with norm $\leq \gamma$. So we can extend L uniquely to a continuous linear operator from \mathcal{H} to \mathcal{G} with preserving norm. It is obvious that minimum of γ coincides with norm. \square

Theorem A.1.8. Let $L^2(\Omega, \mu)$ consists of all measurable functions $f(\omega)$ on Ω such that

$$\int_{\Omega} |f(\omega)|^2 d\mu(\omega) < \infty, \quad (\text{A.9})$$

where (Ω, μ) is a measure space such that Ω is the union of subsets of finite positive measure. Then $L^2(\Omega, \mu)$ becomes a Hilbert space with respect to the inner product

$$\langle f, g \rangle := \int_{\Omega} f(\omega) \overline{g(\omega)} d\mu(\omega). \quad (\text{A.10})$$

Proof. We have

$$|f(\omega) + g(\omega)|^2 \leq 2\{|f(\omega)|^2 + |g(\omega)|^2\}$$

which implies that $L^2(\Omega, \mu)$ is a linear space. If we expand the left handside, we have

$$|f(\omega) + g(\omega)|^2 = |f(\omega)|^2 + |g(\omega)|^2 + 2f(\omega)g(\omega) \leq 2\{|f(\omega)|^2 + |g(\omega)|^2\}$$

which implies that

$$|f(\omega)\overline{g(\omega)}| \leq \frac{1}{2}\{|f(\omega)|^2 + |g(\omega)|^2\}.$$

So, $\langle f, g \rangle$ in (A.10) is well-defined and the linear space $L^2(\Omega, \mu)$ becomes a pre-Hilbert space.

Completeness:

Let us take a Cauchy sequence $(f_n)_{n \geq 1}$ in $L^2(\Omega, \mu)$. Suppose that

$$\sum_{n=1}^{\infty} \|f_{n+1} - f_n\| \equiv \gamma < \infty.$$

For any m , consider the following integration,

$$\int_{\Omega} \sum_{n=1}^m |f_{n+1}(\omega) - f_n(\omega)|^2 d\mu(\omega) \leq \sum_{n=1}^m \|f_{n+1} - f_n\|^2 = \gamma^2 < \infty.$$

Then by the Lebesgue theorem,

$$\int_{\Omega} \sum_{n=1}^{\infty} |f_{n+1}(\omega) - f_n(\omega)|^2 d\mu(\omega) \leq \gamma^2 < \infty$$

which implies

$$g(\omega) \equiv \sum_{n=1}^m |f_{n+1}(\omega) - f_n(\omega)| < \infty \quad \text{for almost all } \omega,$$

and

$$\int_{\Omega} g(\omega)^2 d\mu(\omega) = \int_{\Omega} \sum_{n=1}^{\infty} |f_{n+1}(\omega) - f_n(\omega)|^2 d\mu(\omega) \leq \gamma^2 < \infty.$$

Hence for almost all ω , the sequence $(f_n(\omega))_{n \geq 1}$ converges. To find the limit of $f(\omega)$, consider the following

$$\begin{aligned} |f_{n+1}(\omega) - f_n(\omega)|^2 &\leq \sum_{k=n}^{\infty} |f_{k+1}(\omega) - f_k(\omega)|^2 \\ &\leq \sum_{n=1}^m |f_{n+1}(\omega) - f_n(\omega)| = g(\omega) \end{aligned}$$

and again by Lebesgue theorem,

$$\begin{aligned} \lim_{n \rightarrow \infty} \|f_n - f\|^2 &= \lim_{n \rightarrow \infty} \langle f_n - f, f_n - f \rangle = \lim_{n \rightarrow \infty} \int_{\Omega} |f_n(\omega) - f(\omega)|^2 d\mu(\omega) \\ &\leq \int_{\Omega} \lim_{n \rightarrow \infty} |f_n(\omega) - f(\omega)|^2 d\mu(\omega) = 0. \end{aligned}$$

Therefore f_n converges strongly to f , which means $L^2(\Omega, \mu)$ is complete and it is a Hilbert space with respect to the given inner product. \square

Let \mathcal{H} and \mathcal{G} be Hilbert spaces. The algebraic direct sum of \mathcal{H} and \mathcal{G} , consisting of $f \oplus g$ with the functions $f \in \mathcal{H}$ and $g \in \mathcal{G}$, becomes a Hilbert space, denoted by $\mathcal{H} \oplus \mathcal{G}$ with respect to the inner product

$$\langle f \oplus g, f' \oplus g' \rangle = \langle f, f' \rangle_{\mathcal{H}} + \langle g, g' \rangle_{\mathcal{G}}.$$

Let \mathcal{H} and \mathcal{G} be Hilbert spaces. The algebraic tensor product of \mathcal{H} and \mathcal{G} , consisting of finite sums of $f \otimes g$ with $f \in \mathcal{H}$ and $g \in \mathcal{G}$, becomes a pre-Hilbert space with respect to the inner product

$$\left\langle \sum_i f_i \otimes g_i, \sum_j f'_j \otimes g'_j \right\rangle = \sum_{i,j} \langle f_i, f'_j \rangle_{\mathcal{H}} \cdot \langle g_i, g'_j \rangle_{\mathcal{G}}.$$

Definition A.1.9. The canonical completion of the algebraic tensor product of \mathcal{H} and \mathcal{G} , where \mathcal{H} and \mathcal{G} are Hilbert spaces, is called *tensor product Hilbert space* and denoted by $\mathcal{H} \otimes \mathcal{G}$.

A.2 Projection

We start this section with an important theorem.

Theorem A.2.1 (F.Riesz). *For each continuous linear functional ϕ on a Hilbert space \mathcal{H} , there exists uniquely $g \in \mathcal{H}$ such that*

$$\phi(f) = \langle f, g \rangle \text{ for } f \in \mathcal{H}. \tag{A.11}$$

Proof. For the case $\phi = 0$, take $g = 1$, then we are done.

Let $\phi \neq 0$. For this case, we can suppose that $\|\phi\| = 1$. Then there exists a sequence g_n such that

$$\|g_n\| = 1 \quad (n = 1, 2, \dots) \quad \text{and} \quad \phi(g_n) \longrightarrow \|\phi\| = 1 \text{ as } n \rightarrow \infty.$$

Then by parallelogram rule

$$\|g_m - g_n\|^2 + \|g_m + g_n\|^2 = 2\|g_m\|^2 + 2\|g_n\|^2 = 2 + 2 = 4.$$

On the other hand by triangular rule,

$$2 = \|g_m\| + \|g_n\| \geq \|g_m + g_n\|,$$

then we get

$$2 \geq \|g_m + g_n\| \geq |\phi(g_m) + \phi(g_n)| \longrightarrow 2 \quad \text{as } m, n \rightarrow \infty.$$

Thus,

$$\|g_m + g_n\| = 2 \quad \text{and} \quad \|g_m - g_n\|^2 = 4. \text{ So, } \|g_m - g_n\| \longrightarrow 0 \quad \text{as } m, n \rightarrow \infty.$$

Then, due to the metric completeness, g_n converges strongly to some g such that $\|g\| = \phi(g) = 1$.

Let $f \in \mathcal{H}$ be arbitrary. If $\langle f, g \rangle = 0$, then for all $\zeta \in \mathbb{C}$,

$$\|g + \zeta f\|^2 = \|g\|^2 + \bar{\zeta}\langle g, f \rangle + \zeta\langle f, g \rangle + |\zeta|^2\|f\|^2 = 1 + |\zeta|^2\|f\|^2$$

and

$$\begin{aligned} \|g + \zeta f\|^2 &\geq |\phi(g + \zeta f)|^2 = |\phi(g) + \zeta\phi(f)|^2 = |1 + \zeta\phi(f)|^2 \\ &= 1 + 2\operatorname{Re}\zeta\phi(f) + |\zeta|^2|\phi(f)|^2. \end{aligned}$$

Combining these equations, we get

$$1 + |\zeta|^2\|f\|^2 \geq 1 + 2\operatorname{Re}\zeta\phi(f) + |\zeta|^2|\phi(f)|^2.$$

Since ζ is arbitrary, then $\phi(f) = 0$. This implies

$$\phi(f) = \langle f, g \rangle \quad \text{for } f \text{ satisfying } \langle f, g \rangle = 0.$$

Now consider the other case which is $\langle f, g \rangle \neq 0$. Define $h := f - \langle f, g \rangle g$. It follows that,

$$\langle h, g \rangle = \langle f - \langle f, g \rangle g, g \rangle = \langle f, g \rangle - \langle \langle f, g \rangle g, g \rangle = \langle f, g \rangle - \langle f, g \rangle = 0.$$

Then,

$$0 = \phi(h) = \phi(f) - \langle f, g \rangle \phi(g) = \phi(f) - \langle f, g \rangle.$$

Hence we obtain $\phi(f) = \langle f, g \rangle$.

To see the uniqueness of g , suppose that there exists $g' \in \mathcal{H}$ such that for each linear functional ϕ , we have

$$\phi(f) = \langle f, g' \rangle.$$

Also this equality is satisfied for g . Thus we have

$$\phi(f) = \langle f, g' \rangle = \langle f, g \rangle$$

which implies

$$\langle f, g - g' \rangle = 0.$$

Hence $g = g'$. □

Theorem A.2.2. *For any closed subspace \mathcal{M} of a Hilbert space \mathcal{H} ,*

$$\mathcal{H} = \mathcal{M} \oplus \mathcal{M}^\perp \text{ (algebraic direct sum),}$$

that is, each $f \in \mathcal{H}$ can be uniquely written in the form

$$f = f_{\mathcal{M}} + f_{\mathcal{M}^\perp} \text{ with } f_{\mathcal{M}} \in \mathcal{M}, f_{\mathcal{M}^\perp} \in \mathcal{M}^\perp. \quad (\text{A.12})$$

Moreover, $\|f_{\mathcal{M}}\|$ coincides with the distance from f to \mathcal{M}^\perp ;

$$\|f_{\mathcal{M}}\| = \min\{\|f - g\| : g \in \mathcal{M}^\perp\}, \quad (\text{A.13})$$

the distance being attained only at $f_{\mathcal{M}^\perp}$.

Proof. Since \mathcal{M} is closed, it is also a Hilbert space. By Riesz theorem to the continuous functional $h \mapsto \langle h, f \rangle$ restricted to \mathcal{M} , there is $f_{\mathcal{M}} \in \mathcal{M}$, such that

$$\langle h, f \rangle = \langle h, f_{\mathcal{M}} \rangle \text{ for } h \in \mathcal{M}.$$

By this, we get $\langle h, f - f_{\mathcal{M}} \rangle = 0$. Let us define $f_{\mathcal{M}^\perp} := f - f_{\mathcal{M}}$. Since $\langle h, f - f_{\mathcal{M}} \rangle = 0$, then $f_{\mathcal{M}^\perp} \perp h$ for all $h \in \mathcal{M}$. So, $f_{\mathcal{M}^\perp} \in \mathcal{M}^\perp$. Hence

$$f = f_{\mathcal{M}} + f_{\mathcal{M}^\perp} \text{ where } f_{\mathcal{M}} \in \mathcal{M}, f_{\mathcal{M}^\perp} \in \mathcal{M}^\perp.$$

For the uniqueness, let $f = f'_M + f'_{M^\perp}$ be any other decomposition of f , where $f'_M \in \mathcal{M}$, $f'_{M^\perp} \in \mathcal{M}^\perp$. Then,

$$f = f'_M + f'_{M^\perp} = f_M + f_{M^\perp}$$

which implies that

$$f_M - f'_M = f_{M^\perp} - f'_{M^\perp}$$

and by the last equality we obtain that $f_M = f'_M$.

For the moreover part, let us take any $g \in \mathcal{M}^\perp$. Since $\langle f_{M^\perp} - g, f \rangle = 0$, we get

$$\|f - g\|^2 = \|f_M + f_{M^\perp} - g\|^2 = \|f_M\|^2 + \|f_{M^\perp} - g\|^2.$$

Then we have

$$\|f - g\|^2 \geq \|f_M\|^2 \quad \text{and equality occurs when } g = f_{M^\perp}.$$

Hence,

$$\|f_M\| = \min\{\|f - g\| : g \in \mathcal{M}^\perp\}.$$

□

Corollary A.2.3. *In a Hilbert space, the closed linear span of any subset A coincides with $(A^\perp)^\perp$.*

Proof. Let \mathcal{M} be the closed linear span of A . So, we have $\mathcal{M} \subseteq (A^\perp)^\perp$. It remains to show the other side of inclusion. As $(A^\perp)^\perp$ is a closed subspace of \mathcal{H} , so it is also a Hilbert space. Now by using the Theorem A.2.2, for each $f \in (A^\perp)^\perp$, f has a decomposition such that $f = g + h$ where $g \in \mathcal{M}$, $h \in (A^\perp)^\perp$ and h is orthogonal to all of \mathcal{M} . Since $A \subseteq \mathcal{M}$, h must be orthogonal to all of A . In other words, h must be in A^\perp . But h is both in A^\perp and $(A^\perp)^\perp$. Hence $h = 0$ and $h = 0 = f - g$ implies $f = g \in \mathcal{M}$ i.e. $(A^\perp)^\perp \subseteq \mathcal{M}$. Consequently, $\mathcal{M} = (A^\perp)^\perp$. □

Definition A.2.4. A subset of a Hilbert space \mathcal{H} is said to be *total (complete)* in \mathcal{H} if 0 is the only element that is orthogonal to all of A , i.e. $A^\perp = \{0\}$.

So by the Corollary A.2.3, A is total if and only if every element of \mathcal{H} can be approximated by linear combinations of elements of A .

Definition A.2.5. When \mathcal{M} is a closed subspace of \mathcal{H} , the map $f \mapsto f_{\mathcal{M}}$ determines a linear operator from \mathcal{H} to \mathcal{M} with norm ≤ 1 . This operator is called the *orthogonal projection* to \mathcal{M} and is denoted by $P_{\mathcal{M}}$.

If I is the identity operator, then $I - P_{\mathcal{M}}$ becomes the orthogonal projection to \mathcal{M}^{\perp} and

$$\|f\|^2 = \|P_{\mathcal{M}}f\|^2 + \|(I - P_{\mathcal{M}})f\|^2. \quad (\text{A.14})$$

Theorem A.2.6 (Abstract Interpolation). Let \mathcal{H} be a Hilbert space, X an index set and γ a positive number. Given a map $x \mapsto f_x$ from X to \mathcal{H} and one $x \mapsto a_x$ from X to \mathbb{C} , there exists an element $g \in \mathcal{H}$ such that

$$\|g\| \leq \gamma \quad \text{and} \quad \langle f_x, g \rangle = a_x \quad \text{for} \quad x \in X, \quad (\text{A.15})$$

if and only if

$$\sum_{x,y} \bar{\xi}_y \xi_x \bar{a}_y a_x \leq \gamma^2 \sum_{x,y} \bar{\xi}_y \xi_x \langle f_x, f_y \rangle \quad \text{for} \quad [\xi_x], \quad (\text{A.16})$$

where $[\xi_x]$ denotes, in general, a function on X which vanishes outside a finite number of points.

Proof. Suppose that (A.15) is satisfied for $g \in \mathcal{H}$. Then

$$\sum_{x,y} \bar{\xi}_y \xi_x \bar{a}_y a_x = \left| \sum_x \xi_x a_x \right|^2 = \left| \sum_x \xi_x \langle f_x, g \rangle \right|^2 = \left| \langle \sum_x \xi_x f_x, g \rangle \right|^2,$$

by the Schwarz inequality

$$\leq \left\| \sum_x \xi_x f_x \right\|^2 \|g\|^2 \leq \gamma^2 \left\| \sum_x \xi_x f_x \right\|^2 = \gamma^2 \sum_{x,y} \bar{\xi}_y \xi_x \langle f_x, f_y \rangle.$$

So (A.16) is satisfied.

Now suppose (A.16) is satisfied. Then the correspondence

$$\sum_x \xi_x f_x \mapsto \sum_x \xi_x a_x$$

defines a linear functional on the linear span of $\{f_x : x \in X\}$ with norm $\leq \gamma$. This bounded functional can be extended to the closure of this linear span, and then by using Riesz's Representation Theorem (see Theorem A.2.1) the vector g can be found such that $\|g\| \leq \gamma$ and $\langle f_x, g \rangle = a_x$ for all $x \in X$. \square

A.3 Weak Topology

Definition A.3.1. *The weak topology* of a Hilbert space \mathcal{H} is the weakest topology that makes continuous all linear functionals of the form $f \mapsto \langle f, g \rangle$.

For an element f in the weak topology, a fundamental system of neighborhoods of f consists of subsets of the form

$$U(f; A, \epsilon) = \{h : |\langle f, g \rangle - \langle h, g \rangle| < \epsilon \text{ for } g \in A\},$$

where A is a finite subset of H and ϵ is a positive number. Then a directed net $\{f_\lambda\}$ converges weakly to f if and only if

$$\langle f_\lambda, g \rangle \xrightarrow{\lambda} \langle f, g \rangle$$

for all $g \in \mathcal{H}$.

Definition A.3.2. *The (operator) weak topology* in the space $B(\mathcal{H}, \mathcal{G})$ of continuous linear operators from \mathcal{H} to a Hilbert space \mathcal{G} , is the weakest topology that makes continuous all linear functionals of the form $L \mapsto \langle Lf, g \rangle$ for $f \in \mathcal{H}$ and $g \in \mathcal{G}$.

Then a directed net $\{L_\lambda\}$ converges (operator) weakly to L if and only if

$$\langle L_\lambda f, g \rangle \xrightarrow{\lambda} \langle Lf, g \rangle.$$

Definition A.3.3. *The (operator) strong topology* is the weakest topology that makes continuous all linear operators of the form $L \mapsto Lf$ for $f \in \mathcal{H}$.

Then a directed net $\{L_\lambda\}$ converges (operator) strongly to L if and only if

$$\|L_\lambda f - Lf\| \xrightarrow{\lambda} 0$$

for all $f \in \mathcal{H}$.

Theorem A.3.4 ([1]). *The closed unit ball $U := \{f : \|f\| \leq 1\}$ of a Hilbert space \mathcal{H} is weakly compact.*

Theorem A.3.5 ([1]). *The closed unit ball $\{L : \|L\| \leq 1\}$ of $B(\mathcal{H}, \mathcal{G})$ is (operator) weakly compact.*

Theorem A.3.6. *If a subset A of a Hilbert space \mathcal{H} is weakly bounded in the sense*

$$\sup_{f \in A} |\langle f, g \rangle| < \infty \quad \text{for } g \in \mathcal{H}, \quad (\text{A.17})$$

then it is strongly bounded, that is, $\sup_{f \in A} \|f\| < \infty$.

Proof. Define

$$B_n := \{g : |\langle f, g \rangle| \leq n \text{ for } f \in A\}, \quad (n = 1, 2, 3, \dots).$$

Then for each n , B_n will be a closed subset of \mathcal{H} and so $\mathcal{H} = \bigcup_n B_n$. By Baire category theorem, since \mathcal{H} is a complete metric space with respect to the strong topology and \mathcal{H} is covered by a countable number of closed sets, then at least one of them has nonempty interior, say B_{n_0} . So, for some g_0 and $\epsilon > 0$,

$$V \equiv \{g : \|g - g_0\| \leq \epsilon\} \subset B_{n_0}$$

Let g be in \mathcal{H} . Since

$$\|g_0 + \epsilon g / \|g\| - g_0\| = \|\epsilon g / \|g\|\| \leq \epsilon,$$

then $g_0 + \epsilon g / \|g\|$ will be in V . Then

$$\epsilon |\langle f, g \rangle| / \|g\| \leq |\langle f, g_0 + \epsilon g / \|g\| \rangle| + |\langle f, g_0 \rangle| \leq n_0 + n_0 = 2n_0 \quad \text{for } f \in A.$$

This implies,

$$|\langle f, g \rangle| \leq 2n_0 \|g\| / \epsilon \quad \text{for } f \in A.$$

Putting f instead of g in the last inequality and using Schwarz inequality, we get

$$\sup_{f \in A} \leq 2n_0/\epsilon < \infty.$$

□

Theorem A.3.7. *Let \mathcal{H} and \mathcal{G} be Hilbert spaces. For a linear operator L from \mathcal{H} to \mathcal{G} , strong continuity and weak continuity are the same thing.*

Proof. Let L be weakly continuous. Since by Theorem A.3.4 the unit ball U of a Hilbert space is weakly compact, it follows that $L(U)$ is also weakly compact and so weakly bounded in G . Since Theorem A.3.6 states that weakly boundedness implies strongly boundedness, $L(U)$ will be strongly bounded and hence it is strongly continuous by Theorem A.1.4

Now suppose that L is strongly continuous. So we have, for each $g \in \mathcal{G}$, the linear functional $f \mapsto \langle Lf, g \rangle_{\mathcal{G}}$ is continuous on \mathcal{H} . Now by Riesz theorem, there is $g' \in \mathcal{H}$ such that,

$$f \mapsto \langle f, g' \rangle_{\mathcal{H}} \text{ for } f \in \mathcal{H},$$

then

$$\langle Lf, g \rangle_{\mathcal{G}} = \langle f, g' \rangle_{\mathcal{H}} \text{ for } f \in \mathcal{H}.$$

Hence L is weakly continuous. □

Theorem A.3.8 ([1]). *Let \mathcal{H} and \mathcal{G} be Hilbert spaces. Then the following assertions for a subset \mathbf{L} of $B(\mathcal{H}, \mathcal{G})$ are mutually equivalent:*

(i) \mathbf{L} is (operator) weakly bounded;

$$\sup_{L \in \mathbf{L}} |\langle Lf, g \rangle| < \infty \text{ for } f \in \mathcal{H}, \quad g \in \mathcal{G}.$$

(ii) \mathbf{L} is (operator) strongly bounded;

$$\sup_{L \in \mathbf{L}} \|Lf\| < \infty \text{ for } f \in \mathcal{H}.$$

(iii) \mathbf{L} is norm bounded (or uniformly bounded);

$$\sup_{L \in \mathbf{L}} \|L\| < \infty.$$

Definition A.3.9. A linear operator L from \mathcal{H} to \mathcal{G} , where \mathcal{H} and \mathcal{G} are Hilbert spaces, is said to be *closed* if its graph

$$G_L := \{f \oplus Lf : f \in \mathcal{H}\} \quad (\text{A.18})$$

is a closed subspace of the direct sum space $\mathcal{H} \oplus \mathcal{G}$, that is, if $\|f_n - f\| \rightarrow 0$ in \mathcal{H} as $(n \rightarrow \infty)$ and $\|Lf_n - g\| \rightarrow 0$ in \mathcal{G} as $(n \rightarrow \infty)$, then $g = Lf$.

Theorem A.3.10. *Any closed linear operator L from a Hilbert space \mathcal{H} to another Hilbert space \mathcal{G} is continuous.*

Proof. Let G_L be the graph of L . Since L is closed, then G_L is a closed subspace and so it is a Hilbert space. Consider the linear operator $\phi : f \oplus Lf \mapsto f$ from G_L to \mathcal{H} . ϕ is a bijection because of closedness of G_L . Let U_L be the unit ball of G_L . By Theorem A.3.4 U_L is weakly compact. This implies $\phi(U_L)$ is weakly compact and so is $\bigcup_{n=1}^{\infty} n\phi(U_L) = \mathcal{H}$. Now by Baire theorem, $\phi(U_L)$ has non-empty interior. By the proof of Theorem A.3.6, there exists $0 \leq \gamma < \infty$ such that

$$\|f \oplus Lf\| \leq \gamma \|f\| \text{ for } f \in \mathcal{H}.$$

This implies

$$\|f \oplus Lf\|^2 = \langle \|f \oplus Lf\|, \|f \oplus Lf\| \rangle = \langle f, f \rangle + \langle Lf, Lf \rangle = \|f\|^2 + \|Lf\|^2 \leq \gamma^2 \|f\|^2.$$

Then we obtain

$$\|Lf\| \leq (\gamma^2 - 1)^{1/2} \|f\|.$$

Hence L is continuous. □

Corollary A.3.11 ([1]). *Every continuous linear bijection between Hilbert spaces has continuous inverse.*

A.4 Self-adjoint Operators

Definition A.4.1. Let \mathcal{H} and \mathcal{G} are Hilbert spaces. A function $\Phi : \mathcal{H} \times \mathcal{G} \rightarrow \mathbb{C}$ is a *sesquilinear form (or function)* if for $f, h \in \mathcal{H}$, $g, k \in \mathcal{G}$ and $\alpha, \beta \in \mathbb{C}$,

$$(i) \quad \Phi(\alpha f + \beta h, g) = \alpha \Phi(f, g) + \beta \Phi(h, g) \quad (\text{A.19})$$

$$(ii) \quad \Phi(f, \alpha g + \beta k) = \bar{\alpha}\Phi(f, g) + \bar{\beta}\Phi(f, k) \quad (\text{A.20})$$

are satisfied.

If $L \in B(\mathcal{H}, \mathcal{G})$, then the sesquilinear form Φ is defined as

$$\Phi(f, g) = \langle Lf, g \rangle_{\mathcal{G}} \quad (\text{A.21})$$

and it is bounded,

$$|\Phi(f, g)| \leq \gamma \|f\|_{\mathcal{H}} \|g\|_{\mathcal{G}} \quad \text{for } f \in \mathcal{H}, g \in \mathcal{G}, \quad (\text{A.22})$$

where γ can take any value not less than $\|L\|$.

If we are given a sesqui-linear form Φ satisfies boundedness condition (A.22), then for fixed $f \in \mathcal{H}$, the linear functional $g \mapsto \overline{\Phi(f, g)}$ is continuous on \mathcal{G} . Applying Riesz theorem, there exists uniquely $f' \in \mathcal{G}$ such that

$$\|f'\|_{\mathcal{G}} \leq \gamma \|f\|_{\mathcal{H}} \quad \text{and} \quad \Phi(f, g) = \langle f', g \rangle_{\mathcal{G}} \quad \text{for } g \in \mathcal{G}.$$

Then $f \mapsto f'$ becomes linear, so we obtain $\Phi(f, g) = \langle f', g \rangle_{\mathcal{G}} = \langle Lf, g \rangle_{\mathcal{G}}$.

Definition A.4.2. If $L \in B(\mathcal{H}, \mathcal{G})$, then there exists a unique operator in $B(\mathcal{G}, \mathcal{H})$, called *adjoint* of L , denoted by L^* satisfying

$$\Phi(f, g) = \langle f, L^*g \rangle_{\mathcal{H}} \quad \text{for } f \in \mathcal{H}, g \in \mathcal{G}. \quad (\text{A.23})$$

By the definition of L and L^* , we have the following relation,

$$\langle Lf, g \rangle_{\mathcal{G}} = \langle f, L^*g \rangle_{\mathcal{H}} \quad \text{for } f \in \mathcal{H}, g \in \mathcal{G}. \quad (\text{A.24})$$

Some other properties of adjoint operator:

$$\|L\| = \|L^*\|. \quad (\text{A.25})$$

Given \mathcal{H}, \mathcal{G} and \mathcal{K} are Hilbert spaces, if $K \in B(\mathcal{G}, \mathcal{K})$ and $L \in B(\mathcal{H}, \mathcal{G})$, then

$$KL \in B(\mathcal{H}, \mathcal{K}) \quad \text{and} \quad (KL)^* = L^*K^*. \quad (\text{A.26})$$

$$\ker L = (\text{Ran}L^*)^\perp \text{ and } (\ker L)^\perp = \text{closure}\{\text{Ran}L^*\} \quad (\text{A.27})$$

where $\ker L$ is the kernel of L and $\text{Ran}L$ is the range of L .

Theorem A.4.3 ([1]). *For $L, M \in B(\mathcal{H}, \mathcal{G})$, the following assertions are mutually equivalent.*

(i) $\text{Ran}(M) \subseteq \text{Ran}(L)$.

(ii) *There exists $K \in B(\mathcal{H})$ such that $M = LK$.*

(iii) *There exists $0 \leq \gamma < \infty$ such that*

$$\|M^*g\| \leq \gamma\|L^*g\| \quad \text{for } g \in \mathcal{G}.$$

Definition A.4.4. Let \mathcal{H} be a Hilbert space. A function $\phi : \mathcal{H} \rightarrow \mathbb{C}$ is a *quadratic form* if for all $f \in \mathcal{H}$ and $\zeta \in \mathbb{C}$,

$$\phi(\zeta f) = |\zeta|^2 \phi(f) \quad (\text{A.28})$$

and

$$\phi(f+g) + \phi(f-g) = 2\{\phi(f) + \phi(g)\} \quad (\text{A.29})$$

are satisfied.

For $L \in B(\mathcal{H})$, the quadratic form ϕ on \mathcal{H} is defined as

$$\phi(f) = \langle Lf, f \rangle \text{ for } f \in \mathcal{H}, \quad (\text{A.30})$$

and it is bounded

$$|\phi(f)| \leq \gamma\|f\|^2 \text{ for } f \in \mathcal{H}, \quad (\text{A.31})$$

where γ can take any value not less than $\|L\|$.

The sesqui-linear form Φ associated with L can be recaptured from quadratic form ϕ by the following equation:

$$\Phi(f, g) = \frac{1}{4}\{\phi(f+g) - \phi(f-g)\} + \{\phi(f+ig) - \phi(f-ig)\} \quad (\text{A.32})$$

for all $f, g \in \mathcal{H}$.

Definition A.4.5. A continuous linear operator L on a Hilbert space \mathcal{H} is said to be *self-adjoint* if $L = L^*$.

Remark A.4.6. L is self adjoint if and only if the associated sesqui-linear form Φ is Hermitian.

Remark A.4.7. If L is self-adjoint, norm of L coincides with the minimum of γ in (A.31) for the associated quadratic form.

Theorem A.4.8. *If L is a continuous self-adjoint operator, then*

$$\|L\| = \sup\{|\langle Lf, f \rangle| : \|f\| \leq 1\}. \quad (\text{A.33})$$

Proof. Let $\gamma = \sup\{|\langle Lf, f \rangle| : \|f\| \leq 1\}$.

It is obvious that $\gamma \leq \|L\|$. Consider the reverse direction. By the Remark A.4.6, since L is self-adjoint, then associated quadratic form is Hermitian, so real-valued.

Let fix $f, g \in \mathcal{H}$ and choose $e^{i\theta}$ such that

$$\langle L(e^{i\theta} f), g \rangle = |\langle Lf, g \rangle|.$$

Then using the equality (A.32), it follows that

$$\begin{aligned} |\langle Lf, g \rangle| &= \operatorname{Re} \langle L(e^{i\theta} f), g \rangle \\ &= \frac{1}{4} \{ \langle L(e^{i\theta} f + g), e^{i\theta} f + g \rangle - \langle L(e^{i\theta} f - g), e^{i\theta} f - g \rangle \} \\ &\leq \frac{\gamma}{4} \{ \|e^{i\theta} f + g\|^2 + \|e^{i\theta} f - g\|^2 \} = \frac{\gamma}{2} \{ \|f\|^2 + \|g\|^2 \}. \end{aligned}$$

Now replace f and g by λf and $\frac{g}{\lambda}$ respectively, and take the minimum in λ , we obtain

$$|\langle Lf, g \rangle| \leq \gamma \|f\| \|g\| \text{ for } f, g \in \mathcal{H}.$$

Hence we have

$$\|L\| \leq \gamma.$$

□

Definition A.4.9. A self-adjoint operator $L \in B(\mathcal{H})$ is said to be *positive definite* if $\langle Lf, f \rangle \geq 0$ for all $f \in \mathcal{H}$. If $\langle Lf, f \rangle = 0$ only when $f = 0$, then L is said to be strictly positive definite.

Remark A.4.10. For any positive definite operator $L \in B(\mathcal{H})$, a generalized Schwarz inequality holds:

$$|\langle Lf, g \rangle|^2 \leq \langle Lf, f \rangle \cdot \langle Lg, g \rangle. \quad (\text{A.34})$$

Theorem A.4.11. *Let L and M be continuous positive definite operators on \mathcal{H} and \mathcal{G} respectively. A continuous linear operator K from \mathcal{H} to \mathcal{G} satisfies the inequality*

$$|\langle Kf, g \rangle_{\mathcal{G}}|^2 \leq \langle Lf, f \rangle_{\mathcal{H}} \langle Mg, g \rangle_{\mathcal{G}} \quad \text{for } f \in \mathcal{H}, g \in \mathcal{G} \quad (\text{A.35})$$

if and only if the continuous linear operator

$$\begin{bmatrix} L & K^* \\ K & M \end{bmatrix}$$

on the direct sum Hilbert space $\mathcal{H} \oplus \mathcal{G}$:

$$f \oplus g \longmapsto (Lf + K^*g) \oplus (Kf + Mg)$$

is positive definite.

Proof. Suppose that (A.35) is satisfied. Let S stands for the matrix

$$\begin{bmatrix} L & K^* \\ K & M \end{bmatrix}$$

such that

$$\langle S(f \oplus g), f \oplus g \rangle = \langle Lf + K^*g, f \rangle_{\mathcal{H}} + \langle Kf + Mg, g \rangle_{\mathcal{G}},$$

then by using the property of adjoint operator and Schwarz inequality, the above equation follows as

$$\begin{aligned} \langle S(f \oplus g), f \oplus g \rangle &= \langle Lf, f \rangle_{\mathcal{H}} + \langle K^*g, f \rangle_{\mathcal{H}} + \langle Kf, g \rangle_{\mathcal{G}} + \langle Mg, g \rangle_{\mathcal{G}} \\ &= \langle Lf, f \rangle_{\mathcal{H}} + \langle g, Kf \rangle_{\mathcal{G}} + \langle Kf, g \rangle_{\mathcal{G}} + \langle Mg, g \rangle_{\mathcal{G}} \\ &= \langle Lf, f \rangle_{\mathcal{H}} + 2\text{Re}\langle Kf, g \rangle_{\mathcal{G}} + \langle Mg, g \rangle_{\mathcal{G}} \\ &\geq \langle Lf, f \rangle_{\mathcal{H}} + \langle Mg, g \rangle_{\mathcal{G}} - 2\sqrt{\langle Lf, f \rangle_{\mathcal{H}} \langle Mg, g \rangle_{\mathcal{G}}} \\ &= \{\sqrt{\langle Lf, f \rangle_{\mathcal{H}}} - \sqrt{\langle Mg, g \rangle_{\mathcal{G}}}\}^2 \geq 0. \end{aligned}$$

Thus S is positive definite.

Conversely, let S be positive definite. By the above relation we have

$$\begin{aligned} 0 \leq \langle S(f \oplus g), f \oplus g \rangle &= \langle Lf + K^*g, f \rangle_{\mathcal{H}} + \langle Kf + Mg, g \rangle_{\mathcal{G}} \\ &= \langle Lf, f \rangle_{\mathcal{H}} + \langle K^*g, f \rangle_{\mathcal{H}} + \langle Kf, g \rangle_{\mathcal{G}} + \langle Mg, g \rangle_{\mathcal{G}} \end{aligned}$$

then replacing g with a suitable $e^{i\theta}g$, we get

$$\langle Lf, f \rangle_{\mathcal{H}} - 2|\langle Kf, g \rangle_{\mathcal{G}}| + \langle Mg, g \rangle_{\mathcal{G}} \geq 0,$$

or

$$|\langle Kf, g \rangle_{\mathcal{G}}| \leq \frac{1}{2} \{ \langle Lf, f \rangle_{\mathcal{H}} + \langle Mg, g \rangle_{\mathcal{G}} \}.$$

Then replacing f and g by λf and $\frac{g}{\lambda}$ respectively, we obtain

$$|\langle Kf, g \rangle_{\mathcal{G}}|^2 \leq \langle Lf, f \rangle_{\mathcal{H}} \langle Mg, g \rangle_{\mathcal{G}}.$$

□

Theorem A.4.12 ([1]). *Let L be a continuous positive definite operator. Then there exists uniquely a positive definite operator called the square root of L , denoted by $L^{1/2}$, such that $(L^{1/2})^2 = L$.*

Definition A.4.13. For a continuous linear operator L , the square root of the positive definite operator L^*L is called the *modulus (operator)* of L .

Definition A.4.14. A linear operator V between Hilbert spaces \mathcal{H} and \mathcal{G} is called *isometric* or an *isometry* if it preserves the norm, that is,

$$\|Vf\|_{\mathcal{G}} = \|f\|_{\mathcal{H}} \quad \text{for } f \in \mathcal{H}. \quad (\text{A.36})$$

By (A.36), a continuous linear operator V is isometric if and only if $V^*V = I_{\mathcal{H}}$, in other words, V preserves inner product :

$$\langle Vf, Vg \rangle_{\mathcal{G}} = \langle f, g \rangle_{\mathcal{H}} \quad \text{for } f, g \in \mathcal{H}. \quad (\text{A.37})$$

Definition A.4.15. A linear operator $U : \mathcal{H} \longrightarrow \mathcal{H}$ that is a surjective isometry is called a *unitary (operator)*.

If $U \in B(H)$ is a unitary operator, then $U^* = U^{-1}$.

Definition A.4.16. A continuous linear operator U between Hilbert spaces \mathcal{H} and \mathcal{G} is called a *partial isometry* if for $f \in (\ker U)^\perp = \text{Ran}(U^*)$, $\|Uf\| = \|f\|$. The space $(\ker U)^\perp$ is called the *initial space* of U and the space $\text{Ran}(U)$ is called the *final space* of U .

For a partial isometry U , its adjoint U^* is again a partial isometry.

Corollary A.4.17 ([1]). *Each continuous linear operator L on \mathcal{H} admits a unique decomposition*

$$L = U\tilde{L}, \tag{A.38}$$

where \tilde{L} is positive definite and U is a partial isometry with initial space the closure of $\text{Ran}(\tilde{L})$. Indeed, \tilde{L} must be the modulus $|L|$.

Bibliography

- [1] T. ANDO, *Reproducing Kernel Spaces and Quadratic Inequalities*, Lecture Notes, Hokkaido University, Research Institute of Applied Electricity, Division of Applied Mathematics, Sapporo, Japan, 1987.
- [2] N. ARONSZAJN, *Theory of reproducing kernels*, Trans. Amer. Math. Soc. **68**(1950), 337–404.
- [3] J.B. CONWAY, *A Course in Functional Analysis*, Springer Verlag, Berlin - Heidelberg - New York, 1989.
- [4] P.L. DUREN, *Theory of H_p Spaces*, Academic Press, INC., New York, 1970.
- [5] P.L. DUREN, A. SCHUSTER, *Bergman Spaces*, Amer. Math. Soc., Providence R.I. 2004.
- [6] H. DYM, *J -Contractive Matrix Functions, Reproducing Kernel Hilbert Spaces and Interpolation*, CBMS Regional Conference Series in Mathematics, Amer. Math. Soc., Providence R.I. 1989.
- [7] P. KOOSIS, *Introduction to H_p Spaces*, Cambridge Mathematical Press, Cambridge, 1970.
- [8] S. SAITOH, *Theory of Reproducing Kernels and its Applications*, Pitman Research Notes in Mathematics Series, vol. 189, Longman Scientific and Technical, Harlow, 1988.
- [9] J. WEIDMAN, *Linear Operators in Hilbert Spaces*, Springer Verlag, Berlin - Heidelberg - New York, 1980.