

# Almost periodic solutions of retarded SICNNs with functional response on piecewise constant argument

Marat Akhmet<sup>1</sup> · Mehmet Onur Fen<sup>2</sup> · Mokhtar Kirane<sup>3</sup>

Received: 3 January 2015 / Accepted: 9 August 2015  
© The Natural Computing Applications Forum 2015

**Abstract** We consider a new model for shunting inhibitory cellular neural networks, retarded functional differential equations with piecewise constant argument. The existence and exponential stability of almost periodic solutions are investigated. An illustrative example is provided.

**Keywords** Shunting inhibitory cellular neural networks · Retarded functional differential equations · Alternate constancy of argument · Bohr almost periodic solutions · Exponential stability

## 1 Introduction

Cellular neural networks (*CNNs*) have been paid much attention in the past two decades [1–10]. Exceptional role in psychophysics, speech, perception, robotics, adaptive pattern recognition, vision, and image processing is played by shunting inhibitory cellular neural networks (*SICNNs*), which was introduced by Bouzerdoum and Pinter [11]. One of the most attractive subjects for this type of neural networks is the existence of almost periodic solutions. This

problem has been investigated for models with different types of activation functions [12–20]. In the present study, we investigate a new model of *SICNNs* by considering deviated as well as piecewise constant time arguments and prove the existence of exponentially stable almost periodic solutions. All the results are discussed for the general type of activation functions, but they can be easily specified for applications.

Extended information about differential equations with generalized piecewise constant argument [21] can be found in the book [22]. As a subclass, they contain differential equations with piecewise constant argument (*EPCA*) [23–30], where the piecewise constant argument is assumed to be a multiple of the greatest integer function.

Differential equations with piecewise constant argument are very useful as models for neural networks. This was shown in the studies [31–33], where the authors utilized *EPCA*. We propose to involve a new type of systems, retarded functional differential equations with piecewise constant argument of generalized type, in the modeling. It will help to investigate a larger class of neural networks.

In paper [21], differential equations with piecewise constant argument of generalized type (*EPCAG*) were introduced. We not only maximally generalized the argument functions, but also proposed to reduce investigation of *EPCAG* to integral equations. Due to that innovation, it is now possible to analyze essentially nonlinear systems, that is, systems nonlinear with respect to values of solutions at discrete moments of time, where the argument changes its constancy. Previously, the main and unique method for *EPCA* was reduction to discrete equations and, hence, only equations in which values of solutions at the discrete moments appear linearly [23–30] have been considered.

✉ Marat Akhmet  
marat@metu.edu.tr

<sup>1</sup> Department of Mathematics, Middle East Technical University, 06800 Ankara, Turkey

<sup>2</sup> Neuroscience Institute, Georgia State University, Atlanta, GA 30303, USA

<sup>3</sup> Laboratoire de Mathématiques, Image et Applications, Pôle Sciences et Technologies, Université de La Rochelle, Avenue Michel Crépeau, 17042 La Rochelle, France

The crucial novelty of the present paper is that the piecewise constant argument in the functional differential equations is of alternate (advanced-delayed) type. In the literature, biological reasons for the argument to be delayed were discussed [34, 35]. However, the role of advanced arguments has not been analyzed properly yet. Nevertheless, the importance of anticipation for biology was mentioned by some authors. For example, in the paper [36], it is supposed that synchronization of biological oscillators may request anticipation of counterparts behavior. Consequently, one can assume that equations for neural networks may also need anticipation, which is usually reflected in models by advanced argument. Therefore, the systems taken into account in the present study can be useful in future analyses of *SICNNs*. Furthermore, the idea of involving both advanced and delayed arguments in neural networks can be explained by the existence of retarded and advanced actions in a model of classical electrodynamics [37]. Moreover, mixed-type deviation of the argument may depend on traveling waves emergence in *CNNs* [7]. Understanding the structure of such traveling waves is important due to their potential applications including image processing (see, e.g., [1–7]). More detailed analysis of deviated arguments in neural networks can be found in [38–40].

Shunting inhibition is a phenomenon in which the cell is “clamped” to its resting potential when the reversal potential of  $Cl^-$  channels is close to the membrane resting potential of the cell [11, 41]. It occurs through the opposition of an inward current, which would otherwise depolarize the membrane potential to threshold, by an inward flow of  $Cl^-$  ions [41]. From the biological point of view, shunting inhibition has an important role in the dynamics of neurons [42–44]. According to the results of Vida et al. [42], networks with shunting inhibition are advantageous compared to the networks with hyperpolarizing inhibition such that in the former-type networks, oscillations are generated with smaller tonic excitatory drive, network frequencies are tuned to the  $\gamma$  band, and robustness against heterogeneity in the excitatory drive is markedly improved. It was demonstrated by Mitchell and Silver [43] that shunting inhibition can modulate the gain and offset of the relationship between output firing rate and input frequency in granule cells when excitation and/or inhibition is mediated by time-dependent synaptic input. Besides, Borg-Graham et al. [44] proposed that nonlinear shunting inhibition may act during the initial stage of visual cortical processing, setting the balance between opponent ‘on’ and ‘off’ responses in different locations of the visual receptive field [44]. On the other hand, shunting neural networks are important for various engineering applications [11, 45–53]. For example, in vision, shunting lateral inhibition enhances edges and contrast, mediates directional selectivity, and causes adaptation of the organization of the spatial receptive field and of the contrast

sensitivity function [11, 45–50]. Moreover, such networks are appropriate to be used in medical diagnosis [53]. Therefore, the investigation of the dynamics of *SICNNs*, which are biologically inspired networks designed upon the shunting inhibition concept [11], is important for the improvement in the techniques used in medical diagnosis, adaptive pattern recognition, image processing, etc. [45–53] and may shed light on neuronal activities concerning shunting inhibition [42–44].

Exponential stability of neural networks has been widely studied in the literature (see, e.g., [15–18, 54–60]). According to Liao et al. [54], the exponential stability has importance in neural networks when the exponentially convergence rate is used to determine the speed of neural computations. The studies [54, 55] were concerned with the exponential stability and estimation of exponential convergence rates in neural networks. In the paper [54], Lyapunov–Krasovskii functionals and the linear matrix inequality (LMI) approaches were combined to investigate the problem, whereas the boundedness of the Dini derivative of the neuron input–output activations was required in [55]. The exponential stabilization problem of memristive neural networks was considered in [56] by means of the Lyapunov–Krasovskii functional and free-weighting matrix techniques. Additionally, the Lyapunov–Krasovskii functional method was considered by Wen et al. [61] to analyze the passivity of stochastic impulsive memristor-based piecewise linear systems, and the free-weighting matrix approach was utilized in [57] to derive an LMI-based delay-dependent exponential stability criterion for neural networks with a time-varying delay. On the other hand, exponential stability criteria were derived by Dan et al. [58] for an error system in order to achieve lag synchronization of coupled delayed chaotic neural networks. The concept of lag synchronization was taken into account also within the scope of the papers [62] and [63] for memristive neural networks and for a class of switched neural networks with time-varying delays, respectively. Furthermore, the Banach fixed point theorem and the variant of a certain integral inequality with explicit estimate were used to investigate the global exponential stability of pseudo-almost periodic solutions of *SICNNs* with mixed delays in the study [16].

Almost periodic and in particular quasi-periodic motions are important for the theory of neural networks. According to Pasemann et al. [64], periodic and quasi-periodic solutions have many fundamental importances in biological and artificial systems, as they are associated with central pattern generators, establishing stability properties and bifurcations (leading to the discovery of periodic solutions). Besides, the sinusoidal shape of neural output signals is, in general, associated with appropriate quasi-periodic attractors for discrete-time dynamical systems. In the book [65], the dynamics of the brain activity is

considered as a system of many coupled oscillators with different incommensurable periods. Signals from the neurons have a phase shift of  $\pi/2$  and may be useful for various kinds of applications, for instance controlling the gait of legged robots [66]. Furthermore, an alternative discrete-time model of coupled quasi-periodic and chaotic neural network oscillators was considered by Wang [67].

Let us describe the model of *SICNNs* in its most original form [11]. Consider a two-dimensional grid of processing cells arranged into  $m$  rows and  $n$  columns, and let  $C_{ij}, i = 1, 2, \dots, m, j = 1, 2, \dots, n$ , denote the cell at the  $(i, j)$  position of the lattice. In *SICNNs*, neighboring cells exert mutual inhibitory interactions of the shunting type. The dynamics of a cell  $C_{ij}$  are described by the following nonlinear ordinary differential equation,

$$\frac{dx_{ij}}{dt} = -a_{ij}x_{ij} - \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} f(x_{kl}(t))x_{ij} + L_{ij}(t), \tag{1.1}$$

where  $x_{ij}$  is the activity of the cell  $C_{ij}$ ;  $L_{ij}(t)$  is the external input to the cell  $C_{ij}$ ; the constant  $a_{ij} > 0$  represents the passive decay rate of the cell activity;  $C_{ij}^{kl} \geq 0$  is the coupling strength of postsynaptic activity of the cell  $C_{kl}$  transmitted to the cell  $C_{ij}$ ; the activation function  $f(x_{kl})$  is a positive continuous function representing the output or firing rate of the cell  $C_{kl}$ ; and the  $r$ -neighborhood of the cell  $C_{ij}$  is defined as

$$N_r(i, j) = \{C_{kl} : \max(|k - i|, |l - j|) \leq r, 1 \leq k \leq m, 1 \leq l \leq n\}.$$

It is worth noting that even if the activation function is supposed to be globally bounded and Lipschitzian, these properties are not valid for the nonlinear terms in the right-hand sides of the differential equations describing the dynamics of *SICNNs*, and this is one of the reasons why a sophisticated mathematical analysis is required for *SICNNs* in general. Another reason is that the connections between neurons in *SICNNs* act locally only in  $r$ -neighborhoods. This causes special ways of evaluations different than those customized for earlier developed neural networks in the mathematical analyses of the models.

It is reasonable to say that the usage of deviated arguments in neural networks makes the models much closer to applications. For example, in [13], the model was considered with variable delays,

$$\frac{dx_{ij}}{dt} = -a_{ij}x_{ij} - \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} f(x_{kl}(t - \tau(t)))x_{ij} + L_{ij}(t). \tag{1.2}$$

In the present study, we introduce and investigate more general neural networks. The model will be described in the next section.

## 2 Preliminaries

Let  $\mathbb{Z}$  and  $\mathbb{R}$  denote the sets of all integers and real numbers, respectively. Throughout the paper, the norm  $\|u\| = \max_{(i,j)} |u_{ij}|$ , where  $u = \{u_{ij}\} = (u_{11}, \dots, u_{1n}, \dots, u_{m1}, \dots, u_{mn}) \in \mathbb{R}^{m \times n}$ , will be used.

Suppose that  $\theta = \{\theta_p\}$  and  $\zeta = \{\zeta_p\}$ ,  $p \in \mathbb{Z}$ , are sequences of real numbers such that the first one is strictly ordered,  $|\theta_p| \rightarrow \infty$  as  $|p| \rightarrow \infty$ , and the second one satisfies  $\theta_p \leq \zeta_p \leq \theta_{p+1}$  for all  $p \in \mathbb{Z}$ . The sequence  $\zeta$  is not necessarily strictly ordered. We say that a function is of  $\gamma$ -type and denote it by  $\gamma(t)$ , if  $\gamma(t) = \zeta_p$  for  $\theta_p \leq t < \theta_{p+1}$ ,  $p \in \mathbb{Z}$ . One can affirm, for example, that  $2 \lfloor \frac{t+1}{2} \rfloor$  is a  $\gamma$ -type function with  $\theta_p = 2p - 1$ ,  $\zeta_p = 2p$ .

Fix a nonnegative number  $\tau \in \mathbb{R}$  and let  $\mathcal{C}^0$  be the set of all continuous functions mapping the interval  $[-\tau, 0]$  into  $\mathbb{R}$ , with the uniform norm  $\|\phi\|_0 = \max_{t \in [-\tau, 0]} |\phi(t)|$ . Moreover, we denote by  $\mathcal{C}$  the set consisting of continuous functions mapping the interval  $[-\tau, 0]$  into  $\mathbb{R}^{m \times n}$ , with the uniform norm  $\|\phi\|_0 = \max_{t \in [-\tau, 0]} \|\phi(t)\|$ .

In the present study, we propose to investigate retarded *SICNNs* with functional response on piecewise constant argument of the following form,

$$\frac{dx_{ij}}{dt} = -a_{ij}x_{ij} - \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} f(x_{klt}, x_{kly(t)})x_{ij} + L_{ij}(t), \tag{2.1}$$

where  $f : \mathcal{C}^0 \times \mathcal{C}^0 \rightarrow \mathbb{R}$  is a continuous function.

In network (2.1), the terms  $x_{klt}$  and  $x_{kly(t)}$  must be understood in the way used for functional differential equations [68–70]. That is,  $x_{klt}(s) = x_{kl}(t + s)$  and  $x_{kly(t)}(s) = x_{kl}(\gamma(t) + s)$  for  $s \in [-\tau, 0]$ . Let us clarify that the argument function  $\gamma(t)$  is of the alternate type. Fix an integer  $p$  and consider the function on the interval  $[\theta_p, \theta_{p+1})$ . Then, the function  $\gamma(t)$  is equal to  $\zeta_p$ . If the argument  $t$  satisfies  $\theta_p \leq t < \zeta_p$ , then  $\gamma(t) > t$ , and it is of advanced type. Similarly, if  $\zeta_p < t < \theta_{p+1}$ , then  $\gamma(t) < t$  and, hence, it is of delayed type. Consequently, it is worth noting that the *SICNN* (2.1) is with *alternate constancy* of argument. It is known that  $\gamma(t)$  is the most general among piecewise constant argument functions [22]. Our model is much more general than the equations investigated in [71–74], where the delay is constant  $\tau = 1$ , and it is equal to the step of the greatest integer function  $\lceil t \rceil$ . Differential equations with functional response on the piecewise constant argument were first introduced in the paper [75]. In the present study, we apply the theory to the analysis of neural networks. All previous authors were at most busy with terms of the form  $x(\gamma(t))$ . Thus, one can say that retarded functional differential equations with piecewise constant argument in the most general form are investigated in this paper.

Since the model (2.1) is a new one, we have to investigate not only the existence of almost periodic solutions and their stability, but also common problems of the existence and uniqueness of solutions, their continuation to infinity and boundedness.

One can easily see that system (1.2) is a particular case of (2.1). Additionally, results of the present paper are true or can be easily adapted to the following systems,

$$\frac{dx_{ij}}{dt} = -a_{ij}x_{ij} - \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} f(x_{kl}(\gamma(t)))x_{ij} + L_{ij}(t), \quad (2.2)$$

that is, differential equations with piecewise constant argument, *EPCAG*,

$$\begin{aligned} \frac{dx_{ij}}{dt} = & -a_{ij}x_{ij} - \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} f(x_{kl}(t - \tau(t)), x_{kl}(\gamma(t)) \\ & - \tau(t))x_{ij} + L_{ij}(t), \end{aligned} \quad (2.3)$$

differential equations with variable delay and piecewise constant argument,

$$\begin{aligned} \frac{dx_{ij}}{dt} = & -a_{ij}x_{ij} - \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} f(x_{kl}(t - \tau(t)))x_{ij} \\ & - \sum_{C_{kl} \in N_r(i,j)} D_{ij}^{kl} g(x_{kl}(\gamma(t)))x_{ij} + L_{ij}(t). \end{aligned} \quad (2.4)$$

In other words, what we have suggested are sufficiently general models, which can be easily specified for concrete applications.

Let us introduce the initial condition for *SICNN* (2.1). Fix a number  $\sigma \in \mathbb{R}$  and functions  $\phi = \{\phi_{ij}\}, \psi = \{\psi_{ij}\} \in \mathcal{C}, i = 1, 2, \dots, m, j = 1, 2, \dots, n$ . In the case  $\gamma(\sigma) < \sigma$ , we say that a solution  $x(t) = \{x_{ij}(t)\}$  of (2.1) satisfies the initial condition and write  $x(t) = x(t, \sigma, \phi, \psi), t \geq \sigma$ , if  $x_\sigma(s) = \phi(s), x_{\gamma(\sigma)}(s) = \psi(s)$  for  $s \in [-\tau, 0]$ . In what follows, we assume that if the set  $[\gamma(\sigma) - \tau, \gamma(\sigma)] \cup [\sigma - \tau, \sigma]$  is connected, then the equation  $\phi(s) = \psi(s + \sigma - \gamma(\sigma))$  is true for all  $s \in [-\tau, \gamma(\sigma) - \sigma]$ . If  $\gamma(\sigma) \geq \sigma$ , then we look for a solution  $x(t) = x(t, \sigma, \phi), t \geq \sigma$ , such that  $x_\sigma(s) = \phi(s), s \in [-\tau, 0]$ . Thus, if  $\theta_p \leq \sigma < \theta_{p+1}$  for some  $p \in \mathbb{Z}$ , then there are two cases of the initial condition:

- (IC<sub>1</sub>)  $x_\sigma(s) = \phi(s), \phi \in \mathcal{C}, s \in [-\tau, 0]$  if  $\theta_p \leq \sigma \leq \zeta_p < \theta_{p+1}$ ;
- (IC<sub>2</sub>)  $x_\sigma(s) = \phi(s), x_{\gamma(\sigma)}(s) = \psi(s), \phi, \psi \in \mathcal{C}, s \in [-\tau, 0]$ , if  $\theta_p \leq \zeta_p < \sigma < \theta_{p+1}$ .

Considering *SICNN* (2.1) with these conditions, we shall say about the initial value problem (*IVP*) for (2.1). To be short, we shall say only about *IVP* in the form  $x(t, \sigma, \phi, \psi)$ , specifying  $x(t, \sigma, \phi)$  for (*IC*<sub>1</sub>), if needed. Thus, we can provide the following definitions now.

**Definition 2.1** A function  $x(t) = \{x_{ij}(t)\}, i = 1, 2, \dots, m, j = 1, 2, \dots, n$ , is a solution of (2.1) with (*IC*<sub>1</sub>) or (*IC*<sub>2</sub>) on an interval  $[\sigma, \sigma + a)$  if:

- (i) it satisfies the initial condition;
- (ii)  $x(t)$  is continuous on  $[\sigma, \sigma + a)$ ;
- (iii) the derivative  $x'(t)$  exists for  $t \geq \sigma$  with the possible exception of the points  $\theta_p$ , where one-sided derivatives exist;
- (iv) equation (2.1) is satisfied by  $x(t)$  for all  $t > \sigma$  except possibly at the points of  $\theta$ , and it holds for the right derivative of  $x(t)$  at the points  $\theta_p$ .

**Definition 2.2** A function  $x(t) = \{x_{ij}(t)\}, i = 1, 2, \dots, m, j = 1, 2, \dots, n$ , is a solution of (2.1) on  $\mathbb{R}$  if:

- (i)  $x(t)$  is continuous;
- (ii) the derivative  $x'(t)$  exists for all  $t \in \mathbb{R}$  with the possible exception of the points  $\theta_p, p \in \mathbb{Z}$ , where one-sided derivatives exist;
- (iv) equation (2.1) is satisfied by  $x(t)$  for all  $t \in \mathbb{R}$  except at the points of  $\theta$ , and it holds for the right derivative of  $x(t)$  at the points  $\theta_p, p \in \mathbb{Z}$ .

The existence and uniqueness of solutions of (2.1) will be investigated in the next section.

### 3 Existence and uniqueness

Throughout the paper, we suppose in *SICNN* (2.1) that  $\gamma_0 = \min_{(i,j)} a_{ij} > 0$  and  $C_{ij}^{kl}$  are nonnegative numbers.

The following assumptions are required.

- (C1) The functional  $f$  satisfies the Lipschitz condition  $\|f(\phi_1, \psi_1) - f(\phi_2, \psi_2)\| \leq L(\|\phi_1 - \phi_2\|_0 + \|\psi_1 - \psi_2\|_0)$ , for some positive constant  $L$ , where  $(\phi_1, \psi_1)$  and  $(\phi_2, \psi_2)$  are from  $\mathcal{C}^0 \times \mathcal{C}^0$ ;
- (C2) There exists a positive number  $M$  such that  $\sup_{(\phi, \psi) \in \mathcal{C}^0 \times \mathcal{C}^0} |f(\phi, \psi)| \leq M$ ;
- (C3) There exists a positive number  $\bar{\theta}$  such that  $\theta_{p+1} - \theta_p \leq \bar{\theta}$  for all  $p \in \mathbb{Z}$ ;
- (C4)  $|L_{ij}(t)| \leq L_{ij}$  for all  $i, j$  and  $t \in \mathbb{R}$ , where  $L_{ij}$  are nonnegative real constants.

In the remaining parts of the paper, the notations  $\mu = \max_{(i,j)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl}, \bar{c} = \max_{(i,j)} \frac{\sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl}}{a_{ij}}, \bar{d} = \max_{(i,j)} \frac{\sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl}}{2a_{ij} - \gamma_0}, \bar{L} = \max_{(i,j)} L_{ij}$  and  $\bar{l} = \max_{(i,j)} \frac{L_{ij}}{a_{ij}}$  will be used. We assume that  $\mu\bar{\theta}M < 1$  and  $M\bar{c} < 1$ .

Let us denote  $\mathcal{C}_{H_0} = \{\phi \in \mathcal{C} : \|\phi\|_0 \leq H_0\}$ , where  $H_0$  is a positive number.

**Lemma 3.1** Suppose that the conditions (C1)–(C4) hold and fix an integer  $p$ . If  $H_0$  is a positive number such that  $\mu\bar{\theta}\left[M + \frac{2L(H_0 + \bar{\theta}L)}{1 - \mu\bar{\theta}M}\right] < 1$ , then for every  $(\sigma, \phi, \psi) \in [\theta_p, \theta_{p+1}] \times \mathcal{C}_{H_0} \times \mathcal{C}_{H_0}$ , there exists a unique solution  $x(t) = x(t, \sigma, \phi, \psi)$  of (2.1) on  $[\sigma, \theta_{p+1}]$ .

*Proof* We assume without the loss of generality that  $\theta_p \leq \sigma \leq \zeta_p < \theta_{p+1}$ . That is, we consider (IC<sub>1</sub>) and the solution  $x(t, \sigma, \phi)$ .

Fix an arbitrary function  $\phi \in \mathcal{C}_{H_0}$ . Let us denote by  $\mathcal{A}$  the set of continuous functions  $u(t) = \{u_{ij}(t)\}$ ,  $i = 1, 2, \dots, m, j = 1, 2, \dots, n$ , defined on  $[\sigma - \tau, \theta_{p+1}]$  such that  $u_\sigma(t) = \phi(t)$ ,  $t \in [-\tau, 0]$ , and  $\|u\|_1 \leq K_0$ , where  $\|u\|_1 = \max_{t \in [\sigma, \theta_{p+1}]} \|u(t)\|$  and  $K_0 = \frac{H_0 + \bar{\theta}L}{1 - \mu\bar{\theta}M}$ .

Define on  $\mathcal{A}$  an operator  $\Phi$  such that

$$(\Phi u(t))_{ij} = \begin{cases} \phi_{ij}(t - \sigma), & t \in [\sigma - \tau, \sigma], \\ e^{-a_{ij}(t-\sigma)}\phi_{ij}(0) - \int_\sigma^t e^{-a_{ij}(t-s)} \left[ \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} \right. \\ \left. \times f(u_{kls}, u_{kl\gamma(s)})u_{ij}(s) - L_{ij}(s) \right] ds, & t \in [\sigma, \theta_{p+1}]. \end{cases}$$

One can confirm that  $\|(\Phi u(t))_{ij}\| \leq H_0 + (MK_0 \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} + \bar{L})\bar{\theta}$ ,  $t \in [\sigma, \theta_{p+1}]$ . Accordingly, the inequality  $\|\Phi u\|_1 \leq H_0 + (\mu MK_0 + \bar{L})\bar{\theta} = K_0$  is valid. Therefore,  $\Phi(\mathcal{A}) \subseteq \mathcal{A}$ .

On the other hand, if  $u(t) = \{u_{ij}(t)\}$  and  $v(t) = \{v_{ij}(t)\}$  belong to  $\mathcal{A}$ , then we have for  $t \in [\sigma, \theta_{p+1}]$  that

$$\begin{aligned} & \left| (\Phi u(t))_{ij} - (\Phi v(t))_{ij} \right| \\ & \leq \int_\sigma^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} |f(u_{kls}, u_{kl\gamma(s)}) - f(v_{kls}, v_{kl\gamma(s)})| |u_{ij}(s) - v_{ij}(s)| ds \\ & + \int_\sigma^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} |f(u_{kls}, u_{kl\gamma(s)}) - f(v_{kls}, v_{kl\gamma(s)})| |v_{ij}(s)| ds \\ & \leq \bar{\theta}(M + 2K_0L) \|u - v\|_1 \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl}. \end{aligned}$$

Hence, the inequality  $\|\Phi u - \Phi v\|_1 \leq \mu\bar{\theta}(M + 2K_0L) \|u - v\|_1$  holds. Because  $\mu\bar{\theta}(M + 2K_0L) = \mu\bar{\theta}\left[M + \frac{2L(H_0 + \bar{\theta}L)}{1 - \mu\bar{\theta}M}\right] < 1$ , the operator  $\Phi$  is a contraction. Consequently, there exists a unique solution of (2.1) on  $[\sigma, \theta_{p+1}]$ .  $\square$

The next assertion can be proved exactly in the way that is used to verify Lemma 2.2 from [22], if we use Lemma 3.1.

**Lemma 3.2** Suppose that the conditions (C1)–(C4) hold and fix an integer  $p$ . If  $H_0$  is a positive number such that  $\mu\bar{\theta}\left[M + \frac{2L(H_0 + \bar{\theta}L)}{1 - \mu\bar{\theta}M}\right] < 1$ , then for every  $(\sigma, \phi, \psi) \in [\theta_p, \theta_{p+1}] \times \mathcal{C}_{H_0} \times \mathcal{C}_{H_0}$ , there exists a unique solution  $x(t) =$

$x(t, \sigma, \phi, \psi)$ ,  $t \geq \sigma$ , of (2.1), and it satisfies the integral equation

$$x_{ij}(t) = e^{-a_{ij}(t-\sigma)}\phi_{ij}(0) - \int_\sigma^t e^{-a_{ij}(t-s)} \left[ \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} f(x_{kls}, x_{kl\gamma(s)})x_{ij}(s) - L_{ij}(s) \right] ds. \tag{3.1}$$

### 4 Bounded solutions

In this section, we will investigate the existence of a unique bounded solution of SICNN (2.1). Moreover, the exponential stability of the bounded solution will be considered. An auxiliary result is presented in the following lemma.

**Lemma 4.1** Assume that the conditions (C1)–(C4) are fulfilled. If  $H_0$  is a positive number such that  $\mu\bar{\theta}\left[M + \frac{2L(H_0 + \bar{\theta}L)}{1 - \mu\bar{\theta}M}\right] < 1$ , then a function  $x(t) = \{x_{ij}(t)\}$ ,  $i = 1, 2, \dots, m, j = 1, 2, \dots, n$ , satisfying  $\sup_{t \in \mathbb{R}} \|x(t)\| \leq H_0$  is a solution of (2.1) if and only if it satisfies the following integral equation

$$x_{ij}(t) = - \int_{-\infty}^t e^{-a_{ij}(t-s)} \left[ \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} f(x_{kls}, x_{kl\gamma(s)})x_{ij}(s) - L_{ij}(s) \right] ds. \tag{4.1}$$

*Proof* We consider only sufficiency. The necessity can be proved by using (3.1) in a very similar way to the ordinary differential equations case. One can obtain that

$$\begin{aligned} & \left| \int_{-\infty}^t e^{-a_{ij}(t-s)} \left[ \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} f(x_{kls}, x_{kl\gamma(s)})x_{ij}(s) - L_{ij}(s) \right] ds \right| \\ & \leq \frac{1}{a_{ij}} \left( MH_0 \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} + L_{ij} \right). \end{aligned}$$

Therefore, the integral in (4.1) is convergent. Differentiate (4.1) to verify that it is a solution of (2.1).  $\square$

The following conditions are needed.

- (C5)  $\mu\bar{\theta}\left[M + \frac{2L(H + \bar{\theta}L)}{1 - \mu\bar{\theta}M}\right] < 1$ , where  $H = \frac{\bar{L}}{1 - M\bar{c}}$ ;
- (C6)  $(M + 2LH)\bar{c} < 1$ ;
- (C7)  $2\bar{d}\left[M + LH e^{\gamma_0\tau/2} \left(1 + e^{\gamma_0\bar{\theta}/2}\right)\right] < 1$ .

The main result concerning the existence and exponential stability of bounded solutions of (2.1) is mentioned in the next theorem.

**Theorem 4.1** Suppose that the conditions (C1)–(C6) hold. Then, (2.1) admits a unique bounded on  $\mathbb{R}$  solution, which satisfies (4.1). If, additionally, the condition (C7) is

valid, then the solution is exponentially stable with exponential convergence rate  $\gamma_0/2$ .

*Proof* Let  $C_0(\mathbb{R})$  be the set of uniformly continuous functions defined on  $\mathbb{R}$  such that if  $u(t) \in C_0(\mathbb{R})$ , then  $\|u\|_\infty \leq H$ , where  $\|u\|_\infty = \sup_{t \in \mathbb{R}} \|u(t)\|$ . Define on  $C_0(\mathbb{R})$  the operator  $\Pi$  as

$$(\Pi u(t))_{ij} \equiv - \int_{-\infty}^t e^{-a_{ij}(t-s)} \left[ \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} f(u_{kls}, u_{kl\gamma(s)}) u_{ij}(s) - L_{ij}(s) \right] ds. \tag{4.2}$$

If  $u(t) = \{u_{ij}(t)\}$ ,  $i = 1, 2, \dots, m$ ,  $j = 1, 2, \dots, n$ , belongs to  $C_0(\mathbb{R})$ , then we have that

$$\begin{aligned} |(\Pi u(t))_{ij}| &\leq \int_{-\infty}^t e^{-a_{ij}(t-s)} \left( \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} MH + L_{ij} \right) ds \\ &= \frac{1}{a_{ij}} \left( \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} MH + L_{ij} \right). \end{aligned}$$

Utilizing the last inequality, one can show that  $\|(\Pi u)\|_\infty \leq \bar{c}MH + \bar{l} = H$ . Therefore,  $\Pi u(t) \in C_0(\mathbb{R})$ .

Let us verify that this operator is contractive. Indeed, if  $u(t) = \{u_{ij}(t)\}$  and  $v(t) = \{v_{ij}(t)\}$  belong to  $C_0(\mathbb{R})$ , then

$$\begin{aligned} |(\Pi u(t))_{ij} - (\Pi v(t))_{ij}| &\leq \int_{-\infty}^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} |f(u_{kls}, u_{kl\gamma(s)}) - f(v_{kls}, v_{kl\gamma(s)})| |u_{ij}(s) - v_{ij}(s)| ds \\ &\quad + \int_{-\infty}^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} |f(u_{kls}, u_{kl\gamma(s)}) - f(v_{kls}, v_{kl\gamma(s)})| |v_{ij}(s)| ds \\ &\leq \int_{-\infty}^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} M |u_{ij}(s) - v_{ij}(s)| ds \\ &\quad + \int_{-\infty}^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} HL \left( \|u_{kls} - v_{kls}\|_0 + \|u_{kl\gamma(s)} - v_{kl\gamma(s)}\|_0 \right) ds \\ &\leq (M + 2LH) \frac{\sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl}}{a_{ij}} \|u - v\|_\infty. \end{aligned}$$

Hence, the inequality  $\|\Pi u - \Pi v\|_\infty \leq (M + 2LH)\bar{c}\|u - v\|_\infty$  is valid. In accordance with condition (C6), the operator  $\Pi$  is contractive. Consequently, SICNN (2.1) admits a unique solution  $\tilde{v}(t) = \{\tilde{v}_{ij}(t)\}$  that belongs to  $C_0(\mathbb{R})$ .

We will continue with the investigation of the exponential stability. Fix an arbitrary number  $\epsilon > 0$  and let  $\delta$  be a sufficiently small positive number such that  $\alpha_1 < 1$ ,  $\alpha_2 < 1$ ,  $\alpha_3 < 1$  and  $\mathcal{K}(\delta) < \epsilon$ , where  $\mathcal{K}(\delta) = \frac{\delta}{1 - 2\bar{d}[M + LH e^{\gamma_0/2} (1 + e^{\gamma_0/2})]}$ ,  $\alpha_1 = \mu\bar{\theta} \left[ M + \frac{2L(H + \delta + \bar{\theta}L)}{1 - \mu\bar{\theta}M} \right]$ ,  $\alpha_2 = (M + 2LH)\bar{c} + 4L\bar{d}\mathcal{K}(\delta)$  and  $\alpha_3 = \mu\bar{\theta}(M + 2LH) + 2\mu\bar{\theta}L\mathcal{K}(\delta)$ .

Suppose that  $\tilde{v}_\sigma(s) = \eta(s)$ ,  $s \in [-\tau, 0]$ . Let  $u(t) = \{u_{ij}(t)\}$  be a solution of the network (2.1) with  $u_\sigma(s) = \phi(s)$ ,  $s \in [-\tau, 0]$ , where the function  $\phi$  satisfies the inequality  $\|\phi - \eta\|_0 < \delta$ . Without the loss of generality, we assume that  $\gamma(\sigma) \geq \sigma$ . Using Lemma 3.2, one can verify for  $t \geq \sigma$  that

$$\begin{aligned} u_{ij}(t) - \tilde{v}_{ij}(t) &= e^{-a_{ij}(t-\sigma)} (\phi_{ij}(0) - \eta_{ij}(0)) \\ &\quad - \int_\sigma^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} [f(u_{kls}, u_{kl\gamma(s)}) u_{ij}(s) \\ &\quad - f(\tilde{v}_{kls}, \tilde{v}_{kl\gamma(s)}) \tilde{v}_{ij}(s)] ds. \end{aligned}$$

Denote by  $w(t) = \{w_{ij}(t)\}$ , the difference  $u(t) - \tilde{v}(t)$ . Then,  $w(t)$  satisfies the relation

$$\begin{aligned} w_{ij}(t) &= e^{-a_{ij}(t-\sigma)} (\phi_{ij}(0) - \eta_{ij}(0)) \\ &\quad - \int_\sigma^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} [f(\tilde{v}_{kls} \\ &\quad + w_{kls}, \tilde{v}_{kl\gamma(s)} + w_{kl\gamma(s)}) (\tilde{v}_{ij}(s) + w_{ij}(s)) \\ &\quad - f(\tilde{v}_{kls}, \tilde{v}_{kl\gamma(s)}) \tilde{v}_{ij}(s)] ds. \end{aligned} \tag{4.3}$$

We will consider equation (4.3) with  $\sigma = 0$ . Let  $\Psi_\delta$  be the set of all continuous functions  $w(t) = \{w_{ij}(t)\}$  which are defined on  $[-\tau, \infty)$  such that:

- (i)  $w(t) = \phi(t) - \eta(t)$ ,  $t \in [-\tau, 0]$ ;
- (ii)  $w(t)$  is uniformly continuous on  $[0, +\infty)$ ;
- (iii)  $\|w(t)\| \leq \mathcal{K}(\delta)e^{-\gamma_0 t/2}$  for  $t \geq 0$ .

Define on  $\Psi_\delta$  an operator  $\tilde{\Pi}$  such that

$$(\tilde{\Pi} w(t))_{ij} = \begin{cases} \phi_{ij}(t) - \eta_{ij}(t), & t \in [-\tau, 0], \\ e^{-a_{ij}t} (\phi_{ij}(0) - \eta_{ij}(0)) - \int_0^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} [f(\tilde{v}_{kls} + w_{kls}, \tilde{v}_{kl\gamma(s)} + w_{kl\gamma(s)}) (\tilde{v}_{ij}(s) + w_{ij}(s)) - f(\tilde{v}_{kls}, \tilde{v}_{kl\gamma(s)}) \tilde{v}_{ij}(s)] ds, & t > 0. \end{cases}$$

We shall show that  $\tilde{\Pi} : \Psi_\delta \rightarrow \Psi_\delta$ . Indeed, it is true for  $t \geq 0$  that

$$\begin{aligned}
 |(\tilde{I}w(t))_{ij}| &\leq e^{-a_{ij}t} \delta + \int_0^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} \\
 &\times |f(\tilde{v}_{kls} + w_{kls}, \tilde{v}_{kl\gamma(s)} + w_{kl\gamma(s)}) \\
 &- f(\tilde{v}_{kls}, \tilde{v}_{kl\gamma(s)})| |\tilde{v}_{ij}(s)| ds \\
 &+ \int_0^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} |f(\tilde{v}_{kls} + w_{kls}, \tilde{v}_{kl\gamma(s)} \\
 &+ w_{kl\gamma(s)})| |w_{ij}(s)| ds \leq e^{-a_{ij}t} \delta \\
 &+ \int_0^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} HL(\|w_{kls}\|_0 + \|w_{kl\gamma(s)}\|_0) ds \\
 &+ \int_0^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} MK(\delta) e^{-\gamma_0 s/2} ds \\
 &\leq e^{-a_{ij}t} \delta + \int_0^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} HLK(\delta) \\
 &\left( e^{\gamma_0 \tau/2} + e^{\gamma_0(\bar{\theta} + \tau)/2} \right) e^{-\gamma_0 s/2} ds \\
 &+ \int_0^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} MK(\delta) e^{-\gamma_0 s/2} ds \\
 &= e^{-a_{ij}t} \delta + \left( \frac{2 \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl}}{2a_{ij} - \gamma_0} \right) K(\delta) [M + LHe^{\gamma_0 \tau/2} \\
 &(1 + e^{\gamma_0 \bar{\theta}/2})] e^{-\gamma_0 t/2}.
 \end{aligned}$$

Thus, the inequality

$$\|\tilde{I}w(t)\| \leq e^{-\gamma_0 t} \delta + 2\bar{d}K(\delta)[M + LHe^{\gamma_0 \tau/2}(1 + e^{\gamma_0 \bar{\theta}/2})] e^{-\gamma_0 t/2} \leq K(\delta) e^{-\gamma_0 t/2}$$

is valid for  $t \geq 0$ .

Now, let  $w^1(t) = \{w_{ij}^1(t)\}$ ,  $w^2(t) = \{w_{ij}^2(t)\}$  be elements of  $\Psi_\delta$ . One can confirm for  $t \geq 0$  that

$$\begin{aligned}
 &|(\tilde{I}w^1(t))_{ij} - (\tilde{I}w^2(t))_{ij}| \\
 &\leq \int_0^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} (|\tilde{v}_{ij}(s)| + |w_{ij}^2(s)|) \\
 &\times |f(\tilde{v}_{kls} + w_{kls}^1, \tilde{v}_{kl\gamma(s)} + w_{kl\gamma(s)}^1) \\
 &- f(\tilde{v}_{kls} + w_{kls}^2, \tilde{v}_{kl\gamma(s)} + w_{kl\gamma(s)}^2)| ds \\
 &+ \int_0^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} |f(\tilde{v}_{kls} \\
 &+ w_{kls}^1, \tilde{v}_{kl\gamma(s)} + w_{kl\gamma(s)}^1) - f(\tilde{v}_{kls} \\
 &+ w_{kls}^2, \tilde{v}_{kl\gamma(s)} + w_{kl\gamma(s)}^2)| |w_{ij}^1(s) - w_{ij}^2(s)| ds \\
 &\leq \int_0^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} L(H + K(\delta)) e^{-\gamma_0 s/2} \\
 &\left( \|w_{kls}^1 - w_{kls}^2\|_0 + \|w_{kl\gamma(s)}^1 - w_{kl\gamma(s)}^2\|_0 \right) ds
 \end{aligned}$$

$$\begin{aligned}
 &+ \int_0^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} M |w_{ij}^1(s) - w_{ij}^2(s)| ds \\
 &\leq (M + 2LH) \sup_{t \geq 0} \|w^1(t) - w^2(t)\| \\
 &\frac{\sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl}}{a_{ij}} (1 - e^{-a_{ij}t}) \\
 &+ 4LK(\delta) \sup_{t \geq 0} \|w^1(t) - w^2(t)\| \\
 &\frac{\sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl}}{2a_{ij} - \gamma_0} (e^{-\gamma_0 t/2} - e^{-a_{ij}t}).
 \end{aligned}$$

Therefore, we have that  $\sup_{t \geq 0} \|\tilde{I}w^1(t) - \tilde{I}w^2(t)\| \leq \alpha_2 \sup_{t \geq 0} \|w^1(t) - w^2(t)\|$ . Since  $\alpha_2 < 1$ , one can conclude by using a contraction mapping argument that there exists a unique fixed point  $\tilde{w}(t) = \{\tilde{w}_{ij}(t)\}$  of the operator  $\tilde{I} : \Psi_\delta \rightarrow \Psi_\delta$ , which is a solution of (4.3).

To complete the proof, we need to show that there does not exist a solution of (4.3) with  $\sigma = 0$  different from  $\tilde{w}(t)$ . Suppose that  $\theta_p \leq 0 < \theta_{p+1}$  for some  $p \in \mathbb{Z}$ . Assume that there exists a solution  $\bar{w}(t) = \{\bar{w}_{ij}(t)\}$  of (4.3) different from  $\tilde{w}(t)$ . Denote by  $z(t) = \{z_{ij}(t)\}$  the difference  $\bar{w}(t) - \tilde{w}(t)$ , and let  $\max_{t \in [0, \theta_{p+1}]} \|z(t)\| = \bar{m}$ . It can be verified for  $t \in [0, \theta_{p+1}]$  that

$$\begin{aligned}
 |z_{ij}(t)| &\leq \int_0^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} |\tilde{v}_{ij}(s) + \tilde{w}_{ij}(s)| \\
 &\times |f(\tilde{v}_{kls} + \bar{w}_{kls}, \tilde{v}_{kl\gamma(s)} + \bar{w}_{kl\gamma(s)}) \\
 &- f(\tilde{v}_{kls} + \tilde{w}_{kls}, \tilde{v}_{kl\gamma(s)} + \tilde{w}_{kl\gamma(s)})| ds \\
 &+ \int_0^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} |f(\tilde{v}_{kls} + \bar{w}_{kls}, \tilde{v}_{kl\gamma(s)} \\
 &+ \bar{w}_{kl\gamma(s)})| |z_{ij}(s)| ds \\
 &\leq \int_0^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} L(H + K(\delta)) \\
 &\left( \|z_{kls}\|_0 + \|z_{kl\gamma(s)}\|_0 \right) ds \\
 &+ \int_0^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} M |z_{ij}(s)| ds \\
 &\leq \bar{\theta} \bar{m} [M + 2L(H + K(\delta))] \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl}.
 \end{aligned}$$

The last inequality yields  $\|z(t)\| \leq \alpha_3 \bar{m}$ . Because  $\alpha_3 < 1$ , we obtain a contradiction. Therefore,  $\bar{w}(t) = \tilde{w}(t)$  for  $t \in [0, \theta_{p+1}]$ . Utilizing induction, one can easily prove the uniqueness for all  $t \geq 0$ .  $\square$

*Remark 4.1* In the proof of Theorem 4.1, we make use of the contraction mapping principle to prove the exponential stability. In the literature, Lyapunov–Krasovskii functionals, LMI technique, free-weighting matrix method, and

differential inequality technique were used to investigate the exponential stability in neural networks [12, 54, 57]. They may also be considered in the future to prove the exponential stability in networks of the form (2.1).

The next section is devoted to the existence as well as the exponential stability of almost periodic solutions of (2.1).

### 5 Almost periodic solutions

Let us denote by  $B_0(\mathbb{R})$  the set of all bounded and continuous functions defined on  $\mathbb{R}$ . For  $g \in B_0(\mathbb{R})$  and  $\alpha \in \mathbb{R}$ , a translation of  $g$  by  $\alpha$  is a function  $Q_\alpha g(t) = g(t + \alpha)$ ,  $t \in \mathbb{R}$ . A number  $\alpha \in \mathbb{R}$  is called an  $\epsilon$ -translation number of a function  $g \in B_0(\mathbb{R})$  if  $\|Q_\alpha g(t) - g(t)\| < \epsilon$  for every  $t \in \mathbb{R}$ . Besides, a set  $S \subset \mathbb{R}$  is said to be relatively dense if there exists a number  $h > 0$  such that  $[\vartheta, \vartheta + h] \cap S \neq \emptyset$  for all  $\vartheta \in \mathbb{R}$ . A function  $g \in B_0(\mathbb{R})$  is said to be almost periodic, if for every positive number  $\epsilon$  there exists a relatively dense set of  $\epsilon$ -translation numbers of  $g$  [76].

On the other hand, an integer  $k_0$  is called an  $\epsilon$ -almost period of a sequence  $\{a_p\}$ ,  $p \in \mathbb{Z}$ , of real numbers if  $|a_{p+k_0} - a_p| < \epsilon$  for any  $p \in \mathbb{Z}$  [77]. Let  $\zeta_p^q = \zeta_{p+q} - \zeta_p$  and  $\theta_p^q = \theta_{p+q} - \theta_p$  for all  $p$  and  $q$ . We call the family of sequences  $\{\zeta_p^q\}$ ,  $q \in \mathbb{Z}$ , equipotentially almost periodic [22, 77, 78] if for an arbitrary positive number  $\epsilon$  there exists a relatively dense set of  $\epsilon$ -almost periods, common for all sequences  $\{\zeta_p^q\}$ ,  $q \in \mathbb{Z}$ .

The following conditions are required.

- (C8) The sequences  $\{\zeta_p^q\}$ ,  $q \in \mathbb{Z}$ , as well as the sequences  $\{\theta_p^q\}$ ,  $q \in \mathbb{Z}$ , are equipotentially almost periodic;
- (C9) There exist positive numbers  $\underline{\theta}$  and  $\underline{\zeta}$  such that  $\theta_{p+1} - \theta_p \geq \underline{\theta}$  and  $\zeta_{p+1} - \zeta_p \geq \underline{\zeta}$  for all  $p \in \mathbb{Z}$ .

It follows from condition (C8) that there exists a positive number  $\bar{\theta}$  such that condition (C3) is valid, and  $|\theta_p|, |\zeta_p| \rightarrow \infty$  as  $|p| \rightarrow \infty$  [22, 77, 78].

The next assertion can be proved by the method of common almost periods developed in [79] (see also [77, 78]).

**Lemma 5.1** [78] Assume that  $L(t) = \{L_{ij}(t)\}$ ,  $i = 1, 2, \dots, m, j = 1, 2, \dots, n$  is almost periodic and the conditions (C8), (C9) are valid. Then, for arbitrary

$\eta > 0, 0 < v < \eta$ , there exist relatively dense sets of real numbers  $\Omega$  and integers  $Q$  such that

- (i)  $\|L(t + \alpha) - L(t)\| < \eta, t \in \mathbb{R}$ ;
- (ii)  $|\zeta_p^q - \alpha| < v, p \in \mathbb{Z}$ ;
- (iii)  $|\theta_p^q - \alpha| < v, p \in \mathbb{Z}, \alpha \in \Omega, q \in Q$ .

The existence and exponential stability of the almost periodic solution of the network (2.1) are mentioned in the following theorem.

**Theorem 5.1** Assume that the conditions (C1), (C2), (C4)–(C6), (C8), and (C9) are fulfilled. Then, the SICNN (2.1) admits a unique almost periodic solution. If, additionally, the condition (C7) is valid, then the solution is exponentially stable with exponential convergence rate  $\gamma_0/2$ .

*Proof* It follows from Theorem 4.1 that (2.1) admits a unique bounded on  $\mathbb{R}$  solution  $u(t) = \{u_{ij}(t)\}, i = 1, 2, \dots, m, j = 1, 2, \dots, m$ , which is exponentially stable provided that the condition (C7) is valid. We will show that it is an almost periodic function.

Consider the operator  $\Pi$  defined by equation (4.2) again. It is sufficient to verify that  $\Pi u(t)$  is almost periodic, if  $u(t)$  is.

Let us denote  $\beta = \max_{(i,j)} \left[ \frac{1}{a_{ij}} + (M + 3LH) \frac{\sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl}}{a_{ij}} + \frac{4LH^2 \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl}}{1 - e^{-a_{ij}\underline{\theta}}} \right]$ . Fix an arbitrary positive number  $\epsilon$ . Because  $\Pi u$  is uniformly continuous, there exists a positive number  $\eta$  satisfying  $\eta < \frac{\epsilon}{5}$  and  $\eta \leq \frac{\epsilon}{3\beta}$  such that if  $|t' - t''| < 4\eta$ , then

$$\|\Pi u(t') - \Pi u(t'')\| < \frac{\epsilon}{3}. \tag{5.1}$$

Next, we take into account a number  $v$  with  $0 < v < \eta$  such that  $\|u(t') - u(t'')\| < \eta$  whenever  $|t' - t''| < v$ , and let  $\alpha$  and  $q$  be numbers as mentioned in Lemma 5.1 such that  $\alpha$  is an  $\eta$ -translation number for  $u(t)$ .

Assume that  $t \in (\theta_{\bar{p}} + \eta, \theta_{\bar{p}+1} - \eta)$  for some  $\bar{p} \in \mathbb{Z}$ . Making use of the equation

$$\begin{aligned} (\Pi u(t + \alpha))_{ij} - (\Pi u(t))_{ij} &= - \int_{-\infty}^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} \\ &\quad \times [f(u_{kl}(s+\alpha), u_{kl\gamma}(s+\alpha))u_{ij}(s + \alpha) - f(u_{kls}, u_{kl\gamma}(s))u_{ij}(s)] ds \\ &\quad + \int_{-\infty}^t e^{-a_{ij}(t-s)} [L_{ij}(s + \alpha) - L_{ij}(s)] ds, \end{aligned}$$

we obtain that

$$\begin{aligned}
 & \left| (\Pi u(t + \alpha))_{ij} - (\Pi u(t))_{ij} \right| \\
 & \leq \int_{-\infty}^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} M |u_{ij}(s + \alpha) - u_{ij}(s)| ds \\
 & + \int_{-\infty}^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} LH \|u_{kl}(s+\alpha) - u_{kl}(s)\|_0 ds \\
 & + \int_{-\infty}^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} LH \|u_{kl\gamma}(s+\alpha) - u_{kl\gamma}(s)\|_0 ds \\
 & + \int_{-\infty}^t e^{-a_{ij}(t-s)} |L_{ij}(s + \alpha) - L_{ij}(s)| ds.
 \end{aligned}
 \tag{5.2}$$

According to Lemma 5.1, (i), the inequality

$$\int_{-\infty}^t e^{-a_{ij}(t-s)} |L_{ij}(s + \alpha) - L_{ij}(s)| ds < \frac{\eta}{a_{ij}}$$

is valid. Moreover, since  $\alpha$  is an  $\eta$ -translation number for  $u(t)$ , one can confirm that

$$\int_{-\infty}^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} M |u_{ij}(s + \alpha) - u_{ij}(s)| ds < \frac{M\eta}{a_{ij}} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl}$$

and

$$\int_{-\infty}^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} LH \|u_{kl}(s+\alpha) - u_{kl}(s)\|_0 ds < \frac{LH\eta}{a_{ij}} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl}.$$

On the other hand, we have

$$\begin{aligned}
 & \int_{-\infty}^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} LH \|u_{kl\gamma}(s+\alpha) - u_{kl\gamma}(s)\|_0 ds \\
 & \leq \int_{\theta_p+\eta}^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} LH \|u_{kl\gamma}(s+\alpha) - u_{kl\gamma}(s)\|_0 ds \\
 & + \sum_{\lambda=0}^{\infty} \int_{\theta_{p-\lambda-1}+\eta}^{\theta_{p-\lambda}-\eta} e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} LH \|u_{kl\gamma}(s+\alpha) - u_{kl\gamma}(s)\|_0 ds \\
 & + \sum_{\lambda=0}^{\infty} \int_{\theta_{p-\lambda}-\eta}^{\theta_{p-\lambda}+\eta} e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} LH \|u_{kl\gamma}(s+\alpha) - u_{kl\gamma}(s)\|_0 ds \\
 & + \int_{-\infty}^{\theta_{p-\lambda}-\eta} e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} LH \|u_{kl\gamma}(s+\alpha) - u_{kl\gamma}(s)\|_0 ds.
 \end{aligned}
 \tag{5.3}$$

For any  $p \in \mathbb{Z}$ , if  $s \in (\theta_p + \eta, \theta_{p+1} - \eta)$ , then one can show by using Lemma 5.1, (iii) that the number  $s + \alpha$  belongs to the interval  $(\theta_{p+q}, \theta_{p+q+1})$  so that

$$\|u_{\gamma(s+\alpha)} - u_{\gamma(s)}\|_0 = \max_{\kappa \in [-\tau, 0]} |u(\kappa + \zeta_{p+q}) - u(\kappa + \zeta_p + \alpha)| < \eta,$$

since  $|(\kappa + \zeta_{p+q}) - (\kappa + \zeta_p + \alpha)| = |\zeta_p^q - \alpha| < \nu$  by Lemma 5.1, (ii). Besides, the inequality

$$\sum_{\lambda=0}^{\infty} \int_{\theta_{p-\lambda}-\eta}^{\theta_{p-\lambda}+\eta} e^{-a_{ij}(t-s)} ds \leq 2\eta \sum_{\lambda=0}^{\infty} e^{-a_{ij}\theta\lambda} = \frac{2\eta}{1 - e^{-a_{ij}\theta}}$$

is valid. Therefore, (5.3) yields

$$\begin{aligned}
 & \int_{-\infty}^t e^{-a_{ij}(t-s)} \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl} LH \|u_{kl\gamma}(s+\alpha) - u_{kl\gamma}(s)\|_0 ds \\
 & < LH\eta \left( \frac{2 \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl}}{a_{ij}} + \frac{4H \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl}}{1 - e^{-a_{ij}\theta}} \right).
 \end{aligned}$$

It can be verified by means of (5.2) that

$$\begin{aligned}
 & \left| (\Pi u(t + \alpha))_{ij} - (\Pi u(t))_{ij} \right| \\
 & < \eta \left[ \frac{1}{a_{ij}} + (M + 3LH) \frac{\sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl}}{a_{ij}} + \frac{4LH^2 \sum_{C_{kl} \in N_r(i,j)} C_{ij}^{kl}}{1 - e^{-a_{ij}\theta}} \right].
 \end{aligned}$$

Hence,

$$\|\Pi u(t + \alpha) - \Pi u(t)\| < \beta\eta \leq \epsilon/3 \tag{5.4}$$

for each  $t$  that belongs to the intervals  $(\theta_p + \eta, \theta_{p+1} - \eta)$ ,  $p \in \mathbb{Z}$ .

The inequality  $\eta < \theta/5$  ensures that  $t + 3\eta \in (\theta_p + \eta, \theta_{p+1} - \eta)$  if  $|t - \theta_p| \leq \eta$ . Now, by means of the inequalities (5.1) and (5.4), we attain for  $|t - \theta_p| \leq \eta$ ,  $p \in \mathbb{Z}$ , that

$$\begin{aligned}
 & \|\Pi u(t + \alpha) - \Pi u(t)\| \leq \|\Pi u(t + \alpha) - \Pi u(t + \alpha + 3\eta)\| \\
 & + \|\Pi u(t + \alpha + 3\eta) - \Pi u(t + 3\eta)\| + \|\Pi u(t + 3\eta) - \Pi u(t)\| \\
 & < \epsilon.
 \end{aligned}$$


The last inequality implies that  $\alpha$  is an  $\epsilon$ -translation number of  $\Pi u(t)$ . Consequently, the SICNN (2.1) admits a unique almost periodic solution.  $\square$

**Remark 5.1** The Bohr definition of almost periodicity is also suitable for the application of Lyapunov functional method and the technique of Young’s inequality [59] to show the existence, uniqueness, and exponential stability of almost periodic solutions in CNNs.

### 6 An example

Consider the sequence  $\theta = \{\theta_p\}$  defined as  $\theta_p = p + \frac{1}{4} |\sin(p) - \cos(p\sqrt{2})|$ ,  $p \in \mathbb{Z}$ . Utilizing the technique provided in [77, 80], one can verify that the sequences  $\{\theta_p^q\}$ ,  $q \in \mathbb{Z}$ , are equipotentially almost periodic. We take the function  $\gamma(t)$  with  $\zeta_p = \theta_p$ . One can confirm that the conditions (C3) and (C9) hold with  $\bar{\theta} = 3/2$  and  $\underline{\theta} = 1/2$ , respectively.

Let us take into account the SICNN



**Fig. 1** The unique almost periodic solution of *SICNN* (6.1)

$$\frac{dx_{ij}}{dt} = -a_{ij}x_{ij} - \sum_{C_{kl} \in N_1(i,j)} C_{ij}^{kl} f(x_{kl}(\gamma(t) - \tau))x_{ij} + L_{ij}(t), \tag{6.1}$$

in which  $i, j = 1, 2, 3, f(s) = \frac{s^2}{2}$  if  $|s| \leq 0.1, f(s) = 0.005$  if  $|s| > 0.1, \tau = 0.3,$

Consider the constant function  $\phi(t) = \{\phi_{ij}(t)\}$  such that  $\phi_{11}(t) = -0.025, \phi_{12}(t) = 0.036, \phi_{13}(t) = -0.014, \phi_{21}(t) = 0.012, \phi_{22}(t) = -0.021, \phi_{23}(t) = 0.042, \phi_{31}(t) = 0.023, \phi_{32}(t) = -0.015, \phi_{33}(t) = 0.012.$  We depict in Figure 1 the solution  $x(t) = \{x_{ij}(t)\}$  of (6.1) with  $x(t) = \phi(t), t \leq \sigma = \theta_0 = \frac{1}{4}.$  Figure 1 supports the result of The-

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} = \begin{pmatrix} 9 & 3 & 5 \\ 6 & 5 & 4 \\ 3 & 12 & 9 \end{pmatrix},$$

$$\begin{pmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{pmatrix} = \begin{pmatrix} 0.08 & 0.01 & 0.02 \\ 0.05 & 0.03 & 0.06 \\ 0.04 & 0.07 & 0.02 \end{pmatrix},$$

$$\begin{pmatrix} L_{11}(t) & L_{12}(t) & L_{13}(t) \\ L_{21}(t) & L_{22}(t) & L_{23}(t) \\ L_{31}(t) & L_{32}(t) & L_{33}(t) \end{pmatrix} = \begin{pmatrix} 0.1 \cos(t) + 0.2 \sin(\sqrt{2}t) & 0.2 \cos(\pi t) + 0.1 \sin(\sqrt{2}t) & 0.15 \cos(2t) - 0.12 \cos(\pi t) \\ 0.15 \cos(3t) - 0.1 \sin(\pi t) & 0.2 \cos(t) - 0.15 \sin(\sqrt{2}t) & 0.1 \sin(t) + 0.2 \cos(\sqrt{3}t) \\ 0.2 \cos(\sqrt{2}t) + 0.14 \sin(\pi t) & 0.2 \cos(\sqrt{2}t) + 0.1 \sin(t) & 0.15 \cos(\sqrt{2}t) - 0.13 \cos(4t) \end{pmatrix}.$$

One can calculate that  $\sum_{C_{kl} \in N_1(1,1)} C_{11}^{kl} = 0.17, \sum_{C_{kl} \in N_1(1,2)} C_{12}^{kl} = 0.25, \sum_{C_{kl} \in N_1(1,3)} C_{13}^{kl} = 0.12, \sum_{C_{kl} \in N_1(2,1)} C_{21}^{kl} = 0.28, \sum_{C_{kl} \in N_1(2,2)} C_{22}^{kl} = 0.38, \sum_{C_{kl} \in N_1(2,3)} C_{23}^{kl} = 0.21, \sum_{C_{kl} \in N_1(3,1)} C_{31}^{kl} = 0.19, \sum_{C_{kl} \in N_1(3,2)} C_{32}^{kl} = 0.27, \sum_{C_{kl} \in N_1(3,3)} C_{33}^{kl} = 0.18.$  The conditions (C5)–(C7) are valid for (6.1) with  $\gamma_0 = 3, \mu = 0.38, \bar{c} = \bar{d} = 0.25/3, M = 0.005, L = 0.1, \bar{L} = 0.35, \bar{l} = 0.34/3.$  According to Theorem 5.1, the network (6.1) has a unique almost periodic solution, which is exponentially stable with the rate of convergence 3/2.

orem 5.1 such that the represented solution converges to the unique almost periodic solution of *SICNN* (6.1).

### 7 Conclusion

In this paper, we investigate the existence as well as the exponential stability of almost periodic solutions in a new model of *SICNNs*. The usage of the functional response on alternate (advanced-delayed) type of piecewise constant

arguments is the main novelty of our study, and it is useful for the investigation of a large class of neural networks. An illustrative example is provided to show the effectiveness of the theoretical results.

Our approach concerning exponential stability may be used in the future to investigate synchronization of chaos in coupled neural networks and control of chaos in large communities of neural networks with piecewise constant argument.

Differential equations with functional response on piecewise constant argument can be applied for the development of other kinds of recurrent networks such as Hopfield and Cohen-Grossberg neural networks [81, 82] and others. This will provide new opportunities for the analysis and applications of neural networks.

**Acknowledgments** The authors wish to express their sincere gratitude to the referees for the helpful criticism and valuable suggestions, which helped to improve the paper significantly. The second author is supported by the 2219 Scholarship Programme of TÜBİTAK, the Scientific and Technological Research Council of Turkey.

## References

- Chua LO (1998) CNN: a paradigm for complexity. World Scientific, Singapore
- Chua LO, Yang L (1988) Cellular neural networks: theory. *IEEE Trans Circuits Syst* 35:1257–1272
- Chua LO, Yang L (1988) Cellular neural networks: applications. *IEEE Trans Circuits Syst* 35:1273–1290
- Chua LO, Roska T (1990) Cellular neural networks with nonlinear and delay-type template elements. In: *Proceedings of IEEE international workshop on cellular neural networks and their applications*, pp 12–25
- Chua LO, Roska T (1992) Cellular neural networks with nonlinear and delay type template elements and non-uniform grids. *Int J Circuit Theory Appl* 20:449–451
- Hsu C-H, Lin S-S, Shen W (1999) Traveling waves in cellular neural networks. *Int J Bifurcat Chaos* 9:1307–1319
- Weng P, Wu J (2003) Deformation of traveling waves in delayed cellular neural networks. *Int J Bifurcat Chaos* 13:797–813
- Zou F, Schwartz S, Nossek J (1990) Cellular neural network design using a learning algorithm. In: *Proceedings of IEEE international workshop on cellular neural networks and their applications*, pp 73–81
- Li L, Fang Z, Yang Y (2012) A shunting inhibitory cellular neural network with continuously distributed delays of neutral type. *Nonlinear Anal Real World Appl* 13:1186–1196
- Rosko T, Boros T, Thiran P, Chua LO (1990) Detecting simple motion using cellular neural networks. In: *Proceedings of IEEE international workshop on cellular neural networks and their applications*, pp 127–138
- Bouzerdoum A, Pinter RB (1993) Shunting inhibitory cellular neural networks: derivation and stability analysis. *IEEE Trans Circuits Syst I Fund Theory Appl* 40:215–221
- Chen L, Zhao H (2008) Global stability of almost periodic solution of shunting inhibitory cellular neural networks with variable coefficients. *Chaos Solitons Fractals* 35:351–357
- Ding HS, Liang J, Xiao TJ (2008) Existence of almost periodic solutions for SICNNs with time-varying delays. *Phys Lett A* 372:5411–5416
- Chen A, Cao J (2002) Almost periodic solution of shunting inhibitory CNNs with delays. *Phys Lett A* 298:161–170
- Ou C (2009) Almost periodic solutions for shunting inhibitory cellular neural networks. *Nonlinear Anal Real World Appl* 10:2652–2658
- Cherif F (2012) Existence and global exponential stability of pseudo almost periodic solution for SICNNs with mixed delays. *J Appl Math Comput* 39:235–251
- Hu M, Wang L (2011) Existence and exponential stability of almost periodic solution for Cohen–Grossberg SICNNs with impulses. *World Acad Sci Eng Technol* 52:941–950
- Xia Y, Cao J, Huang Z (2007) Existence and exponential stability of almost periodic solution for shunting inhibitory cellular neural networks with impulses. *Chaos Solitons Fractals* 34:1599–1607
- Huang X, Cao J (2003) Almost periodic solution of shunting inhibitory cellular neural networks with time-varying delay. *Phys Lett A* 314:222–231
- Zhou Q, Xiao B, Yu Y (2006) Existence and stability of almost periodic solutions for shunting inhibitory cellular neural networks with continuously distributed delays. *Electron J Differ Equ* 2006:1–10
- Akhmet M (2006) On the integral manifolds of the differential equations with piecewise constant argument of generalized type. In: Agarwal RP, Perera K (eds) *Proceedings of the conference on differential and difference equations at the Florida Institute of Technology, August 1–5, 2005*. Hindawi Publishing Corporation, Melbourne, pp 11–20
- Akhmet M (2011) *Nonlinear hybrid continuous/discrete time models*. Atlantis Press, Amsterdam
- Aftabizadeh AR, Wiener J (1988) Oscillatory and periodic solutions for systems of two first order linear differential equations with piecewise constant argument. *Appl Anal* 26:327–333
- Busenberg S, Cooke KL (1982) Models of vertically transmitted diseases with sequential-continuous dynamics. In: Lakshmikantham V (ed) *Nonlinear phenomena in mathematical sciences*. Academic Press, New York, pp 179–187
- Cooke KL, Wiener J (1987) Neutral differential equations with piecewise constant argument. *Boll Un Mat Ital* 7:321–346
- Cooke KL, Wiener J (1984) Retarded differential equations with piecewise constant delays. *J Math Anal Appl* 99:265–297
- Dai L (2008) *Nonlinear dynamics of piecewise constant systems and implementation of piecewise constant arguments*. World Scientific, Singapore
- Dai L, Singh MC (1994) On oscillatory motion of spring-mass systems subjected to piecewise constant forces. *J Sound Vib* 173:217–232
- Seifert G (2000) Almost periodic solutions of certain differential equations with piecewise constant delays and almost periodic time dependence. *J Differ Equ* 164:451–458
- Wiener J (1993) *Generalized solutions of functional differential equations*. World Scientific, Singapore
- Zhu H, Huang L (2004) Dynamics of a class of nonlinear discrete-time neural networks. *Comput Math Appl* 48:85–94
- Yang X (2006) Existence and exponential stability of almost periodic solutions for cellular neural networks with piecewise constant argument. *Acta Math Appl Sin* 29:789–800
- Jun YX, Jian WZ (2008) Asymptotic behavior of a neural network model with three piecewise constant arguments. *Hunan Daxue Xuebao* 35:59–62 (Chinese)
- Murray JD (2002) *Mathematical biology: I. An introduction*. Springer, New York

35. Hoppensteadt FC, Peskin CS (1992) *Mathematics in medicine and the life sciences*. Springer, New York
36. Buck J (1988) Synchronous rhythmic flashing of fireflies. II. *Q Rev Biol* 63:265–290
37. Driver RD (1979) Can the future influence the present? *Phys Rev D* 19:1098–1107
38. Akhmet MU, Yılmaz E (2010) Impulsive Hopfield-type neural network system with piecewise constant argument. *Nonlinear Anal Real World Appl* 11:2584–2593
39. Akhmet MU, Aruğaslan D, Yılmaz E (2010) Stability in cellular neural networks with piecewise constant argument. *J Comput Appl Math* 233:2365–2373
40. Akhmet MU, Aruğaslan D, Yılmaz E (2010) Stability analysis of recurrent neural networks with piecewise constant argument of generalized type. *Neural Netw* 23:805–811
41. Shepherd GM (2004) *The synaptic organization of the brain*. Oxford University Press, New York
42. Vida I, Bartos M, Jonas P (2006) Shunting inhibition improves robustness of gamma oscillations in hippocampal interneuron networks by homogenizing firing rates. *Neuron* 49:107–117
43. Mitchell SJ, Silver RA (2003) Shunting inhibition modulates neuronal gain during synaptic excitation. *Neuron* 38:433–445
44. Borg-Graham LJ, Monier C, Frégnac Y (1998) Visual input evokes transient and strong shunting inhibition in visual cortical neurons. *Nature* 393:369–373
45. Bouzerdoum A, Nabet B, Pinter RB (1991) Analysis and analog implementation of directionally sensitive shunting inhibitory cellular neural networks. In: *Visual information processing: from neurons to chips*, vol SPIE-1473, pp 29–38
46. Bouzerdoum A, Pinter RB (1992) Nonlinear lateral inhibition applied to motion detection in the fly visual system. In: Pinter RB, Nabet B (eds) *Nonlinear vision*. CRC Press, Boca Raton, pp 423–450
47. Bouzerdoum A, Pinter RB (1990) A shunting inhibitory motion detector that can account for the functional characteristics of fly motion-sensitive interneurons. *Proc Int Jt Conf Neural Netw* 1:149–153
48. Pinter RB, Olberg RM, Warrant E (1989) Luminance adaptation of preferred object size in identified dragonfly movement detectors. In: *Proceedings of IEEE international conference SMC*, pp 682–686
49. Bouzerdoum A (1993) The elementary movement detection mechanism in insect vision. *Philos Trans R Soc London B* 339:375–384
50. Pinter RB (1983) Product term nonlinear lateral inhibition enhances visual selectivity for small objects and edges. *J Theor Biol* 110:525–531
51. Jernigan ME, Belshaw RJ, McLean GF (1991) Nonlinear lateral inhibition and image processing. In: Nabet B, Pinter RB (eds) *Sensory neural networks: lateral inhibition*. CRC Press, Boca Raton, pp 27–45
52. Fukushima K, Miyake S, Ito T (1983) Neocognitron: a neural network model for a mechanism of visual pattern recognition. *IEEE Trans Syst Man Cybern* 13:826–834
53. Arulampalam G, Bouzerdoum A (2001) Application of shunting inhibitory artificial neural networks to medical diagnosis. *Proceedings of seventh Australian and New Zealand intelligent information systems conference*. Perth, Western Australia, pp 89–94
54. Liao X, Chen G, Sanchez EN (2002) Delay-dependent exponential stability analysis of delayed neural networks: an LMI approach. *Neural Netw* 15:855–866
55. Yi Z, Heng PA, Fu AWC (1999) Estimate of exponential convergence rate and exponential stability for neural networks. *IEEE Trans Neural Netw* 10:1487–1493
56. Wen S, Huang T, Zeng Z, Chen Y, Li P (2015) Circuit design and exponential stabilization of memristive neural networks. *Neural Netw* 63:48–56
57. He Y, Wu M, She J-H (2006) Delay-dependent exponential stability of delayed neural networks with time-varying delay. *IEEE Trans Circuits Syst II Exp Briefs* 53:553–557
58. Dan S, Yang SX, Feng W (2013) Lag synchronization of coupled delayed chaotic neural networks by periodically intermittent control. *Abstr Appl Anal* 2013:501461
59. Jiang H, Zhang L, Teng Z (2005) Existence and global exponential stability of almost periodic solution for cellular neural networks with variable coefficients and time-varying delays. *IEEE Trans Neural Netw* 16:1340–1351
60. Wang L (2010) Existence and global attractivity of almost periodic solutions for delayed high-ordered neural networks. *Neurocomputing* 73:802–808
61. Wen SP, Zeng ZG, Huang TW, Li CJ (2015) Passivity and passification of stochastic impulsive memristor-based piecewise linear system with mixed delays. *Int J Robust Nonlinear Control* 25:610–624
62. Wen S, Zeng Z, Huang T, Zhang Y (2014) Exponential adaptive lag synchronization of memristive neural networks via fuzzy method and applications in pseudorandom number generators. *IEEE Trans Fuzzy Syst* 22:1704–1713
63. Wen S, Zeng Z, Huang T, Meng Q, Yao W (2015) Lag synchronization of switched neural networks via neural activation function and applications in image encryption. *IEEE Trans Neural Netw Learn Syst* 26:1493–1502
64. Pasemann F, Hild M, Zahedi K (2003), SO(2)-networks as neural oscillators. In: Mira J, Álvarez JR (eds) *Computational methods in neural modeling, lecture notes in computer science*, vol 2686, pp 144–151
65. Izhikevich EM (2007) *Dynamical systems in neuroscience: the geometry of excitability and bursting*. The MIT Press, Cambridge
66. Kimura H, Akiyama S, Sakurama K (1999) Realization of dynamic walking and running of the quadruped using neural oscillator. *Auton Robots* 7:247–258
67. Wang X (1992) Discrete-time dynamics of coupled quasi-periodic and chaotic neural network oscillators. *Proceedings of international joint conference on neural networks*. Baltimore, Maryland, pp 517–522
68. Burton TA (1985) *Stability and periodic solutions of ordinary and functional differential equations*. Academic Press, Orlando
69. Hale J (1971) *Functional differential equations*. Springer, New York
70. Kuang Y (1993) *Delay differential equations: with applications in population dynamics*. Academic Press, Boston
71. Seifert G (2003) Second-order neutral delay-differential equations with piecewise constant time dependence. *J Math Anal Appl* 281:1–9
72. Seifert G (2003) Almost periodic solutions of certain neutral functional differential equations. *Commun Appl Anal* 7:437–442
73. Wang G (2007) Periodic solutions of a neutral differential equation with piecewise constant arguments. *J Math Anal Appl* 326:736–747
74. Wang L, Yuan R, Zhang CY (2011) A spectrum relation of almost periodic solution of second order scalar functional differential equations with piecewise constant argument. *Acta Math Sin Engl Ser* 27:2275–2284
75. Akhmet MU (2014) Quasilinear retarded differential with functional dependence on piecewise constant argument. *Commun Pure Appl Anal* 13:929–947
76. Corduneanu C (2009) *Almost periodic oscillations and waves*. Springer, New York
77. Samoilenko AM, Perestyuk NA (1995) *Impulsive differential equations*. World Scientific, Singapore

78. Halanay A, Wexler D (1971) Qualitative theory of impulsive systems. Mir, Moscow (Russian)
79. Wexler D (1966) Solutions périodiques et presque-périodiques des systèmes d'équations différentielles linéaires en distributions. *J Differ Equ* 2:12–32
80. Akhmetov MU, Perestyuk NA, Samoilenko AM (1983) Almost-periodic solutions of differential equations with impulse action. *Akad Nauk Ukrain SSR Inst Mat. Preprint* 26:49 (Russian)
81. Hopfield JJ (1984) Neurons with graded response have collective computational properties like those of two-state neurons. *Proc Natl Acad Sci USA* 81:3088–3092
82. Cohen MA, Grossberg S (1993) Absolute stability of global pattern formation and parallel memory storage by competitive neural networks. *IEEE Trans Syst Man Cybern SMC* 13:815–826